

Central Lancashire Online Knowledge (CLoK)

Title	Gender differences in patellar tendon kinetics during running
Туре	Article
URL	https://clok.uclan.ac.uk/12636/
DOI	https://doi.org/10.1080/02640414.2015.1110318
Date	2015
Citation	Sinclair, Jonathan Kenneth and Taylor, Paul John (2015) Gender differences
	in patellar tendon kinetics during running. Journal of Sports Sciences, 33
	(S1). s21-s24. ISSN 0264-0414
Creators	Sinclair, Jonathan Kenneth and Taylor, Paul John

It is advisable to refer to the publisher's version if you intend to cite from the work. https://doi.org/10.1080/02640414.2015.1110318

For information about Research at UCLan please go to http://www.uclan.ac.uk/research/

All outputs in CLoK are protected by Intellectual Property Rights law, including Copyright law. Copyright, IPR and Moral Rights for the works on this site are retained by the individual authors and/or other copyright owners. Terms and conditions for use of this material are defined in the <u>http://clok.uclan.ac.uk/policies/</u> performed maximal contractions $(\times 3)$ of an isokinetic

knee extension/flexion exercise on a Cybex dynam-

AQ4

ometer at velocities of $30^\circ \cdot \text{s}^{-1}$ and $240^\circ \cdot \text{s}^{-1}$. On 90 the second visit, maximal bilateral and unilateral countermovement $(\times 3)$ and drop jump $(\times 3)$ trials were performed on a force platform (Kistler, 1000 Hz) with lower-body kinematics being concurrently recorded in the frontal plane using a high-speed 95 video camera (Fastec TS3-39, 100 Hz). Independent t-tests examined gender differences in performance $(P_1 < 0.05)$ and Pearson's correlations (two-tailed) examined relationships between isokinetic performance and lower-body jumping kinematics. Females 100 $(4.77^{\circ} \pm 14.20^{\circ})$ displayed significantly greater knee valgus on landing from a bilateral drop jump than males $(-15.82^{\circ} \pm 10.49^{\circ})$, $P_{1} < 0.05^{\circ})$. Importantly, knee valgus during the stance phase of the drop jump displayed a significant negative correlation with 105 concentric hamstring-to-quadriceps ratio (H:Q ratio) AQ5 at $30^{\circ}_{1} \cdot s^{-1}_{1}$ ($r = -0.60, P_{1} < 0.05$). There was a tendency for females $(51.83 \pm 6.85\%)$ and $53.83 \pm 6.85\%$) to display lower H:Q ratios than males $(56.83 \pm 55.42\%$ and $58.17 \pm 10.07\%)$ at 30° AQ6 110 \cdot s⁻¹ and 240° \cdot s⁻¹, respectively, however, this was not statistically significant. Furthermore, there were no significant differences in leg dominance between males and females (P > 0.05). Despite the higher knee injury incidence in females, significant differ-115 ences were only observed between males and

females in landing kinematics during jumping tasks. Present findings suggest that knee joint kinematics adopted by females may be partially explained by quadriceps-dominant characteristics. 120 Although the link between knee joint kinematics and ACL injuries requires further investigation,

the present findings highlight that examination of two-dimensional knee joint kinematics during jumping assessments may provide a useful means 125 of identifying potential injury risk factors for coaches working in an applied setting.

> 032. Effects of ankle bracing on ground reaction forces during cricket bowling and cricket-specific sprinting and

agility performance measures 130

> ADAM HAWKEY^{1,2*}, COURTNEY WILKINSON¹, ASHLEY RICHARDSON¹, IAN GIBBS² & GRAHAM ARNOLD²

¹Abertay University; ²University of Dundee

135 *Corresponding author: a.hawkey@abertay.ac.uk @a_hawkey

The ankle joint is the most common injury site in cricket (Fong et al., 2007, Sports Medicine, 37(1), 73-94). While previous research has reported a reduction in injury risk with the application of a 140 semi-rigid ankle brace (Hawkey et al., 2012, Journal of Sports Therapy, 5(1), 33-40), there is limited research into the effects of ankle bracing on ground reaction forces (GRFs) during bowling and on cricket-specific performance. Therefore, the current 145 study was designed to investigate the effects of wearing ankle braces on both GRF and performance in cricket. Following institutional ethics approval, 20 male university cricket players (age 20 \pm 2.5 years, height 1.78 ± 0.2 m, mass 75 ± 5 kg; Mean [SD]) 150 performed three trials, in both braced (Aircast A60) and non-braced conditions, in each of two cricket performance tests: 20 m sprint (simulating a single run between the wicket) and an adapted 505 agility test (imitating a turn at the wicket), in accordance 155 with Hawkey et al. (2009, Journal of Sports Sciences, 27(4), S137). A paired samples t-test reported a statistically, but not practically, significant difference (P < 0.001) between the 30 m sprint times in the braced $(3.33 \pm 0.26 \text{ s})$ and non-braced 160 $(3.32 \pm 0.26 \text{ s})$ conditions. However, no significant difference (P = 0.29) was found between the agility times for the braced $(2.38 \pm 0.1 \text{ s})$ and non-braced $(2.36 \pm 0.1 \text{ s})$ conditions. Of these 20 participants, six bowlers then delivered an over (six balls), with 165 the foot of their supporting leg contacting a force platform during ball release, in both braced (Aircast A60) and non-braced conditions. An independent samples *t*-test revealed no significant difference between the vertical (P = 0.418), anterior-posterior 170 (P = 0.181) and the mediolateral (P = 0.830) GRFs in the braced and non-braced conditions. The results show no reduction in GRF with the application of an ankle brace, possibly due to the linear nature of the bowling action. However, the non-sig-175 nificant effect on cricket-specific performance, which is comparable to the results of previous cricket-based research (Hawkey et al., 2009, Journal of Sports Sciences, 27(4), \$137), suggests that cricketers may benefit from wearing ankle braces to reduce the 180 potential risk of an ankle injury.

033. Gender differences in patellar tendon kinetics during running

JONATHAN SINCLAIR* & PAUL JOHN TAYLOR

190

AQ7 195

- **.** 170
 - 200

205

210

injury site (Robinson and Nee, JOSPT, 37, 232-238). There is currently a paucity of information regarding the influence of gender on the loads experienced by the patellar tendon during running. The aim of the current investigation was, therefore, to determine whether female recreational runners exhibit distinct patellar tendon loading patterns in relation to their male counterparts. Twelve male (age 26.55 ± 4.11 years, height 1.78 ± 0.11 m, mass 77.11 \pm 5.06 kg) and 12 female (age 26.67 ± 5.34 years, height 1.67 ± 0.12 m, mass 63.28 ± 9.75 kg) runners ran over a force platform which operated at 1000 Hz, at 4.0 m . s⁻¹. Ethical approval was granted by the author's institution. Lower limb kinematics were collected using an eight-camera optoelectric motion capture system which operated at 250 Hz. Patellar tendon loads

It has been shown that 19.4-79.3% of all who parti-

cipate in recreational running activities will suffer

from a chronic pathology over the course of 1 year

(Van Gent et al., 2007, British Journal of Sports

Medicine, 41, 469–480). Female runners are known

to be at increased risk from chronic injuries in rela-

tion to males, with the knee being the most common

- were examined using a predictive algorithm, whereby the knee extensor moment was divided by the patellar tendon moment arm (Janssen et al., 2012, <u>Medicine and Science in Sports Exercise</u>, 45, 927–934; Herzog and Read, 1993, fournal of
- *Anatomy*, 182, 213–230). Sex differences in patellar tendon loads were examined statistically using independent samples *t*-tests. The results indicate that peak patellar tendon force (male = 6.49 ± 2.28 and female = 7.03 ± 1.35 BW) and patellar tendon loading rate (male = 92.41 ± 32.51 and female = 111.05 ± 48.58 BW s^{-1}) were significantly
- higher in female runners. On the basis that patellar tendon pathology is considered to be a function of excessive tendon loading, the current study indicates that female runners may be at increased risk of patellar tendon pathologies.

034. Influence of footwear temperature on the kinetics and kinematics of running

JONATHAN SINCLAIR*, SARAH JANE HOBBS & HANNAH SHORE

University of Central Lancashire

235 *Corresponding author: jksinclair@uclan.ac.uk

The most frequently utilised material for running shoe midsoles is a copolymer called ethylene-vinyl acetate. Like most polymers, ethylene-vinyl acetate

exhibits viscoelastic properties (Knauss et al., 2008, Mechanics of Polymers: Viscoelasticity (pp. 49–96), 240 Springen). It has long been established that the mechanical properties of most polymers are highly temperature dependent (Dib et al., 2001, Journal of Sport Medicine, 15, 172–176); at lower temperatures, the materials become less elastic, whereas the oppo-245 site occurs at higher temperatures. As such, it has been proposed that the cushioning characteristics of running shoes may differ in different environmental temperature conditions. The aim of the current investigation was to examine the effects of cooled 250 footwear on the kinetics and kinematics of running in comparison to footwear at normal temperature. Twelve participants (age 21.45 ± 2.98 years, height 1.66 ± 0.06 m, mass (60.87 ± 4.37) ran at 4.0 m · $\mathbf{s}^{-1} \pm 5\%$ in both cooled and normal temperature 255 footwear conditions over a force platform (1000 Hz). Ethical approval was granted by the author's institution. Two identical footwear were worn, one of which was cooled for 30 min. Lower extremity kinematics were obtained using a motion capture system 260 (250 Hz), and tibial accelerations (1000 Hz) were measured using а tri-axial accelerometer. Differences between cooled and normal footwear temperatures were contrasted using paired samples *t*-tests. The results showed that midsole temperature 265 (P = 0.004) and deformation (P = 0.001) were significantly reduced in the cooled footwear. In addition, instantaneous loading rate (P = 0.02), peak tibial acceleration (P = 0.01) and tibial acceleration slope (IP = 0.007) were significantly greater in the 270 cooled footwear. Finally, peak eversion (P = 0.02) and tibial internal rotation (P = 0.01) were also shown to be significantly larger in the cooled footwear condition. This study indicates that running in cooler footwear places runners at greater risk from 275 the kinetic and kinematic parameters linked to the aetiology of injuries.

035. The effect of a real-time gaitretraining programme on knee angle and ground reaction forces in a group of recreational runners

LOULIA HADJIIOANNOU, ANDREW BARNES*, SEAN CLARKSON & JONATHAN WHEAT

Sheffield Hallam University *Corresponding author: a.barnes@shu.ac.uk

Gait-retraining using real-time visual feedback is an effective intervention for modifying factors AQ8

285

280