

Central Lancashire Online Knowledge (CLoK)

Title	Repetitive task training after stroke: A Cochrane review [Abstract no.116]
Type	Article
URL	https://clock.uclan.ac.uk/16742/
DOI	https://doi.org/10.1177/1747493016669275
Date	2016
Citation	Thomas, Lois Helene, Coupe, Jacqueline, McMahon, Naoimh, Connell, Louise, French, Beverley, Harrison, Joanna, Sutton, Chris J, Tishkovskaya, Svetlana, Leathley, Michael John et al (2016) Repetitive task training after stroke: A Cochrane review [Abstract no.116]. International Journal of Stroke, 11(4) (4_suppl). p. 59. ISSN 1747-4930
Creators	Thomas, Lois Helene, Coupe, Jacqueline, McMahon, Naoimh, Connell, Louise, French, Beverley, Harrison, Joanna, Sutton, Chris J, Tishkovskaya, Svetlana, Leathley, Michael John and Watkins, Caroline Leigh

It is advisable to refer to the publisher's version if you intend to cite from the work.
<https://doi.org/10.1177/1747493016669275>

For information about Research at UCLan please go to <http://www.uclan.ac.uk/research/>

All outputs in CLoK are protected by Intellectual Property Rights law, including Copyright law. Copyright, IPR and Moral Rights for the works on this site are retained by the individual authors and/or other copyright owners. Terms and conditions for use of this material are defined in the <http://clock.uclan.ac.uk/policies/>

Repetitive task training after stroke: A Cochrane review

L Thomas, J Coupe, N McMahon, L Connell, B French, J Harrison, C Sutton, S Tishkovskaya, M Leathley and C Watkins

University of Central Lancashire, Preston, UK

Introduction

Repetitive task training involves the active practice of task-specific motor activities. We updated our Cochrane Review published in 2007.

Method

We searched MEDLINE (01/10/2006 – 08/03/2016), EMBASE (01/10/2006 – 07/03/2016) and the Cochrane Stroke Trials Register (04/03/2016). 2 authors independently screened abstracts, extracted data and appraised trials. Quality of evidence within each study and outcome group was determined using the Cochrane Collaboration Risk of Bias Tool (CCRB) and GRADE criteria.

Results

29 trials with 1759 participants were included. Results were statistically significant for arm function (standardised mean difference (SMD) 0.25, 95% CI 0.11 to 0.40), hand function (SMD 0.28, 95% CI 0.12 to 0.44), sitting balance/reach (SMD 0.28, 95% CI 0.01 to 0.55), walking distance (MD 38.80, 95% CI 24.75 to 52.86), walking speed (SMD 0.33, 95% CI 0.18 to 0.49), functional ambulation (SMD 0.26, 95% CI 0.08 to 0.43), sit-to-stand (Standardised effect 0.35, 95% CI 0.13 to 0.56), lower limb functional measures (SMD 0.29, 95% CI 0.10 to 0.48), standing balance/reach (SMD 0.27, 95% CI 0.09 to 0.45) and global motor function (SMD 0.38, 95% CI 0.11 to 0.65). Follow-up measures were significant for both upper and lower limb outcomes up to 6 months post-treatment.

Conclusion

Repetitive task training resulted in improvement in upper and lower limb function; improvements were sustained up to 6 months post-treatment. Further research should focus on the type and amount of training, including measuring the number of repetitions performed.