

Central Lancashire Online Knowledge (CLoK)

Title	Should Consultation Recording Use be a Practice Standard? A Systematic
	Review of the Effectiveness and Implementation of Consultation Recordings
Туре	Article
URL	https://clok.uclan.ac.uk/20867/
DOI	https://doi.org/10.1002/pon.4592
Date	2018
Citation	Rieger, Kendra, Hack, Thomas, Beaver, Kinta and Schofield, Penelope
	(2018) Should Consultation Recording Use be a Practice Standard? A
	Systematic Review of the Effectiveness and Implementation of Consultation
	Recordings. Psycho-Oncology, 27 (4). pp. 1121-1128. ISSN 1057-9249
Creators	Rieger, Kendra, Hack, Thomas, Beaver, Kinta and Schofield, Penelope

It is advisable to refer to the publisher's version if you intend to cite from the work. https://doi.org/10.1002/pon.4592

For information about Research at UCLan please go to http://www.uclan.ac.uk/research/

All outputs in CLoK are protected by Intellectual Property Rights law, including Copyright law. Copyright, IPR and Moral Rights for the works on this site are retained by the individual authors and/or other copyright owners. Terms and conditions for use of this material are defined in the <u>http://clok.uclan.ac.uk/policies/</u> **Psycho-Oncology**



Should Consultation Recording Use be a Practice Standard? A Systematic Review of the Effectiveness and Implementation of Consultation Recordings

Journal:	Psycho-Oncology
Manuscript ID	PON-17-0444.R1
Wiley - Manuscript type:	Review Article
Date Submitted by the Author:	n/a
Complete List of Authors:	Rieger, Kendra; University of Manitoba , College of Nursing, Rady Faculty of Health Sciences Hack, Thomas; St. Boniface General Hospital Research Centre, Psychosocial Oncology and Cancer Control Beaver, Kinta; University of Central Lancashire, School of Nursing & Caring Sciences Schofield , Penelope; Swinburne University of Technology, Department of Psychological Sciences
Keywords:	Cancer, Oncology, Audiovisual aids, Communication, Consultation, Review, Tape recording



http://mc.manuscriptcentral.com/pon

Should Consultation Recording Use be a Practice Standard? A Systematic Review of the

Effectiveness and Implementation of Consultation Recordings

Kendra L. Rieger, PhD

Assistant Professor College of Nursing, Rady Faculty of Health Sciences, University of Manitoba, Canada Helen Glass Centre for Nursing, 89 Curry Place University of Manitoba, Winnipeg, Manitoba, Canada R3T 2N2 Phone: 204.474.7107 Fax: 204.474.7682 Email: Kendra.Rieger@umanitoba.ca

Thomas F. Hack, PhD

Professor

College of Nursing, Rady Faculty of Health Sciences, University of Manitoba, Canada Director, Psychosocial Oncology & Cancer Nursing Research, I.H. Asper Clinical Research Institute

Senior Scientist, Research Institute in Oncology and Hematology at CancerCare Manitoba Visiting Professor, University of Central Lancashire, Preston, UK

Kinta Beaver, PhD

Professor of Cancer Care School of Health Sciences, University of Central Lancashire, United Kingdom

Penelope Schofield, PhD

Professor, Department of Psychology, School of Health Sciences, Faculty of Health, Arts and Design, Swinburne University of Technology, Hawthorn, Australia.

Department of Cancer Experiences Research, Peter MacCallum Cancer Centre, Melbourne, Australia

Sir Peter MacCallum Department of Oncology, Faculty of Medicine, Dentistry and Health Sciences, The University of Melbourne, Parkville, Australia

Abstract

Objective: To conduct a systematic review of the effectiveness of consultation recordings, and identify factors contributing to their successful implementation in healthcare settings.

Methods: A systematic review was conducted for quantitative studies examining the effectiveness of consultation recordings in healthcare. Two independent reviewers assessed the relevance and quality of retrieved quantitative studies using standardized criteria. Study findings were examined to determine consultation recording effectiveness and to identify barriers and facilitators to implementation. A supplementary review of qualitative evidence was performed to further explicate implementation factors.

Results: Of the 3,373 articles retrieved in the quantitative search, 26 satisfied the standardized inclusion criteria (12 randomized controlled trials, 1 quasi-experiment, and 13 cross-sectional studies). The majority of patients found consultation recordings beneficial. Statistically significant evidentiary support was found for the beneficial impact of consultation recordings on the following patient reported outcomes: knowledge, perception of being informed, information recall, decision-making factors, anxiety, and depression. Implementation barriers included strength of evidence concerns, patient distress, impact of the recording on consultation quality, clinic procedures, medico-legal issues, and resource costs. Facilitators included comfort with being recorded, clinical champions, legal strategies, efficient recording procedures, and a positive consultation recording experience.

Conclusions: Consultation recordings are valuable to patients and positively associated with patient reported outcomes. Successful integration of consultation recording use into clinical practice requires an administratively supported, systematic approach to addressing implementation factors.

Psycho-Oncology

Systematic Review of Consultation Recordings

Background

Patients commonly seek detailed diagnostic and treatment information to gain a fuller understanding of their disease and its management.^{1,2} Patients can be overwhelmed, however, by the complexity of information provided during healthcare consultations.³ Elevated levels of anxiety may hinder efforts to process and retain information imparted during these consultations.⁴ making it difficult to remember.^{5, 6}

The most recent Cochrane review, conducted almost one decade ago, concluded that consultation recordings in oncology "may benefit most adults with cancer...most patients find them very useful".^{5 p.2} The Cochrane review also identified further questions worth exploring, many of which have been the focus of subsequent research studies that have not been subjected to systematic review. Providing patients with audio-recordings of important consultations can reduce their anxiety, improve recall of information, and enhance their perception that they have been adequately informed about their disease and treatment.⁵⁻⁹ **Recordings** enable greater patient participation in decisions surrounding disease management, reduce treatment decision regret, and facilitate patient satisfaction with care.^{4, 5, 7, 8, 10} Despite the established benefits of **recordings**, use of this intervention in clinical practice is sporadic.

The provision of recordings of key consultations is topical and controversial. While some researchers and patient advocacy groups have argued for its routine use in clinical practice, many clinicians are hesitant to record consultations fearing litigation.¹¹ Many believe the evidence base is mixed and inconclusive, while others believe the resources required to implement the intervention preclude its use in standard care.¹² If implementation barriers are perceived to be strong enough to preclude implementation of an empirically validated intervention, then these barriers need to be remedied. The objectives of the present review were to: 1) Conduct a systematic review of the

effectiveness of **consultation recording** use; and 2) Review the **consultation recording** literature to identify key implementation factors and the best ways to address them.

Methods

Review Objective One

Inclusion and exclusion criteria. To be included, studies had to involve adults or children having a clinical consultation in a healthcare setting, after which the patient or family received a consultation recording (tape or digital). Primary quantitative study designs eligible for inclusion were randomized controlled trials (RCTs), quasi-experimental studies (QE), analytical crosssectional (observational) studies, and descriptive cross-sectional studies. Studies from all countries were considered if they were written in English. We excluded studies that investigated written summaries or video recordings of consultations alone, did not provide the **recording** to the patient, or examined the use of standardized audio-recorded education materials. An a priori systematic review protocol guided this work. Research ethics board approval was not required for this systematic review.

Search strategy, study selection, and analysis. Online databases including MEDLINE, CINAHL, PsychINFO, EMBASE, Web of Science, Sociological Abstracts, and ProQuest Dissertations & Theses were systematically searched from January 1, 2002, to August 30, 2016 (see supporting information Table S1). A grey literature search was conducted of Conference Proceedings Citation Index, Grey Matters, Google Scholar, and relevant websites. Two independent reviewers (KR, TH) screened titles/abstracts of retrieved articles for potential inclusion criteria and one reviewer read the full-text of relevant articles to confirm eligibility. Two independent reviewers (KR, TH) critically assessed eligible RCTs with two widely used quality appraisal tools: the Joanna Briggs Institute (JBI) Critical Appraisal Tool for Randomized Controlled Trials¹³ and the Effective Public Health Practice Project Quality Assessment Tool for Quantitative Studies (EPHPP).¹⁴ Quasi-

Psycho-Oncology

Systematic Review of Consultation Recordings

experimental studies were assessed by two reviewers (KR, TH) with the JBI Critical Appraisal Tool for Quasi-Experimental Studies.¹⁵ To be included in the review, an RCT or quasi-experimental study had to meet key criteria determined by two reviewers (see supporting information Table S2). Disagreements were resolved through discussion between the two reviewers. The PRISMA statement for reporting systematic reviews was utilized to frame the search and study selection process.¹⁶ The heterogeneity of the samples, methods, and outcomes in the reviewed studies precluded a meta-analysis; therefore, the extracted data was analyzed using a narrative synthesis.

Review Objective Two

In addition to the descriptive findings and narrative comments contained within the quantitative papers from the first objective, a systematic review of the qualitative literature was conducted using terms associated with healthcare consultations, consultation recordings, and qualitative research.¹⁷ (see supporting information Table S1) MEDLINE, CINAHL, and the grey literature were searched from January 1, 2002, to February 5, 2017. Two reviewers screened retrieved titles/abstracts for relevance (KR, TH), and all qualitative designs were included. Only data addressing the barriers and facilitators contributing to the successful implementation of **recordings** were extracted and thematically analyzed from the included quantitative and qualitative studies.

Results

Quantitative Studies for Inclusion

The database search for objective one retrieved 3359 articles, and an additional 14 articles were retrieved through searching the grey literature, reference lists, and Scopus using the forward search function (see Figure 1). After removing duplicates (n=1374), 1999 titles/abstracts were examined and 1911 articles were excluded for not meeting the inclusion criteria. The full texts of the remaining articles (n=88) were reviewed in-depth, and a further 61 articles were excluded.

Systematic Review of Consultation Recordings

Reasons for exclusion are provided in Figure 1. After critically appraising the RCT and QE studies, 12 RCTs, 1 QE study, and 13 cross-sectional descriptive (CS) studies were included. One RCT was excluded as it did not meet our quality criteria.¹⁸

Description and Quality Appraisal of Quantitative Studies

The majority of studies (see supporting information Table S3 for study details) examined **consultation recordings** for patients with cancer (n=19).^{1-3, 8, 9, 19-32} Some unique oncology contexts included patients transitioning from curative to palliative care,⁸ parents of children with leukemia having an initial consultation,²⁵ and migrant patients with cancer who requested an interpreter.²² Three studies examined the usefulness of **recordings** for enhancing informed consent.^{4, 20, 21} The remaining seven studies occurred outside of the oncology context.^{4, 10, 33-37} The geographical settings included the United Kingdom, ^{2, 4, 10, 26, 28-30, 33, 34} United States,^{1, 19, 24, 32, 36} Canada,^{9, 21, 27, 31} Australia,^{3, 22, 23, 37} Sweden,²⁰ The Netherlands,⁸ Italy,²⁵ and Denmark.³⁵ Supporting information table S2 summarizes the RCT and QE quality appraisals. On the JBI checklist,¹³ 11 RCTs received 8-13 out of a possible 13 points, and one received 5-7 points. On the EPHPP tool,¹⁴ five RCTs received a strong global rating, five received a moderate rating, and two received a weak rating. The one QE²⁶ study received seven out of nine points on the JBI QE tool.¹⁵

Description of Qualitative Studies

In addition to the descriptive data and narrative comments from the quantitative studies, 12 qualitative studies were included to answer the second review objective. These qualitative studies were identified when screening titles/abstracts for objective one or found in the 92 articles retrieved in the systematic search for objective two. Studies explored patients' experiences/perspectives of **consultation recording**^{26, 31, 34, 36, 38-42} and healthcare professionals' experiences of **consultation recording**^{12, 31, 34, 36, 40, 43, 44} The qualitative studies took place in the United States, ^{36, 38, 40, 43, 44} UK, ^{26, 34, 39, 41, 42} Canada, ³¹ and Australia.¹²

Psycho-Oncology

Systematic Review of Consultation Recordings

Objective One: Effectiveness of Consultation Recordings

Satisfaction. Patients had an overall positive perspective of **consultation recordings**. Sixteen studies (7 RCTs, 9 CS) reported that the majority of participants found **recordings** useful or beneficial, with a range of 55-98% across studies.^{1-3, 8, 10, 20, 22-25, 28-30, 34, 36, 37} However, listening to the **recording** was distressing (3-18%)^{2, 20, 29} or a source of anxiety (19-26.8%)^{3, 25} for small proportions of participants. In one study, participants had a marginally significant preference for **consultation recordings** over standardized audio-recordings.²¹ In another study, 17 (57%) patients preferred to receive both a **recording** and summary letter, 7 preferred the audio-recording alone, and 3 preferred the letter alone, with participants more likely to give the **recording** (69%) to a family member or friend than the letter (50%).³ Five RCTs examined if **recordings** had an impact on the patients' satisfaction with communication with the health provider, but none found significant differences.^{9, 21, 23, 27, 37}

Use of consultation recordings. Most studies reported that the majority of participants (51-100%) listened to their **recordings**,^{2-4, 8-10, 19, 20, 22-25, 27-29, 31-34, 37} although one study found that only 31% of participants listened.³⁵ Some participants listened more than once, with an average use of 1.3 to 3.0 times.^{9, 10, 21, 25-28, 30, 31} Reasons cited for not listening included technological issues,^{26, 31, 35} not wanting to revisit their decision,²⁶ feeling they had received sufficient information already,^{3, 26, ^{29, 31} worrying it would evoke negative emotions,^{20, 29} intending the **recording** for a family member,²⁹ being overwhelmed or too busy,^{31, 34} desiring to move on,^{3, 31} preferring another modality,³¹ feeling too sick,³¹ disliking the sound of their voice,³⁴ and lacking the privacy to listen to the **recording**.³⁴ When patients were prompted to record their own consultations, one study found that only 20% did so.³² Between 20 and 100% of participants in 14 studies reported that they shared their **recording** with others.^{1-3, 8-10, 19, 20, 22, 23, 27, 29, 31, 32} Patients shared their **recording** with spouses/partners, children, family members, friends, and doctors. A number of factors were}

Systematic Review of Consultation Recordings

associated with using the **recording** including increased anxiety,²³ being married,³ receiving bad news,³ having longer consultations,³ increasing age,³⁵ and gender (female patients and their relatives replaying together; relatives of male patients replaying alone).³⁵

Knowledge, perception of being informed, and recall of information. Nine (75.0%) RCTs assessed the impact of **recordings** on knowledge, perception of being informed, or recall of information, and five found significant positive results. Of the four RCTs specifically measuring knowledge, three investigated knowledge relevant to enhancing informed consent^{4, 20, 21} and only one demonstrated significant findings.⁴ Two^{9, 27} of the three^{9, 21, 27} studies examining patients' perception of being informed, and two^{2, 37} of the three^{2, 33, 37} studies measuring recall, showed that the **recording** provided significant benefit.

Psychological health and well-being. Seven (58.3%) RCTs measured anxiety, depression, and/or stress,^{2, 4, 10, 23, 30, 33, 37} and significant differences were observed in three studies. In two studies, **recordings** significantly reduced participants' anxiety^{4, 33} and in one study it reduced their depression.⁴ Another RCT found significant decreases in anxiety and depression when the analysis was repeated with only those who had listened to the recording.²³ There were no other RCTs that reported a statistically significant impact of **recordings** on psychological adjustment,^{9, 27, 30} quality of life,^{8, 9, 27} treatment adherence,¹⁰ or openness to discussing cancer-related issues in the family.⁸

Decision making. Of the three studies assessing the impact of **consultation recordings** on concepts associated with decision making, all had significant findings. One study found significantly higher decisional self-efficacy,³⁰ and two found lower decisional regret,^{26, 30} for those who had received a recording. In the QE study, receiving a **recording** was positively associated with less decision regret, and was even a stronger predictor of decision regret than erectile dysfunction or incontinence.²⁶ Another RCT measured health locus of control, and found that the

Psycho-Oncology

Systematic Review of Consultation Recordings

consultation recording group reported a significantly higher sense of control regarding their own health than the control and standardized recording groups.⁴

Objective Two: Barriers and Facilitators to Consultation Recording

The synthesis for the second question revealed noteworthy individual, interpersonal, and system barriers to implementing **consultation recordings**, as well as critical facilitators.¹²

Perceived value of consultation recording and the evidence. The qualitative findings showed that both patients and practitioners valued **consultation recordings**.^{12, 31, 36, 39, 42} Patients appreciated **recordings** for the purposes of enhancing patient-centred communication and fostering empowerment.³⁹ Some practitioners questioned whether **recordings** benefited those with a passive decision making style.¹² Healthcare professionals had differing assessments of the empirical evidence on **recordings**, with the majority being unaware of the supporting evidence base.^{12, 31} The provision of high quality, accessible evidence such as systematic reviews and clinical practice guidelines was identified as an essential facilitator to ensuring staff were adequately informed, and promoting practice change.³¹

Clinician's motivation. Positive feedback motivated clinicians to continue with **recordings**. Piloting and evaluating **consultation recording** was identified as a means by which to promote the "buy-in" of stakeholders and to inspire routine implementation,¹² through exposure to **the** benefits.^{29, 40, 43} One study offered a feedback letter to oncologists espousing the patient-reported benefits of **recordings** to reinforce the importance of continuing the practice.³¹

Fear of distressing patients. Clearly identified barriers included concerns that **recordings** could increase patients' anxiety when replaying distressing content or impose an undesired active decision-making role on patients.^{2, 12, 29} Practitioners noted that certain visits might not be wholly appropriate for **recording,** such as consultations in which bad news was given, the patient became upset, or the patient was extremely anxious.^{12, 29}

Impact on the consultation. Variation in perspectives regarding the impact of **recording** on the consultation determined whether this factor was a barrier or facilitator. Some healthcare workers reported feeling anxious or self-conscious about what they said when being recorded as well as concerned that the **recording** would make the communication more formal, factual, or structured.^{12, 20, 29} Practitioners noted that consultations are not always a straightforward provision of information and that a desire to create a coherent recording could be a barrier.¹² In contrast, others felt at ease with being recorded⁴⁰ or thought that more detailed information was given with **recordings**.²⁹ Some patients thought the intervention might violate the traditional etiquette of the doctor-patient relationship; thus, challenging the clinicians' status or damaging the relationship.³⁹ Others believed that the **recording** process would not hinder physicians from sharing information, might increase their accountability, and could enhance respect for clinicians who were willing to be recorded.39

Clinical champions. An essential facilitator of successful implementation was leadership. that is, an identified, respected champion with the administrative or social power to advocate for the intervention, and obtain necessary funds and staff resources for a consultation recording program.³¹ Support from physicians conducting the consultations was crucial for successful implementation, as was their consistency in following the recording procedures.⁴⁴

Legal and privacy concerns. There was evidence that legal and privacy concerns were substantial barriers to recordings. Physicians were concerned about medico-legal implications, such as who owns the recording, whether the treatment center should retain a copy, how to store the recordings within existing medical records, if the recording might be used in a lawsuit, and with whom the recording would be shared.^{12, 29, 31} There were also concerns about a breach of confidentiality if the consultation was shared without permission, and a loss of control of the recording if it was posted on the internet.^{1, 34, 39} An important facilitator was assuring that legal

6

Psycho-Oncology

Systematic Review of Consultation Recordings

requirements were met.¹² One strategy involved creating a disclosure/consent form for patients, providers, and the organization, and ensuring the consent process communicated the patients' and providers' rights, obligations, and the appropriate forms of distribution for the **recording**.¹² One center created a disclaimer which clearly stated that the **recording** had not been reviewed by physicians, could contain errors or omissions, and that it was primarily for the convenience and personal use of patients.⁴³

Required resources and technology. There were reservations about the amount of time that recordings might require in an already overloaded clinical setting.^{12, 24, 36, 43} However, researchers reported that recordings did not substantially increase the length of consultations,^{1, 9, 10, 24, 29} and some even found it was an efficient means of promoting the doctor-patient partnership.^{10, 12, 34} Intervention and resource costs were identified as other primary barriers to the sustainable implementation of recordings.^{12, 31, 36, 40} One cost-saving facilitator was integrating recording procedures into existing staff roles.⁴⁰ Technology was highlighted as a facilitator when no significant technical problems occurred,⁸ and as a barrier when malfunctioning or inaccessible equipment hindered the intervention.^{9, 10, 22, 29, 34, 37} Researchers cited poor recording quality as a barrier to use.^{3, 8, 24, 26, 34} With an array of recording options, selecting the most appropriate recording technology for the target patient population was an important consideration.^{34, 41} The growing accessibility of smartphones for self-recording of consultations was acknowledged as holding potential for facilitating recordings.^{12, 39}

Logistics and procedures. Logistical facilitators included **recordings** posing a minimal burden to the clinical environment and embedding **recordings** into usual care.¹² Clear procedures and staff support were critical to preventing problems with **recordings** and avoiding increased consultation time.^{8, 12} Significant logistical components of supporting **recordings** included organized scheduling, informing patients of the benefits of **recordings** and their option to record,

ensuring the recording equipment was functioning properly, obtaining consent to record, ensuring the recording occurred despite time pressures, and delivering the recording to the patient.^{12, 24, 31, 38,} ⁴³ To address concerns of **recordings** causing distress, patients should be offered the choice of receiving a **recording** rather than systematically providing one to all patients.¹² To avoid covert self-recordings, procedures should be in place to allow patients to record their own consultations.^{12, ³⁹}

Conclusions

This systematic review builds upon previous work by including the most recent literature and elucidating the barriers and facilitators of routine implementation. The findings demonstrate that **consultation recordings** enhance decision-making and patient satisfaction, and that patients frequently listen to their recordings and share them with others. Although patients derive benefits from **recordings**, there were equivocal findings regarding knowledge, information recall, the perception of being informed, and psychological well-being.

Only a proportion of the studies reported significant results for anxiety and depression. It may be unreasonable to expect patients' psychological well-being to be significantly improved by a recording only a short time after a life-altering diagnosis.⁹ There was also variation in patients' perceptions of recordings and identified factors that contributed to their use. Recordings may also resonate more with certain individuals depending on their preferred coping mechanisms. If patients prefer to use denial as a coping mechanism, recordings may not be valuable to them, and may actually cause anxiety.⁶

The findings of this present review had notable differences with the earlier Cochrane review⁵ A distinct finding, which was only evident in this review, was that **recordings** had an unequivocally positive impact on concepts associated with decision making. Lower decision regret

Psycho-Oncology

Systematic Review of Consultation Recordings

may be a result of **consultation recordings** enhancing patients' understanding of the risks, benefits, and outcomes of various treatments, all contributing to more fully informed decision making.²⁶

This review also extended the Cochrane review by identifying key implementation factors and strategies to address them. Barriers to implementation of **recordings** included a lack of awareness of the empirical evidence on **recordings**, feeling self-conscious about being recorded, perceptions that the recording process will negatively impact the consultation, variation in physician cooperation with **recording** procedures, lack of staff and financial resources, and medico-legal concerns. Important implementation facilitators included acceptance of research evidence, valuing **recordings** as a communication tool, champions to promote **recordings**, positive feedback for clinicians who use **recordings**, and dedicated staff to support **recordings**. Producing research knowledge alone is often insufficient to change practice. This systematic identification of **consultation recording** implementation factors is vital to establishing and maintaining a **recording** service.⁴⁵

Although none of the reviewed studies were conducted in low-income countries, some of the studies included participants from disadvantaged populations.^{2, 22, 41} **Recordings** may be especially useful for those with low literacy or for non-English speakers, as the intervention does not rely on written information alone.^{2, 22, 41} A patient in one study who spoke English as a second language thought the **recordings** allowed a careful review of sentences not initially understood.³ This potential of **recordings** is important, as literacy is identified as a determinant of health and **recordings** may promote health literacy through a more accessible medium.⁴⁶ Further, one study reported that the commonly observed positive association between deprivation and anxiety was not found in patients who received a **recording**.² Thus, **recordings** may reduce the negative impact of socio-economic factors. These observations are noteworthy as a review of audio-recordings for health literacy found no studies of **recordings** with disadvantaged or low-literacy populations.⁴⁶

Study limitations. First, we included evidence from cross-sectional descriptive studies to capture informative data about the use of, and satisfaction with, **recordings** in diverse settings. This study design is especially vulnerable to the impact of confounding variables and alternative explanations for the results.⁴⁷ Second, there were a number of studies in which **consultation recording** was part of a complex intervention that included other decision or communication aids.^{19, 30, 32, 36} In a complex intervention, it may not be possible to attribute findings to a particular component of an intervention such as **the recording**. Lastly, all of the studies took place in Canada, US, Europe, or Australia. These findings may not be generalizable to countries in South America, Asia or Africa where there may be appreciable differences in healthcare provider-patient communication and the impact of **recordings**.

Implications for research. Although there has been some preliminary work carried out in regard to who benefits most from **recordings** and what types of consultations should be recorded, more research is needed.^{7, 8, 10} The mechanisms by which benefits occur with **recordings** need to be better understood as this knowledge could inform outcome variables and the most appropriate measurement time-points.^{7, 27} More research on the factors associated with valuing and utilizing **recordings** is needed. Continued investigation of **recordings** amongst diverse and disadvantaged populations would be insightful and support broader use of this intervention.⁴⁰ Further study is needed regarding healthcare providers' perspectives of patient self-recording, why more patients do not record their consultations, and how healthcare providers could enable patients to do so.³² Given the **growing body of evidence supporting consultation recordings**, systematic implementation studies targeting the barriers and facilitators identified in this review are also needed.³¹

Clinical implications. The current review provides convincing evidence of the effectiveness of **recordings** in supporting informed decision making and demonstrates that patients value and use **recordings**. Findings also suggest that **recordings** can foster recall of information, the perception

Page 15 of 73

Psycho-Oncology

Systematic Review of Consultation Recordings

of being informed, knowledge, and psychosocial adjustment in some situations. Although there were equivocal findings on certain outcomes, **recordings were** highly valued by patients and **are** not burdensome in regard to time and costs. This strength of the evidence suggests that **the routine implementation of recording** use should be **considered** in oncology, i.e., the clinical area with the largest, most compelling evidence base. The findings of this review also indicate that recordings may be helpful for many patients but not for all. Providing patients with a choice of whether to receive the recording or other communication aids may be important to address personal preferences. Clinicians should also be aware of the potential for distress for some patients and provide adequate psychosocial supports, especially for distressing consultations.

The findings also revealed that despite the positive evidence base for this intervention, several factors impede its uptake into clinical practice. Fears of litigation are a commonly expressed rationale for not recording consultations with patients. Concerns that these recordings may turn up in social media and other public domains are frequent. However, Elwyn and colleagues⁴⁸ argue that liability insurers believe recordings actually protect clinicians, that patients primarily share their recordings with family and caregivers, and that clinicians may be able take legal action against patients who intentionally use a recording to harm their reputation. Although accurate information can allay fears, other implementation hurdles exist. Organizational resource costs must be addressed to enable successful implementation. This review suggests that for implementation to be successful, treatment centers need to be committed to providing a recording service, assign responsibility for the service, ensure recording equipment is available for all consultations, and develop staff education materials among other associated implementation processes and materials. Time is necessary for staff education, and for evaluating and maintaining the service. While senior administrative levels may express concern that there are insufficient resources to support a recording service, it is much less expensive now than in the past. The cost

of recording and memory storage devises have dropped significantly over the past decade, and in some instances patients can record the consultations themselves on their cellular devices, thereby freeing up clinic resources that might otherwise have been used to purchase recording materials. The cost of educating staff in the service, and the cost of recording mediums such as digital recorders and USB memory keys is low when compared against other clinical services.

In the absence of a **recording** service, patients are recording consultations on their own, sometimes covertly, out of concern their clinicians may look upon them with disfavor if they express their intention or desire to have their consultations recorded. In one survey, 15% of respondents reported having covertly recorded a consultation, and 35% admitted that they would consider doing so.³⁹ If increasing numbers of patients are **desiring to** record consultations on their own,^{39, 49-51} should clinical centers establish policies and procedures for addressing this trend? In most western nations, anecdotal legal opinion suggests that the law supports the right of citizens to record personal communications with a second party without second party consent, as both parties engaged in a mutually agreed to conversation are said to "own" the conversation. Moreover, an online public forum in the UK, which attracted legal commentary, concluded that patients did not have to seek permission to record consultations, and that such recordings were legal.¹¹ Additional study of the rights of patients and clinicians surrounding the use of **recordings** is needed to clarify the limit of any legal argument.

Moving forward, decisions should be made at the highest administrative levels as to whether the practice of **recording** is supported, and if so, then further decisions need to be made as to how to establish the service. If clinic staff are responsible for recording consultations, then standardized procedures will be required, including decisions surrounding which consultations to record, what parts of the consultation to record, whether the clinic will retain a copy of the recording, **and**

Psycho-Oncology

Systematic Review of Consultation Recordings

whether to include a disclaimer statement for legal protection. If patients are encouraged to record consultations, then how will patients become aware of the option to record their consultations? Implementation tips have been prepared for those wanting to initiate a **recording** service (see Table 1).⁷ On the other hand, if administration decides not to support the use of **recordings** in practice, then procedures should be established for responding to patients who express a desire to record consultations, keeping in mind that an increasing number of patients are recording consultations covertly, and in most developed countries are likely protected by law in

doing so.

http://mc.manuscriptcentral.com/pon

This work was supported by the St. Boniface Hospital Foundation and a Chair in Psychosocial Oncology and Supportive Care Research awarded to Dr. Thomas Hack from the Canadian Breast Cancer Foundation.

We would like to thank Ms. Lisa Demczuk for her assistance with the search strategy.

Conflicts of interest:

We have no conflicts of interest to declare.

arch.

Psycho-Oncology

Systematic Review of Consultation Recordings

References

- 1. Seider MI, Damato BE. Digital audio recording of initial patient visits to an ocular oncology clinic: a pilot study. *OSLI Retina*. 2015;46(5):597-599. doi: 10.3928/2325816020150521-16
- Stephens MR, Gaskell AL, Gent C, Pellard S, Day-Thompson R, Blackshaw GR, Lewis WG. Prospective randomised clinical trial of providing patients with audiotape recordings of their oesophagogastric cancer consultations. *Patient Educ Couns*. 2008;72(2):218-222. doi: 10.1016/j.pec.2008.04.004
- Knox R, Butow PN, Devine R, Tattersall MH. Audiotapes of oncology consultations: Only for the first consultation? *Ann Oncol.* 2002;13(4):622-627.
- Mishra PK, Mathias H, Millar K, Nagrajan K, Murday A. A randomized controlled trial to assess the effect of audiotaped consultations on the quality of informed consent in cardiac surgery. *Arch Surg* 2010;145(4):383-388. doi: 10.1001/archsurg.2010.45
- Pitkethly M, MacGillivray S, Ryan R. Recordings or summaries of consultations for people with cancer. *Cochrane Database Syst Rev.* 2008;3:1-58. doi:

10.1002/14651858. CD001539. pub2

- 6. McClement SE, Hack TF. Audio-taping the oncology treatment consultation: a literature review. *Patient Educ Couns.* 1999;36(3):229-238.
- Hack TF, Beaver K, Schofield P. Audio-recording cancer consultations for patients and their families – putting evidence into practice. In: Kissane DW, Bultz BD, Butow PN, Bylund CL, Noble S, Wilkinson S eds. *Oxford Textbook of Communication in Oncology and Palliative Care*. 2nd ed. Oxford: Oxford University Press; 2017. p. 56-60
- 8. Uitdehaag MJ, van der Velden L-A, de Boer MF, Spaander MC, Steyerberg EW, Kuipers EJ, et al. Recordings of consultations are beneficial in the transition from curative to palliative cancer

care: a pilot-study in patients with oesophageal or head and neck cancer. *Eur J Oncol Nurs*. 2012;16(2):109-114. doi: 10.1016/j.ejon.2011.04.006

- Hack TF, Pickles T, Bultz BD, Reuther JD, Weir LM, Degner LF, et al. Impact of providing audiotapes of primary adjuvant treatment consultations to women with breast cancer: a multisite, randomized, controlled trial. *J Clin Oncol.* 2003;21(22):4138-4144. doi: 10.1016/j.ejon.2011.04.006
- 10. Liddell C, Rae G, Brown TR, Johnston D, Coates V, Mallett J. Giving patients an audiotape of their GP consultation: a randomised controlled trial. *Br J Gen Pract*. 2004;54(506):667-672.
- 11. Elwyn G. "Patientgate"—digital recordings change everything. *BMJ*. 2014;348:1-2. doi: 10.1136/bmj.g2078
- Moloczij N, Krishnasamy M, Butow P, Hack TF, Stafford L, Jefford M, et al. Barriers and facilitators to the implementation of audio-recordings and question prompt lists in cancer care consultations: a qualitative study. *Patient Educ Couns*. 2017;100(6):1083-1091. doi: 10.1016/j.pec.2017.01.005.
- 13. Joanna Briggs Institute. The Joanna Briggs Institute critical appraisal checklist for randomized controlled trials. Adelaide: Joanna Briggs Institute; 2016. Available from: http://joannabriggs.org/assets/docs/critical-appraisal-tools/JBI_Critical_Appraisal-Checklist for Randomized Controlled Trials.pdf
- 14. Thomas BH, Ciliska D, Dobbins M, Mirucci S. A process for systematically reviewing the literature: providing the research evidence for public health nursing interventions. *Worldviews Evid Based Nurs*. 2004;1(3):176-184. doi: 10.1111/j.1524-475X.2004.04006.x
- 15. Joanna Briggs Institute. The Joanna Briggs institute critical appraisal checklist for quasiexperimental studies. Adelaide: Joanna Briggs Institute; 2016. Available from:

Psycho-Oncology

1	Sy	stematic Review of Consultation Recordings	21
2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 17 18		http://joannabriggs.org/assets/docs/critical-appraisal-tools/JBI_Critical_Appraisal-	
		Checklist_for_QuasiExperimental_Studies.pdf	
	16	. Liberati A, Altman DG, Tetzlaff J, Mulrow C, Gøtzsche PC, Ionnidis JPA, et al. The PRISM	ÍA
		statement for reporting systematic reviews and meta-analyses of studies that evaluate healthc	care
		interventions: explanation and elaboration. BMJ. 2009;339:1-27. doi: 10.1136/bmj.b2700	
	17	. Tricco AC, Antony J, Zarin W, Strifler L, Ghassemi M, Ivory J, et al. A scoping review of ra	apid
		review methods. BMC Medicine. 2015;13:1-15. doi: 10.1186/s12916-015-0465-6	
19 20	18	. Wolderslund M, Kofoed PE, Holst R, Axboe M, Ammentorp J. Digital audio recordings	
21 22		improve the outcomes of patient consultations: A randomised cluster trial. Patient Educ Cou	INS.
23 24 25		2016;100(2):242-249. doi: 10.1016/j.pec.2016.08.029	
26 27	19	. Belkora J, Volz S, Loth M, Teng A, Zarin-Pass M, Moore D, et al. Coaching patients in the u	use
28 29		of decision and communication aids: re-aim evaluation of a patient support program. BMC	
30 31 32		Health Serv Res. 2015;15:1-14. doi: 10.1186/s12913-015-0872-6	
33 34	20	. Bergenmar M, Johansson H, Wilking N, Hatchek T, Brandberg Y. Audio-recorded informati	ion
35 36		to patients considering participation in cancer clinical trials - a randomized study. Acta Onco	əl.
37 38 39		2014;53(9):1197-1204. doi: 10.3109/0284186X.2014.921726	
40 41	21	. Hack TF, Whelan T, Olivotto IA, Weir L, Bultz BD, Magwood B, et al. Standardized audiot	ape
42 43		versus recorded consultation to enhance informed consent to a clinical trial in breast oncolog	gy.
44 45 46		Psycho-Oncology. 2007;16(4):371-376. doi: 10.1002/pon.1070	
47 48	22	. Lipson-Smith R, Hyatt A, Butow P, Hack TF, Jefford M, Hale S, et al. Are audio recordings	the
49 50		answer? — a pilot study of a communication intervention for non-English speaking patients	
51 52 53		with cancer. Psycho-Oncology. 2016;25(10):1237-1240. doi: 10.1002/pon.4193	
54 55			
56 57			
58			

- 23. Lobb E, Butow P, Meiser B, Barratt A, Kirk J, Gattas M, et al. The use of audiotapes in consultations with women from high risk breast cancer families: a randomised trial. *J Med Genet*. 2002;39(9):697-703. doi: 10.1136/jmg.39.9.697
- 24. Bowden JR, Brennan PA, Butler-Keating R, Zaki GA. Use of audiotaped patient consultations in a head and neck oncology clinic and survey of patient attitudes to this facility. *J Laryngol Otol.* 2006;117(11):879-882. doi: 10.1258/002221503322542890
- 25. Masera G, Beltrame F, Corbetta A, Fraschini D, Adamoli L, Jankovic M, et al. Audiotaping communication of the diagnosis of childhood leukemia: parents' evaluation. *J Pediatr Hematol Oncol.* 2003;25(5):368-371.
- 26. Good DW, Delaney H, Laird A, Hacking B, Stewart GD, Mcneill SA. Consultation audiorecording reduces long-term decision regret after prostate cancer treatment: a non-randomised comparative cohort study. *Surg.* 2016;14(6):308-314. doi: 10.1016/j.surge.2014.10.006.
- 27. Hack TF, Pickles T, Bultz BD, Ruether JD, Degner LF. Impact of providing audiotapes of primary treatment consultations to men with prostate cancer: a multi-site, randomized, controlled trial. *Psycho-Oncology*. 2007;16(6):543-552. doi: 10.1002/pon.1094
- Purbrick RMJ, Tu KL, Damato BE. The patient's role in the decision-making process a perspective from the Liverpool Ocular Oncology Centre. *Eye*. 2005;20(9):1034-1039. doi: 10.1038/sj.eye.6702074
- 29. Haslop C. Taped consultations in a thoracic oncology lung cancer service. *Cancer Nurs Pract.* 2005;4(3):35-38. doi: 10.7748/cnp2005.04.4.3.35.c163
- Hacking B, Wallace L, Scott S, Kosmala-Anderson J, Belkora J, Mcneill A. Testing the feasibility, acceptability and effectiveness of a 'decision navigation' intervention for early stage prostate cancer patients in Scotland--a randomised controlled trial. *Psycho-Oncology*. 2013;22(5):1017-1024. doi: 10.1002/pon.3093

Psycho-Oncology

Systematic Review of Consultation Recordings

- 31. Hack TF, Ruether JD, Weir LM, Grenier D, Degner LF. Promoting consultation recording practice in oncology: identification of critical implementation factors and determination of patient benefit. *Psycho-Oncology*. 2013;22(6):1273-1282 doi:10.1002/pon.3135.
- 32. Volz S, Moore DH, Belkora JK. Do patients use decision and communication aids as prompted when meeting with breast cancer specialists? *Health Expect.* 2015;18(3):379-91. doi: 10.1111/hex.12042
- Cope CD, Lyons AC, Donovan V, Rylance M, Kilby MD. Providing letters and audiotapes to supplement a prenatal diagnostic consultation: effects on later distress and recall. *Prenat Diagn*. 2003;23(13):1060-1067. doi:10.1002/pd.747.
- 34. Shepherd L, Salkovskis PM, Morris M. Recording therapy sessions: an evaluation of patient and therapist reported behaviours, attitudes and preferences. *Behav Cogn Psychother*. 2009;37(2):141-150. doi: 10.1017/S1352465809005190.
- 35. Wolderslund M, Kofoed PE, Holst R, Ammentorp Jette. Patients' use of digital audio recordings in four different outpatient clinics. *Int J Qual Health Care*. 2015;27(6):466-72. doi: 10.1093/intqhc/mzv069
- 36. Bozic KJ, Chenok KE, Schindel J, Chan V, Huddleston JI, Braddock C, et al. Patient, surgeon, and healthcare purchaser views on the use of decision and communication aids in orthopaedic surgery: a mixed methods study. *BMC Health Serv Res.* 2014;14:1-10. doi: 10.1186/1472-6963-14-366.
- 37. Koh TH, Butow PN, Coory M, Budge D, Collie LA, Whitehall J, et al. Provision of taped conversations with neonatologists to mothers of babies in intensive care: randomised controlled trial. *BMJ*. 2007;334:28-31. doi: 10.1136/bmj.39017.675648.BE

Systematic Review of Consultation Recordings

- Belkora JK, Loth MK, Volz S, Rugo HS. Implementing decision and communication aids to facilitate patient-centered care in breast cancer: a case study. *Patient Educ Couns*. 2009;77(3):360-368. doi: 10.1016/j.pec.2009.09.012
- 39. Elwyn G, Barr PJ, Grande SW. Patients recording clinical encounters: a path to empowerment? Assessment by mixed methods. *BMJ*. 2015;5:1-8. doi: 10.1136/bmjopen-2015-008566
- 40. Hacking B, Scott SE, Wallace LM, Shepherd SC, Belkora J. Navigating healthcare: a qualitative study exploring prostate cancer patients' and doctors' experience of consultations using a decision-support intervention. *Psycho-Oncology*. 2014;23(6):665-71. doi: 10.1002/pon.3466
- 41. Jackson M, Skinner J. Improving consultations in general practice for Somali patients: a qualitative study. *Diversity in Health & Social Care*. 2007;4:61-67.
- 42. Leahy M, Douglass J, Barley V, Jarman M, Cooper G. Audiotaping the heart surgery consultation: qualitative study of patients' experiences. *Heart*. 2005;91(11):1469-1470. doi: 10.1136/hrt.2004.048769
- 43. Belkora JK, Loth MK, Chen DF, Chen JY, Volz S, Esserman LJ. Monitoring the implementation of consultation planning, recording, and summarizing in a breast care center. *Patient Educ Couns*. 2008;73(3):536-543. doi: 10.1016/j.pec.2008.07.037
- 44. Pass M, Volz S, Teng A, Esserman L, Belkora J. Physician behaviors surrounding the implementation of decision and communication AIDS in a breast cancer clinic: a qualitative analysis of staff intern perceptions. *J Cancer Educ.* 2012;27(4):764-769. doi: 10.1007/s13187-012-0402-8.
- 45. Hack TF, Carlson L, Butler L, Degner LF, Jakulj F, Pickles T, et al. Facilitating the implementation of empirically valid interventions in psychosocial oncology and supportive care. *Support Care Cancer*. 2011;19(8):1097-1105. doi: 10.1007/s00520-011-1159-z

Psycho-Oncology

Systematic Review of Consultation Recordings

- 46. Santo A, Laizner AM, Shohet L. Exploring the value of audiotapes for health literacy: a systematic review. *Patient Educ Couns*. 2005;58(3):235-243. doi: 10.1016/j.pec.2004.07.001
- 47. Polit DF, Beck CT. Nursing research: generating and assessing evidence for nursing practice.
 10th ed. Philadelphia: Lippincott Williams & Wilkins; 2017.
- 48. Elwyn G, Barr P, Castaldo, M. Can patients make recordings of medical encounters? What does the law say? *JAMA*. 2017;318(6):513-514.
- 49. Tsulukidze M, Grande SW, Thompson R, Rudd K, Elwyn, G. Patients covertly recording clinical encounters: Threat or opportunity? A qualitative analysis of online texts. *PLoS ONE*. 2015; 10(5):1-10.
- 50. Adams M. Covert recording by patients of encounters with gastroenterology providers: Path to empowerment or breach of trust? *Clin Gastroenterol Hepatol* 2017;15(1):13-16.
- 51. Grande SW, Castaldo MG, Carpenter-Song E, Griesemer I, Elwyn G. A digital advocate? Reactions of rural people who experience homelessness to the idea of recording clinical encounters. *Health Expect.* 2017; 20(4):618-625.

 Table 1. Implementation Tips for Initiating and Establishing a Consultation Recording

 Service

- Provide high quality and accessible evidence, such as systematic reviews, to ensure staff members are adequately informed and to promote practice change.
- Ensure administrative commitment of financial resources and staff to facilitate the management of recordings.
- Develop standardized procedures for consultation recordings including which consultations to record, what parts of the consultation to record, and whether the clinic will retain a copy of the recording.
- Select the most appropriate recording device for the target patient population, and ensure that technology is functioning and accessible for all consultations.
- Educate all staff, including physicians, on the procedures for recordings.
- Delegate the responsibility for recording the consultation to a specific staff member.
- Identify a respected champion with administrative or social power to promote consultation recordings.
- Inform all patients regarding the opportunity to have their consultation recorded.
- Offer patients a choice of whether to receive the recording or other communication aids to address personal preferences.
- Create a disclaimer and an informed consent form to address medico-legal concerns.
- Pilot consultation recordings so that staff can experience how it works and its impact on patients to promote buy-in.
- Establish policies and procedures for patients who desire to record consultations on their own to avoid covert recordings.

http://mc.manuscriptcentral.com/pon

Should Consultation Recording Use be a Practice Standard? A Systematic Review of the

Effectiveness and Implementation of Consultation Recordings

Kendra L. Rieger, PhD

Assistant Professor College of Nursing, Rady Faculty of Health Sciences, University of Manitoba, Canada Helen Glass Centre for Nursing, 89 Curry Place University of Manitoba, Winnipeg, Manitoba, Canada R3T 2N2 Phone: 204.474.7107 Fax: 204.474.7682 Email: Kendra.Rieger@umanitoba.ca

Thomas F. Hack, PhD

Professor

College of Nursing, Rady Faculty of Health Sciences, University of Manitoba, Canada Director, Psychosocial Oncology & Cancer Nursing Research, I.H. Asper Clinical Research Institute

Senior Scientist, Research Institute in Oncology and Hematology at CancerCare Manitoba Visiting Professor, University of Central Lancashire, Preston, UK

Kinta Beaver, PhD

Professor of Cancer Care School of Health Sciences, University of Central Lancashire, United Kingdom

Penelope Schofield, PhD

Professor, Department of Psychology, School of Health Sciences, Faculty of Health, Arts and Design, Swinburne University of Technology, Hawthorn, Australia.

Department of Cancer Experiences Research, Peter MacCallum Cancer Centre, Melbourne, Australia

Sir Peter MacCallum Department of Oncology, Faculty of Medicine, Dentistry and Health Sciences, The University of Melbourne, Parkville, Australia

Abstract

Objective: To conduct a systematic review of the effectiveness of consultation recordings, and identify factors contributing to their successful implementation in healthcare settings.

Methods: A systematic review was conducted for quantitative studies examining the effectiveness of consultation recordings in healthcare. Two independent reviewers assessed the relevance and quality of retrieved quantitative studies using standardized criteria. Study findings were examined to determine consultation recording effectiveness and to identify barriers and facilitators to implementation. A supplementary review of qualitative evidence was performed to further explicate implementation factors.

Results: Of the 3,373 articles retrieved in the quantitative search, 26 satisfied the standardized inclusion criteria (12 randomized controlled trials, 1 quasi-experiment, and 13 cross-sectional studies). The majority of patients found consultation recordings beneficial. Statistically significant evidentiary support was found for the beneficial impact of consultation recordings on the following patient reported outcomes: knowledge, perception of being informed, information recall, decision-making factors, anxiety, and depression. Implementation barriers included strength of evidence concerns, patient distress, impact of the recording on consultation quality, clinic procedures, medico-legal issues, and resource costs. Facilitators included comfort with being recorded, clinical champions, legal strategies, efficient recording procedures, and a positive consultation recording experience.

Conclusions: Consultation recordings are valuable to patients and positively associated with patient reported outcomes. Successful integration of consultation recording use into clinical practice requires an administratively supported, systematic approach to addressing implementation factors.

Background

Patients commonly seek detailed diagnostic and treatment information to gain a fuller understanding of their disease and its management.^{1, 2} Patients can be overwhelmed, however, by the complexity of information provided during healthcare consultations.³ Elevated levels of anxiety may hinder efforts to process and retain information imparted during these consultations,⁴ making it difficult to remember.^{5, 6}

The most recent Cochrane review, conducted almost one decade ago, concluded that consultation recordings in oncology "may benefit most adults with cancer…most patients find them very useful".^{5 p.2} The Cochrane review also identified further questions worth exploring, many of which have been the focus of subsequent research studies that have not been subjected to systematic review. Providing patients with audio-recordings of important consultations can reduce their anxiety, improve recall of information, and enhance their perception that they have been adequately informed about their disease and treatment.^{5.9} Recordings enable greater patient participation in decisions surrounding disease management, reduce treatment decision regret, and facilitate patient satisfaction with care.^{4, 5, 7, 8, 10} Despite the established benefits of recordings, use of this intervention in clinical practice is sporadic.

The provision of recordings of key consultations is topical and controversial. While some researchers and patient advocacy groups have argued for its routine use in clinical practice, many clinicians are hesitant to record consultations fearing litigation.¹¹ Many believe the evidence base is mixed and inconclusive, while others believe the resources required to implement the intervention preclude its use in standard care.¹² If implementation barriers are perceived to be strong enough to preclude implementation of an empirically validated intervention, then these barriers need to be remedied. The objectives of the present review were to: 1) Conduct a systematic review of the

Page 31 of 73

Psycho-Oncology

Systematic Review of Consultation Recordings

effectiveness of consultation recording use; and 2) Review the consultation recording literature to identify key implementation factors and the best ways to address them.

Methods

Review Objective One

Inclusion and exclusion criteria. To be included, studies had to involve adults or children having a clinical consultation in a healthcare setting, after which the patient or family received a consultation recording (tape or digital). Primary quantitative study designs eligible for inclusion were randomized controlled trials (RCTs), quasi-experimental studies (QE), analytical crosssectional (observational) studies, and descriptive cross-sectional studies. Studies from all countries were considered if they were written in English. We excluded studies that investigated written summaries or video recordings of consultations alone, did not provide the recording to the patient, or examined the use of standardized audio-recorded education materials. An a priori systematic review protocol guided this work. Research ethics board approval was not required for this systematic review.

Search strategy, study selection, and analysis. Online databases including MEDLINE, CINAHL, PsychINFO, EMBASE, Web of Science, Sociological Abstracts, and ProQuest Dissertations & Theses were systematically searched from January 1, 2002, to August 30, 2016 (see supporting information Table S1). A grey literature search was conducted of Conference Proceedings Citation Index, Grey Matters, Google Scholar, and relevant websites. Two independent reviewers (KR, TH) screened titles/abstracts of retrieved articles for potential inclusion criteria and one reviewer read the full-text of relevant articles to confirm eligibility. Two independent reviewers (KR, TH) critically assessed eligible RCTs with two widely used quality appraisal tools: the Joanna Briggs Institute (JBI) Critical Appraisal Tool for Randomized Controlled Trials¹³ and the Effective Public Health Practice Project Quality Assessment Tool for Quantitative Studies (EPHPP).¹⁴ Quasi-

experimental studies were assessed by two reviewers (KR, TH) with the JBI Critical Appraisal Tool for Quasi-Experimental Studies.¹⁵ To be included in the review, an RCT or quasi-experimental study had to meet key criteria determined by two reviewers (see supporting information Table S2). Disagreements were resolved through discussion between the two reviewers. The PRISMA statement for reporting systematic reviews was utilized to frame the search and study selection process.¹⁶ The heterogeneity of the samples, methods, and outcomes in the reviewed studies precluded a meta-analysis; therefore, the extracted data was analyzed using a narrative synthesis.

Review Objective Two

In addition to the descriptive findings and narrative comments contained within the quantitative papers from the first objective, a systematic review of the qualitative literature was conducted using terms associated with healthcare consultations, consultation recordings, and qualitative research.¹⁷ (see supporting information Table S1) MEDLINE, CINAHL, and the grey literature were searched from January 1, 2002, to February 5, 2017. Two reviewers screened retrieved titles/abstracts for relevance (KR, TH), and all qualitative designs were included. Only data addressing the barriers and facilitators contributing to the successful implementation of recordings were extracted and thematically analyzed from the included quantitative and qualitative studies.

Results

Quantitative Studies for Inclusion

The database search for objective one retrieved 3359 articles, and an additional 14 articles were retrieved through searching the grey literature, reference lists, and Scopus using the forward search function (see Figure 1). After removing duplicates (n=1374), 1999 titles/abstracts were examined and 1911 articles were excluded for not meeting the inclusion criteria. The full texts of the remaining articles (n=88) were reviewed in-depth, and a further 61 articles were excluded.

Psycho-Oncology

Systematic Review of Consultation Recordings

Reasons for exclusion are provided in Figure 1. After critically appraising the RCT and QE studies, 12 RCTs, 1 QE study, and 13 cross-sectional descriptive (CS) studies were included. One RCT was excluded as it did not meet our quality criteria.¹⁸

Description and Quality Appraisal of Quantitative Studies

The majority of studies (see supporting information Table S3 for study details) examined consultation recordings for patients with cancer (n=19).^{1-3, 8, 9, 19-32} Some unique oncology contexts included patients transitioning from curative to palliative care,⁸ parents of children with leukemia having an initial consultation,²⁵ and migrant patients with cancer who requested an interpreter.²² Three studies examined the usefulness of recordings for enhancing informed consent.^{4, 20, 21} The remaining seven studies occurred outside of the oncology context.^{4, 10, 33-37} The geographical settings included the United Kingdom, ^{2, 4, 10, 26, 28-30, 33, 34} United States,^{1, 19, 24, 32, 36} Canada,^{9, 21, 27, 31} Australia,^{3, 22, 23, 37} Sweden,²⁰ The Netherlands,⁸ Italy,²⁵ and Denmark.³⁵ Supporting information table S2 summarizes the RCT and QE quality appraisals. On the JBI checklist,¹³ 11 RCTs received 8-13 out of a possible 13 points, and one received 5-7 points. On the EPHPP tool,¹⁴ five RCTs received a strong global rating, five received a moderate rating, and two received a weak rating. The one QE²⁶ study received seven out of nine points on the JBI QE tool.¹⁵

Description of Qualitative Studies

In addition to the descriptive data and narrative comments from the quantitative studies, 12 qualitative studies were included to answer the second review objective. These qualitative studies were identified when screening titles/abstracts for objective one or found in the 92 articles retrieved in the systematic search for objective two. Studies explored patients' experiences/perspectives of consultation recording^{26, 31, 34, 36, 38-42} and healthcare professionals' experiences of consultation recording implementation.^{12, 31, 34, 36, 40, 43, 44} The qualitative studies took place in the United States,^{36, 38, 40, 43, 44} UK, ^{26, 34, 39, 41, 42} Canada,³¹ and Australia.¹²

Objective One: Effectiveness of Consultation Recordings

Satisfaction. Patients had an overall positive perspective of consultation recordings. Sixteen studies (7 RCTs, 9 CS) reported that the majority of participants found recordings useful or beneficial, with a range of 55-98% across studies.^{1-3, 8, 10, 20, 22-25, 28-30, 34, 36, 37} However, listening to the recording was distressing (3-18%)^{2, 20, 29} or a source of anxiety (19-26.8%)^{3, 25} for small proportions of participants. In one study, participants had a marginally significant preference for consultation recordings over standardized audio-recordings.²¹ In another study, 17 (57%) patients preferred to receive both a recording and summary letter, 7 preferred the audio-recording alone, and 3 preferred the letter alone, with participants more likely to give the recording (69%) to a family member or friend than the letter (50%).³ Five RCTs examined if recordings had an impact on the patients' satisfaction with communication with the health provider, but none found significant differences.^{9, 21, 23, 27, 37}

Use of consultation recordings. Most studies reported that the majority of participants (51-100%) listened to their recordings,^{2-4, 8-10, 19, 20, 22-25, 27-29, 31-34, 37} although one study found that only 31% of participants listened.³⁵ Some participants listened more than once, with an average use of 1.3 to 3.0 times.^{9, 10, 21, 25-28, 30, 31} Reasons cited for not listening included technological issues,^{26, 31, 35} not wanting to revisit their decision,²⁶ feeling they had received sufficient information already,^{3, 26,} ^{29, 31} worrying it would evoke negative emotions,^{20, 29} intending the recording for a family member,²⁹ being overwhelmed or too busy,^{31, 34} desiring to move on,^{3, 31} preferring another modality,³¹ feeling too sick,³¹ disliking the sound of their voice,³⁴ and lacking the privacy to listen to the recording.³⁴ When patients were prompted to record their own consultations, one study found that only 20% did so.³² Between 20 and 100% of participants in 14 studies reported that they shared their recording with others.^{1-3, 8-10, 19, 20, 22, 23, 27, 29, 31, 32} Patients shared their recording with spouses/partners, children, family members, friends, and doctors. A number of factors were
Psycho-Oncology

Systematic Review of Consultation Recordings

associated with using the recording including increased anxiety,²³ being married,³ receiving bad news,³ having longer consultations,³ increasing age,³⁵ and gender (female patients and their relatives replaying together; relatives of male patients replaying alone).³⁵

Knowledge, perception of being informed, and recall of information. Nine (75.0%) RCTs assessed the impact of recordings on knowledge, perception of being informed, or recall of information, and five found significant positive results. Of the four RCTs specifically measuring knowledge, three investigated knowledge relevant to enhancing informed consent^{4, 20, 21} and only one demonstrated significant findings.⁴ Two^{9, 27} of the three^{9, 21, 27} studies examining patients' perception of being informed, and two^{2, 37} of the three^{2, 33, 37} studies measuring recall, showed that the recording provided significant benefit.

Psychological health and well-being. Seven (58.3%) RCTs measured anxiety, depression, and/or stress,^{2, 4, 10, 23, 30, 33, 37} and significant differences were observed in three studies. In two studies, recordings significantly reduced participants' anxiety^{4, 33} and in one study it reduced their depression.⁴ Another RCT found significant decreases in anxiety and depression when the analysis was repeated with only those who had listened to the recording.²³ There were no other RCTs that reported a statistically significant impact of recordings on psychological adjustment,^{9, 27, 30} quality of life,^{8, 9, 27} treatment adherence,¹⁰ or openness to discussing cancer-related issues in the family.⁸

Decision making. Of the three studies assessing the impact of consultation recordings on concepts associated with decision making, all had significant findings. One study found significantly higher decisional self-efficacy,³⁰ and two found lower decisional regret,^{26, 30} for those who had received a recording. In the QE study, receiving a recording was positively associated with less decision regret, and was even a stronger predictor of decision regret than erectile dysfunction or incontinence.²⁶ Another RCT measured health locus of control, and found that the consultation

Systematic Review of Consultation Recordings

recording group reported a significantly higher sense of control regarding their own health than the control and standardized recording groups.⁴

Objective Two: Barriers and Facilitators to Consultation Recording

The synthesis for the second question revealed noteworthy individual, interpersonal, and system barriers to implementing consultation recordings, as well as critical facilitators.¹²

Perceived value of consultation recording and the evidence. The qualitative findings showed that both patients and practitioners valued consultation recordings.^{12, 31, 36, 39, 42} Patients appreciated recordings for the purposes of enhancing patient-centred communication and fostering empowerment.³⁹ Some practitioners questioned whether recordings benefited those with a passive decision making style.¹² Healthcare professionals had differing assessments of the empirical evidence on recordings, with the majority being unaware of the supporting evidence base.^{12, 31} The provision of high quality, accessible evidence such as systematic reviews and clinical practice guidelines was identified as an essential facilitator to ensuring staff were adequately informed, and promoting practice change.³¹

Clinician's motivation. Positive feedback motivated clinicians to continue with recordings. Piloting and evaluating consultation recording was identified as a means by which to promote the "buy-in" of stakeholders and to inspire routine implementation,¹² through exposure to the benefits.^{29, 40, 43} One study offered a feedback letter to oncologists espousing the patient-reported benefits of recordings to reinforce the importance of continuing the practice.³¹

Fear of distressing patients. Clearly identified barriers included concerns that recordings could increase patients' anxiety when replaying distressing content or impose an undesired active decision-making role on patients.^{2, 12, 29} Practitioners noted that certain visits might not be wholly appropriate for recording, such as consultations in which bad news was given, the patient became upset, or the patient was extremely anxious.^{12, 29}

Psycho-Oncology

Systematic Review of Consultation Recordings

Impact on the consultation. Variation in perspectives regarding the impact of recording on the consultation determined whether this factor was a barrier or facilitator. Some healthcare workers reported feeling anxious or self-conscious about what they said when being recorded as well as concerned that the recording would make the communication more formal, factual, or structured.^{12, 20, 29} Practitioners noted that consultations are not always a straightforward provision of information and that a desire to create a coherent recording could be a barrier.¹² In contrast, others felt at ease with being recorded⁴⁰ or thought that more detailed information was given with recordings.²⁹ Some patients thought the intervention might violate the traditional etiquette of the doctor-patient relationship; thus, challenging the clinicians' status or damaging the relationship.³⁹ Others believed that the recording process would not hinder physicians from sharing information, might increase their accountability, and could enhance respect for clinicians who were willing to be recorded.³⁹

Clinical champions. An essential facilitator of successful implementation was leadership, that is, an identified, respected champion with the administrative or social power to advocate for the intervention, and obtain necessary funds and staff resources for a consultation recording program.³¹ Support from physicians conducting the consultations was crucial for successful implementation, as was their consistency in following the recording procedures.⁴⁴

Legal and privacy concerns. There was evidence that legal and privacy concerns were substantial barriers to recordings. Physicians were concerned about medico-legal implications, such as who owns the recording, whether the treatment center should retain a copy, how to store the recordings within existing medical records, if the recording might be used in a lawsuit, and with whom the recording would be shared.^{12, 29, 31} There were also concerns about a breach of confidentiality if the consultation was shared without permission, and a loss of control of the recording if it was posted on the internet.^{1, 34, 39} An important facilitator was assuring that legal requirements were met.¹² One strategy involved creating a disclosure/consent form for patients,

Systematic Review of Consultation Recordings

providers, and the organization, and ensuring the consent process communicated the patients' and providers' rights, obligations, and the appropriate forms of distribution for the recording.¹² One center created a disclaimer which clearly stated that the recording had not been reviewed by physicians, could contain errors or omissions, and that it was primarily for the convenience and personal use of patients.⁴³

Required resources and technology. There were reservations about the amount of time that recordings might require in an already overloaded clinical setting.^{12, 24, 36, 43} However, researchers reported that recordings did not substantially increase the length of consultations,^{1, 9, 10, 24, 29} and some even found it was an efficient means of promoting the doctor-patient partnership.^{10, 12, 34} Intervention and resource costs were identified as other primary barriers to the sustainable implementation of recordings.^{12, 31, 36, 40} One cost-saving facilitator was integrating recording procedures into existing staff roles.⁴⁰ Technology was highlighted as a facilitator when no significant technical problems occurred,⁸ and as a barrier when malfunctioning or inaccessible equipment hindered the intervention.^{9, 10, 22, 29, 34, 37} Researchers cited poor recording quality as a barrier to use.^{3, 8, 24, 26, 34} With an array of recording options, selecting the most appropriate recording technology for the target patient population was an important consideration.^{34, 41} The growing accessibility of smartphones for self-recording of consultations was acknowledged as holding potential for facilitating recordings.^{12, 39}

Logistics and procedures. Logistical facilitators included recordings posing a minimal burden to the clinical environment and embedding recordings into usual care.¹² Clear procedures and staff support were critical to preventing problems with recordings and avoiding increased consultation time.^{8, 12} Significant logistical components of supporting recordings included organized scheduling, informing patients of the benefits of recordings and their option to record, ensuring the recording equipment was functioning properly, obtaining consent to record, ensuring the recording

Page 39 of 73

Psycho-Oncology

Systematic Review of Consultation Recordings

occurred despite time pressures, and delivering the recording to the patient.^{12, 24, 31, 38, 43} To address concerns of recordings causing distress, patients should be offered the choice of receiving a recording rather than systematically providing one to all patients.¹² To avoid covert self-recordings, procedures should be in place to allow patients to record their own consultations.^{12, 39}

Conclusions

This systematic review builds upon previous work by including the most recent literature and elucidating the barriers and facilitators of routine implementation. The findings demonstrate that consultation recordings enhance decision-making and patient satisfaction, and that patients frequently listen to their recordings and share them with others. Although patients derive benefits from recordings, there were equivocal findings regarding knowledge, information recall, the perception of being informed, and psychological well-being.

Only a proportion of the studies reported significant results for anxiety and depression. It may be unreasonable to expect patients' psychological well-being to be significantly improved by a recording only a short time after a life-altering diagnosis.⁹ There was also variation in patients' perceptions of recordings and identified factors that contributed to their use. Recordings may also resonate more with certain individuals depending on their preferred coping mechanisms. If patients prefer to use denial as a coping mechanism, recordings may not be valuable to them, and may actually cause anxiety.⁶

The findings of this present review had notable differences with the earlier Cochrane review⁵ A distinct finding, which was only evident in this review, was that recordings had an unequivocally positive impact on concepts associated with decision making. Lower decision regret may be a result of consultation recordings enhancing patients' understanding of the risks, benefits, and outcomes of various treatments, all contributing to more fully informed decision making.²⁶

This review also extended the Cochrane review by identifying key implementation factors and strategies to address them. Barriers to implementation of recordings included a lack of awareness of the empirical evidence on recordings, feeling self-conscious about being recorded, perceptions that the recording process will negatively impact the consultation, variation in physician cooperation with recording procedures, lack of staff and financial resources, and medico-legal concerns. Important implementation facilitators included acceptance of research evidence, valuing recordings as a communication tool, champions to promote recordings, positive feedback for clinicians who use recordings, and dedicated staff to support recordings. Producing research knowledge alone is often insufficient to change practice. This systematic identification of consultation recording implementation factors is vital to establishing and maintaining a recording service.45

Although none of the reviewed studies were conducted in low-income countries, some of the studies included participants from disadvantaged populations.^{2, 22, 41} Recordings may be especially useful for those with low literacy or for non-English speakers, as the intervention does not rely on written information alone.^{2, 22, 41} A patient in one study who spoke English as a second language thought the recordings allowed a careful review of sentences not initially understood.³ This potential of recordings is important, as literacy is identified as a determinant of health and recordings may promote health literacy through a more accessible medium.⁴⁶ Further, one study reported that the commonly observed positive association between deprivation and anxiety was not found in patients who received a recording.² Thus, recordings may reduce the negative impact of socio-economic factors. These observations are noteworthy as a review of audio-recordings for health literacy found no studies of recordings with disadvantaged or low-literacy populations.⁴⁶

Study limitations. First, we included evidence from cross-sectional descriptive studies to capture informative data about the use of, and satisfaction with, recordings in diverse settings. This

Psycho-Oncology

Systematic Review of Consultation Recordings

study design is especially vulnerable to the impact of confounding variables and alternative explanations for the results.⁴⁷ Second, there were a number of studies in which consultation recording was part of a complex intervention that included other decision or communication aids.^{19, 30, 32, 36} In a complex intervention, it may not be possible to attribute findings to a particular component of an intervention such as the recording. Lastly, all of the studies took place in Canada, US, Europe, or Australia. These findings may not be generalizable to countries in South America, Asia or Africa where there may be appreciable differences in healthcare provider-patient communication and the impact of recordings.

Implications for research. Although there has been some preliminary work carried out in regard to who benefits most from recordings and what types of consultations should be recorded, more research is needed.^{7, 8, 10} The mechanisms by which benefits occur with recordings need to be better understood as this knowledge could inform outcome variables and the most appropriate measurement time-points.^{7, 27} More research on the factors associated with valuing and utilizing recordings is needed. Continued investigation of recordings amongst diverse and disadvantaged populations would be insightful and support broader use of this intervention.⁴⁰ Further study is needed regarding healthcare providers' perspectives of patient self-recording, why more patients do not record their consultations, and how healthcare providers could enable patients to do so.³² Given the growing body of evidence supporting consultation recordings, systematic implementation studies targeting the barriers and facilitators identified in this review are also needed.³¹

Clinical implications. The current review provides convincing evidence of the effectiveness of recordings in supporting informed decision making and demonstrates that patients value and use recordings. Findings also suggest that recordings can foster recall of information, the perception of being informed, knowledge, and psychosocial adjustment in some situations. Although there were equivocal findings on certain outcomes, recordings were highly valued by patients and are not

Systematic Review of Consultation Recordings

burdensome in regard to time and costs. This strength of the evidence suggests that the routine implementation of recording use should be considered in oncology, i.e., the clinical area with the largest, most compelling evidence base. The findings of this review also indicate that recordings may be helpful for many patients but not for all. Providing patients with a choice of whether to receive the recording or other communication aids may be important to address personal preferences. Clinicians should also be aware of the potential for distress for some patients and provide adequate psychosocial supports, especially for distressing consultations.

The findings also revealed that despite the positive evidence base for this intervention, several factors impede its uptake into clinical practice. Fears of litigation are a commonly expressed rationale for not recording consultations with patients. Concerns that these recordings may turn up in social media and other public domains are frequent. However, Elwyn and colleagues⁴⁸ argue that liability insurers believe recordings actually protect clinicians, that patients primarily share their recordings with family and caregivers, and that clinicians may be able take legal action against patients who intentionally use a recording to harm their reputation. Although accurate information can allay fears, other implementation hurdles exist. Organizational resource costs must be addressed to enable successful implementation. This review suggests that for implementation to be successful, treatment centers need to be committed to providing a recording service, assign responsibility for the service, ensure recording equipment is available for all consultations, and develop staff education materials among other associated implementation processes and materials. Time is necessary for staff education, and for evaluating and maintaining the service. While senior administrative levels may express concern that there are insufficient resources to support a recording service, it is much less expensive now than in the past. The cost of recording and memory storage devises have dropped significantly over the past decade, and in some instances patients can record the consultations themselves on their cellular devices, thereby freeing up clinic resources

Page 43 of 73

Psycho-Oncology

Systematic Review of Consultation Recordings

that might otherwise have been used to purchase recording materials. The cost of educating staff in the service, and the cost of recording mediums such as digital recorders and USB memory keys is low when compared against other clinical services.

In the absence of a recording service, patients are recording consultations on their own, sometimes covertly, out of concern their clinicians may look upon them with disfavor if they express their intention or desire to have their consultations recorded. In one survey, 15% of respondents reported having covertly recorded a consultation, and 35% admitted that they would consider doing so.³⁹ If increasing numbers of patients are desiring to record consultations on their own, ^{39, 49-51} should clinical centers establish policies and procedures for addressing this trend? In most western nations, anecdotal legal opinion suggests that the law supports the right of citizens to record personal communications with a second party without second party consent, as both parties engaged in a mutually agreed to conversation are said to "own" the conversation. Moreover, an online public forum in the UK, which attracted legal commentary, concluded that patients did not have to seek permission to record consultations, and that such recordings were legal.¹¹ Additional study of the rights of patients and clinicians surrounding the use of recordings is needed to clarify the limit of any legal argument.

Moving forward, decisions should be made at the highest administrative levels as to whether the practice of recording is supported, and if so, then further decisions need to be made as to how to establish the service. If clinic staff are responsible for recording consultations, then standardized procedures will be required, including decisions surrounding which consultations to record, what parts of the consultation to record, whether the clinic will retain a copy of the recording, and whether to include a disclaimer statement for legal protection. If patients are encouraged to record consultations, then how will patients become aware of the option to record their consultations? Implementation tips have been prepared for those wanting to initiate a recording service (see Table

Psycho-Oncology

Systematic Review of Consultation Recordings

1).⁷ On the other hand, if administration decides not to support the use of recordings in practice, then procedures should be established for responding to patients who express a desire to record consultations, keeping in mind that an increasing number of patients are recording consultations covertly, and in most developed countries are likely protected by law in doing so.

Psycho-Oncology

Systematic Review of Consultation Recordings

This work was supported by the St. Boniface Hospital Foundation and a Chair in Psychosocial Oncology and Supportive Care Research awarded to Dr. Thomas Hack from the Canadian Breast Cancer Foundation.

We would like to thank Ms. Lisa Demczuk for her assistance with the search strategy.

Conflicts of interest:

We have no conflicts of interest to declare.

<text>

References

- 1. Seider MI, Damato BE. Digital audio recording of initial patient visits to an ocular oncology clinic: a pilot study. *OSLI Retina*. 2015;46(5):597-599. doi: 10.3928/2325816020150521-16
- Stephens MR, Gaskell AL, Gent C, Pellard S, Day-Thompson R, Blackshaw GR, Lewis WG. Prospective randomised clinical trial of providing patients with audiotape recordings of their oesophagogastric cancer consultations. *Patient Educ Couns*. 2008;72(2):218-222. doi: 10.1016/j.pec.2008.04.004
- 3. Knox R, Butow PN, Devine R, Tattersall MH. Audiotapes of oncology consultations: Only for the first consultation? *Ann Oncol.* 2002;13(4):622-627.
- Mishra PK, Mathias H, Millar K, Nagrajan K, Murday A. A randomized controlled trial to assess the effect of audiotaped consultations on the quality of informed consent in cardiac surgery. *Arch Surg* 2010;145(4):383-388. doi: 10.1001/archsurg.2010.45
- Pitkethly M, MacGillivray S, Ryan R. Recordings or summaries of consultations for people with cancer. *Cochrane Database Syst Rev.* 2008;3:1-58. doi:

10.1002/14651858. CD001539. pub2

- 6. McClement SE, Hack TF. Audio-taping the oncology treatment consultation: a literature review. *Patient Educ Couns.* 1999;36(3):229-238.
- Hack TF, Beaver K, Schofield P. Audio-recording cancer consultations for patients and their families – putting evidence into practice. In: Kissane DW, Bultz BD, Butow PN, Bylund CL, Noble S, Wilkinson S eds. *Oxford Textbook of Communication in Oncology and Palliative Care*. 2nd ed. Oxford: Oxford University Press; 2017. p. 56-60
- 8. Uitdehaag MJ, van der Velden L-A, de Boer MF, Spaander MC, Steyerberg EW, Kuipers EJ, et al. Recordings of consultations are beneficial in the transition from curative to palliative cancer

Psycho-Oncology

Systematic Review of Consultation Recordings

care: a pilot-study in patients with oesophageal or head and neck cancer. *Eur J Oncol Nurs*. 2012;16(2):109-114. doi: 10.1016/j.ejon.2011.04.006

- Hack TF, Pickles T, Bultz BD, Reuther JD, Weir LM, Degner LF, et al. Impact of providing audiotapes of primary adjuvant treatment consultations to women with breast cancer: a multisite, randomized, controlled trial. *J Clin Oncol*. 2003;21(22):4138-4144. doi: 10.1016/j.ejon.2011.04.006
- 10. Liddell C, Rae G, Brown TR, Johnston D, Coates V, Mallett J. Giving patients an audiotape of their GP consultation: a randomised controlled trial. *Br J Gen Pract*. 2004;54(506):667-672.
- 11. Elwyn G. "Patientgate"—digital recordings change everything. *BMJ*. 2014;348:1-2. doi: 10.1136/bmj.g2078
- Moloczij N, Krishnasamy M, Butow P, Hack TF, Stafford L, Jefford M, et al. Barriers and facilitators to the implementation of audio-recordings and question prompt lists in cancer care consultations: a qualitative study. *Patient Educ Couns*. 2017;100(6):1083-1091. doi: 10.1016/j.pec.2017.01.005.
- 13. Joanna Briggs Institute. The Joanna Briggs Institute critical appraisal checklist for randomized controlled trials. Adelaide: Joanna Briggs Institute; 2016. Available from: http://joannabriggs.org/assets/docs/critical-appraisal-tools/JBI_Critical_Appraisal-Checklist for Randomized Controlled Trials.pdf
- 14. Thomas BH, Ciliska D, Dobbins M, Mirucci S. A process for systematically reviewing the literature: providing the research evidence for public health nursing interventions. *Worldviews Evid Based Nurs*. 2004;1(3):176-184. doi: 10.1111/j.1524-475X.2004.04006.x
- 15. Joanna Briggs Institute. The Joanna Briggs institute critical appraisal checklist for quasiexperimental studies. Adelaide: Joanna Briggs Institute; 2016. Available from:

http://joannabriggs.org/assets/docs/critical-appraisal-tools/JBI_Critical_Appraisal-

 $Checklist_for_Quasi_-_Experimental_Studies.pdf$

- 16. Liberati A, Altman DG, Tetzlaff J, Mulrow C, Gøtzsche PC, Ionnidis JPA, et al. The PRISMA statement for reporting systematic reviews and meta-analyses of studies that evaluate healthcare interventions: explanation and elaboration. *BMJ*. 2009;339:1-27. doi: 10.1136/bmj.b2700
- 17. Tricco AC, Antony J, Zarin W, Strifler L, Ghassemi M, Ivory J, et al. A scoping review of rapid review methods. *BMC Medicine*. 2015;13:1-15. doi: 10.1186/s12916-015-0465-6
- Wolderslund M, Kofoed PE, Holst R, Axboe M, Ammentorp J. Digital audio recordings improve the outcomes of patient consultations: A randomised cluster trial. *Patient Educ Couns*. 2016;100(2):242-249. doi: 10.1016/j.pec.2016.08.029
- Belkora J, Volz S, Loth M, Teng A, Zarin-Pass M, Moore D, et al. Coaching patients in the use of decision and communication aids: re-aim evaluation of a patient support program. *BMC Health Serv Res.* 2015;15:1-14. doi: 10.1186/s12913-015-0872-6
- Bergenmar M, Johansson H, Wilking N, Hatchek T, Brandberg Y. Audio-recorded information to patients considering participation in cancer clinical trials - a randomized study. *Acta Oncol.* 2014;53(9):1197-1204. doi: 10.3109/0284186X.2014.921726
- 21. Hack TF, Whelan T, Olivotto IA, Weir L, Bultz BD, Magwood B, et al. Standardized audiotape versus recorded consultation to enhance informed consent to a clinical trial in breast oncology. *Psycho-Oncology*. 2007;16(4):371-376. doi: 10.1002/pon.1070
- 22. Lipson-Smith R, Hyatt A, Butow P, Hack TF, Jefford M, Hale S, et al. Are audio recordings the answer? a pilot study of a communication intervention for non-English speaking patients with cancer. *Psycho-Oncology*. 2016;25(10):1237-1240. doi: 10.1002/pon.4193

Psycho-Oncology

Systematic Review of Consultation Recordings

- 23. Lobb E, Butow P, Meiser B, Barratt A, Kirk J, Gattas M, et al. The use of audiotapes in consultations with women from high risk breast cancer families: a randomised trial. *J Med Genet.* 2002;39(9):697-703. doi: 10.1136/jmg.39.9.697
- 24. Bowden JR, Brennan PA, Butler-Keating R, Zaki GA. Use of audiotaped patient consultations in a head and neck oncology clinic and survey of patient attitudes to this facility. *J Laryngol Otol.* 2006;117(11):879-882. doi: 10.1258/002221503322542890
- 25. Masera G, Beltrame F, Corbetta A, Fraschini D, Adamoli L, Jankovic M, et al. Audiotaping communication of the diagnosis of childhood leukemia: parents' evaluation. *J Pediatr Hematol Oncol.* 2003;25(5):368-371.
- 26. Good DW, Delaney H, Laird A, Hacking B, Stewart GD, Mcneill SA. Consultation audiorecording reduces long-term decision regret after prostate cancer treatment: a non-randomised comparative cohort study. *Surg.* 2016;14(6):308-314. doi: 10.1016/j.surge.2014.10.006.
- 27. Hack TF, Pickles T, Bultz BD, Ruether JD, Degner LF. Impact of providing audiotapes of primary treatment consultations to men with prostate cancer: a multi-site, randomized, controlled trial. *Psycho-Oncology*. 2007;16(6):543-552. doi: 10.1002/pon.1094
- Purbrick RMJ, Tu KL, Damato BE. The patient's role in the decision-making process a perspective from the Liverpool Ocular Oncology Centre. *Eye*. 2005;20(9):1034-1039. doi: 10.1038/sj.eye.6702074
- 29. Haslop C. Taped consultations in a thoracic oncology lung cancer service. *Cancer Nurs Pract.* 2005;4(3):35-38. doi: 10.7748/cnp2005.04.4.3.35.c163
- Hacking B, Wallace L, Scott S, Kosmala-Anderson J, Belkora J, Mcneill A. Testing the feasibility, acceptability and effectiveness of a 'decision navigation' intervention for early stage prostate cancer patients in Scotland--a randomised controlled trial. *Psycho-Oncology*. 2013;22(5):1017-1024. doi: 10.1002/pon.3093

Systematic Review of Consultation Recordings

- 31. Hack TF, Ruether JD, Weir LM, Grenier D, Degner LF. Promoting consultation recording practice in oncology: identification of critical implementation factors and determination of patient benefit. *Psycho-Oncology*. 2013;22(6):1273-1282 doi:10.1002/pon.3135.
- 32. Volz S, Moore DH, Belkora JK. Do patients use decision and communication aids as prompted when meeting with breast cancer specialists? *Health Expect*. 2015;18(3):379-91. doi: 10.1111/hex.12042
- Cope CD, Lyons AC, Donovan V, Rylance M, Kilby MD. Providing letters and audiotapes to supplement a prenatal diagnostic consultation: effects on later distress and recall. *Prenat Diagn*. 2003;23(13):1060-1067. doi:10.1002/pd.747.
- Shepherd L, Salkovskis PM, Morris M. Recording therapy sessions: an evaluation of patient and therapist reported behaviours, attitudes and preferences. *Behav Cogn Psychother*. 2009;37(2):141-150. doi: 10.1017/S1352465809005190.
- 35. Wolderslund M, Kofoed PE, Holst R, Ammentorp Jette. Patients' use of digital audio recordings in four different outpatient clinics. *Int J Qual Health Care*. 2015;27(6):466-72. doi: 10.1093/intqhc/mzv069
- 36. Bozic KJ, Chenok KE, Schindel J, Chan V, Huddleston JI, Braddock C, et al. Patient, surgeon, and healthcare purchaser views on the use of decision and communication aids in orthopaedic surgery: a mixed methods study. *BMC Health Serv Res.* 2014;14:1-10. doi: 10.1186/1472-6963-14-366.
- 37. Koh TH, Butow PN, Coory M, Budge D, Collie LA, Whitehall J, et al. Provision of taped conversations with neonatologists to mothers of babies in intensive care: randomised controlled trial. *BMJ*. 2007;334:28-31. doi: 10.1136/bmj.39017.675648.BE

Psycho-Oncology

Systematic Review of Consultation Recordings

- Belkora JK, Loth MK, Volz S, Rugo HS. Implementing decision and communication aids to facilitate patient-centered care in breast cancer: a case study. *Patient Educ Couns*. 2009;77(3):360-368. doi: 10.1016/j.pec.2009.09.012
- 39. Elwyn G, Barr PJ, Grande SW. Patients recording clinical encounters: a path to empowerment? Assessment by mixed methods. *BMJ*. 2015;5:1-8. doi: 10.1136/bmjopen-2015-008566
- 40. Hacking B, Scott SE, Wallace LM, Shepherd SC, Belkora J. Navigating healthcare: a qualitative study exploring prostate cancer patients' and doctors' experience of consultations using a decision-support intervention. *Psycho-Oncology*. 2014;23(6):665-71. doi: 10.1002/pon.3466
- 41. Jackson M, Skinner J. Improving consultations in general practice for Somali patients: a qualitative study. *Diversity in Health & Social Care*. 2007;4:61-67.
- 42. Leahy M, Douglass J, Barley V, Jarman M, Cooper G. Audiotaping the heart surgery consultation: qualitative study of patients' experiences. *Heart*. 2005;91(11):1469-1470. doi: 10.1136/hrt.2004.048769
- 43. Belkora JK, Loth MK, Chen DF, Chen JY, Volz S, Esserman LJ. Monitoring the implementation of consultation planning, recording, and summarizing in a breast care center. *Patient Educ Couns*. 2008;73(3):536-543. doi: 10.1016/j.pec.2008.07.037
- 44. Pass M, Volz S, Teng A, Esserman L, Belkora J. Physician behaviors surrounding the implementation of decision and communication AIDS in a breast cancer clinic: a qualitative analysis of staff intern perceptions. *J Cancer Educ.* 2012;27(4):764-769. doi: 10.1007/s13187-012-0402-8.
- 45. Hack TF, Carlson L, Butler L, Degner LF, Jakulj F, Pickles T, et al. Facilitating the implementation of empirically valid interventions in psychosocial oncology and supportive care. *Support Care Cancer*. 2011;19(8):1097-1105. doi: 10.1007/s00520-011-1159-z

Systematic Review of Consultation Recordings

- 46. Santo A, Laizner AM, Shohet L. Exploring the value of audiotapes for health literacy: a systematic review. *Patient Educ Couns*. 2005;58(3):235-243. doi: 10.1016/j.pec.2004.07.001
- 47. Polit DF, Beck CT. Nursing research: generating and assessing evidence for nursing practice.
 10th ed. Philadelphia: Lippincott Williams & Wilkins; 2017.
- 48. Elwyn G, Barr P, Castaldo, M. Can patients make recordings of medical encounters? What does the law say? *JAMA*. 2017;318(6):513-514.
- 49. Tsulukidze M, Grande SW, Thompson R, Rudd K, Elwyn, G. Patients covertly recording clinical encounters: Threat or opportunity? A qualitative analysis of online texts. *PLoS ONE*. 2015; 10(5):1-10.
- 50. Adams M. Covert recording by patients of encounters with gastroenterology providers: Path to empowerment or breach of trust? *Clin Gastroenterol Hepatol* 2017;15(1):13-16.
- 51. Grande SW, Castaldo MG, Carpenter-Song E, Griesemer I, Elwyn G. A digital advocate? Reactions of rural people who experience homelessness to the idea of recording clinical encounters. *Health Expect.* 2017; 20(4):618-625.

Psycho-Oncology

Systematic Review of Consultation Recordings

Table 1. Implementation Tips for Initiating and Establishing a Consultation Recording Service

- Provide high quality and accessible evidence, such as systematic reviews, to ensure staff members are adequately informed and to promote practice change.
- Ensure administrative commitment of financial resources and staff to facilitate the management of recordings.
- Develop standardized procedures for consultation recordings including which consultations to record, what parts of the consultation to record, and whether the clinic will retain a copy of the recording.
- Select the most appropriate recording device for the target patient population, and ensure that technology is functioning and accessible for all consultations.
- Educate all staff, including physicians, on the procedures for recordings.
- Delegate the responsibility for recording the consultation to a specific staff member.
- Identify a respected champion with administrative or social power to promote consultation recordings.
- Inform all patients regarding the opportunity to have their consultation recorded.
- Offer patients a choice of whether to receive the recording or other communication aids to address personal preferences.
- Create a disclaimer and an informed consent form to address medico-legal concerns.
- Pilot consultation recordings so that staff can experience how it works and its impact on patients to promote buy-in.
- Establish policies and procedures for patients who desire to record consultations on their own to avoid covert recordings.



Figure 1. PRISMA Flow Diagram of Search and Study Selection Process



From: Moher D, Liberati A, Tetzlaff J, Altman DG, The PRISMA Group (2009). Preferred Reporting Items for Systematic Reviews and Meta-Analyses: The PRISMA Statement. PLoS Med 6(7): e1000097. doi:10.1371/journal.pmed1000097

For more information, visit <u>www.prisma-statement.org</u>.

http://mc.manuscriptcentral.com/pon

2	
3	
1	
4	
5	
6	
7	
Ω.	
0	
9	
10	
11	
12	
12	
13	
14	
15	
16	
17	
17	
18	
19	
20	
21	
21	
22	
23	
24	
25	
20	
26	
27	
28	
20	
29	
30	
31	
32	
33	
24	
34	
35	
36	
37	
20	
38	
39	
40	
41	
10	
42	
43	
44	
45	
16	
40	
47	
48	
49	
50	
50	
51	
52	
53	
51	
54	
55	
56	
57	
58	
50	
59	
60	

Table S1. Quantitative and Qualitative Review Search Strategies for MEDLINE Ovid

	consultation\$
2.	office adj3 visit\$
3.	doctor adj 3 visit\$
1.	clinic adj 3 visit\$
5.	appointment\$
5.	"patient-provider communication"
7.	patient adj3 education
3.	patient adj3 participation
).	patient adj3 communication
10.	family adj3 education
11.	family adj3 participation
12.	family adj3 communication
13.	parent adj3 education
14.	parent adj3 participation
15.	parent adj3 communication
16. 17	sibling adj3 education
I/. 10	sibling adj participation
10. 10	Physician Datient Palations/
19. 70	Professional-Patient Relations/
20. 21	Office Visits/
21.	Health education/
23.	Referral and Consultation/
24.	Decision Making/
25.	Communication/
26.	Appointments and Schedules/
27.	Health Communication/
28.	Professional-Family Relations/
29.	Access to Information/
30.	or/1-29
31.	recording\$ adj4 consultation\$
32. 22	audiotap\$
53. ∑₄	
34. 25	audiorecord\$
55. 26	audio recordo
30. 37	audio adj3 record\$
38.	digital\$ adi3 record\$
30. 39	"tane record\$"
40	"taned conversation\$"
41.	Tape Recording/
42.	or/31-41
43.	quantitative

A A 1' ' 1 1'1 / ' 1¢
44. clinical adj1 trial\$
45. allocate\$ adj2 random\$
46. "follow up" adj l study
47. "follow up" adj1 studies
48. observational adj1 study
49. observational adj1 studies
50. "cross sectional" adj1 study
51. "cross-sectional" adj1 studies
52. randomize\$
53. radomis\$
54. experiment\$
55. investigation
56. quasi-experiment\$
57. "controlled trial\$"
58. quasi-randomised
59. quasi-randomized
60. "non-randomised controlled trial\$"
61. "non-randomized controlled trial\$"
62. survey\$
63. "prospective study"
64. "pilot study"
65. "descriptive statistics"
66. cross-sectional adj3 analysis
67. Randomized Controlled Trial.pt
68. EXP Epidemiological Studies/
69. Random Allocation/
70. Double-Blind Method/
71. Single-Blind Method/
72. EXP Clinical Trial/
73. Controlled Clinical Trial.pt.
74. Clinical trial.pt
75. Multicenter Studies.pt.
76. Prospective Studies Retrospective Studies Longitudinal Studies/
77. Observational study Observational study.pt
78. Surveys and Questionnaires Regression analysis/
79. EXP Analysis of Variance/
80. Cross-sectional studies/
81. Statistics, Nonparametric/
82. or/43-81
83. 30 and 42 and 82
84. limit 83 to yr = 2002-August 30, 2016
85. limit 84 = English language
Qualitative Review Search Strategy for MEDLINE Ovid
1. consultation\$
2. office adj3 visit\$

1	
2	
2	
3	
4	
5	
6	
7	
8	
9	
10	
10	
11	
12	
13	
14	
15	
16	
17	
18	
10	
19	
20	
21	
22	
23	
24	
25	
26	
20	
21	
28	
29	
30	
31	
32	
33	
24	
34	
35	
36	
37	
38	
39	
40	
<u>_</u> 10	
41	
42	
43	
44	
45	
46	
47	
48	
- 1 0 ∕∩	
49	
50	
51	
52	
53	
54	
55	
56	
50	
5/	
58	
59	
60	

3.	doctor adj 3 visit\$
4.	clinic adj 3 visit\$
5.	appointment\$
6.	"patient-provider communication"
7.	patient adj3 education
8.	patient adj3 participation
9.	patient adj3 communication
10). family adj3 education
11	. family adj3 participation
12	2. family adj3 communication
13	B. parent adj3 education
14	l. parent adj3 participation
15	5. parent adj3 communication
16	5. sibling adj3 education
17	v. sibling adj participation
18	3. sibling adj3 communication
19	D. Physician-Patient Relations/
20). Professional-Patient Relations/
21	. Office Visits/
22	2. Health Education/
23	B. Referral and Consultation/
24	l. Decision Making/
25	5. Communication/
26	5. Appointments and Schedules/
27	7. Health Communication/
28	3. Professional-Family Relations/
29	P. Access to Information/
30). or/1-29
31	. consultation\$ adj4 recording\$
32	2. consultation\$ adj4 tape\$
33	B. consultation\$ adj4 digital
34	l. or/31-33
35	5. qualitative
36	5. interviews\$
37	7. "focus group\$"
38	B. themes
39	0. "thematic analysis"
40). Focus Groups/
41	. EXP Qualitative Research/
42	2. Interview/
43	8. or/35-42
44	4. 30 and 34 and 43
45	5. limit 44 to $yr = 2002$ -February 5, 2017
46	5. limit $45 = \text{English language}$

Page 58 of 73

Table S2: Summary of Quality Appraisals

				RCT	Րs: JBI Qua	ality Appr	aisal Tool						
Author(s)	Randomi zation	Allocation Concealment	Similar Groups	Partici pants Blinded	HCP Blinded	Outcome Assessors Blinded	Identical Treatment	Follow- up	ITT	Procedur es for Measure ment	Reliable Measure ment	Statistical Analysis	Trial Desigr
Bergenmar et al. ²⁰	U	U	Y	Y	Y	N/A	Y	N	N	Y	Y	Y	Y
Cope et al. ³³	Y	U	Y	Ν	N	U	Y	Ν	N	Y	Y	Y	Y
Hack et al.9	Y	Y	Y	Y	Y	Y	Y	U	N	Y	Y	Y	Y
Hack et al. ²⁷	Y	Y	Y	Y	Y	Y	Y	U	N	Y	Y	Y	Y
Hack et al. ²¹	Y	U	Y	Y	Y	Y	Y	U	N	Y	Y	Y	Y
Hacking et al. ³⁰	Y	Y	Y	N	N	N	Y	Y	N	Y	Y	Y	Y
Koh et al. ³⁷	Y	Y	Y	U	U	N	Y	Y	Y	Y	Y	U	Y
Liddell et al. ¹⁰	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	U	Y	Y
Lobb et al. ²³	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y
Mishra et al.4	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y
Stephens et al. ²	U	Ν	Y	Ν	N	N	Y	Y	Y	Y	Y	Y	Y
Uitdehaag et al. ⁸	U	Y	U	Y	Y	Y	Y	Y	N	Y	Y	U	Y
Key: 1-yes, N-110, e	,-uncical,			RCTS	: EPHPP Q	uality Ap	praisal Toc	ol	gs on qu		5, 1, 10, 1	1, & 12	
Author(s)	Selecti	on Bias	Study Des	ign	Confounder	rs	Blinding	Data	Collection	Witho	lrawals and	Globa	l Rating
Bergenmar et al. ²⁰	N	M	S		S		М		M		M		S
Cope et al. ³³	1	N	S		S		М		S		S		M
Hack et al. ⁹		S	S		S		S		S		S		S
Hack et al. ²⁷		S	S		S		S		S		S		S
Hack et al. ²¹	N	M	S		S		М		W		W	,	W
Hacking et al. ³⁰	/	N	S		W		М		S		S	,	W
Koh et al. ³⁷	N	N	S		S		М		S		W]	М
	1	M	S		S		М		W		S]	М
Liddell et al. ¹⁰	ſ	*1							~		n		c
Liddell et al. ¹⁰ Lobb et al. ²³	N	M	S		S		М		S		8		3
Liddell et al. ¹⁰ Lobb et al. ²³ Mishra et al. ⁴	N	M M M	S S		S S		M M		S S		S S		S S
Liddell et al. ¹⁰ Lobb et al. ²³ Mishra et al. ⁴ Stephens et al. ²		M M M M	S S S		S S S		M M W		S S S		S S S		S S M

Page 59 of 73

1

Psycho-Oncology

		•	Quasi-exper	rimental Study:	JBI Quality A	ppraisal Tool		•	
Author(s)	Clear IV and DV	Similar Groups	Similar Treatment	Control Group	Multiple Measurements	Follow-up	Procedure for Measurement	Reliable Measurement	Statistical Analysis
Good et al. ²⁶	Y	Y	Y	Y	N	N	Y	Y	Y
Key: Y=yes; N=	no								
Excluded if 2 or	r more N rating	s on questions 2	2, 7, 8, & 9						
			h	ttp://mc manu	scriptcontral	com/non			
				up.//inc.manu	Scriptcentral.	John/poh			

Table S3. Summary of Study Characteristics

	Characteristics of Randomized Controlled Trials and Quasi-experimental Study									
Study	Study Design &	Sample	Intervention	Outcome Measures	Results					
	Setting	Characteristics								
Bergenmar	RCT; participants	N=130	Consultation	Knowledge and	No statistically significant					
et al., 2014	randomly assigned to	Diagnosis:	recording	understanding	differences between groups on					
(Sweden)	an intervention group	breast, gastro-	(CR) of a	Tool: Quality of Informed	knowledge and understanding. In					
	(n=67) or a control	intestinal, or	medical	Consent	the intervention group, 43% (n=29)					
	(n=63) group.	prostate cancer	consultation		listened to the complete recording,					
	Department of	Mean age: 54	with an	Use and perception of audio-	and 9% (6) listened to parts. Of the					
1	oncology.	years (control),	oncologist	recording	patients who reported listening to					
		55 years	about a phase		the CD, 69% found it quite/vey					
		(intervention)	2 or 3		helpful in understanding					
		Gender: 23	clinical trial		information about the clinical drug					
		males and 107			trial and 78% rated it as quite/very					
		females		A	beneficial.					
Cope et al.,	RCT; participants	N=117	CR of a	State and trait anxiety	Women in the audiotape group					
2003	randomly assigned to	Diagnosis:	consultation <	Tool: Speilberger Stat-Trait	reported significantly less anxiety					
(United	four information	suspected fetal	about a mid-	Anxiety Inventory	than the control group two weeks					
Kingdom)	groups: a non-	anomalies	trimester		after the consultation. No					
	technical letter (n=29),	Mean age: 28	fetal scan	Depression	significant differences between					
	an audiotape of the	years		Tool: Beck Depression	information groups on depression					
	consultation (n=29), an	Gender: all		Inventory	scores or recall of information.					
	audiotape and non-	female			53.6% of women used the tape,					
	technical letter (n=30),	Race/		Recall of information	87.0% of women used the letter,					
	or standard	ethnicity:		Tool: semi-structured	24% of the women fully used the					
	information (n=29).	White		telephone interview assessing	tape/letter, and 64% partially used					
	Fetal Medicine Clinic,	(n=105), Asian		free and cued recall	the tape/letter.					
	Birmingham Women's	(n=5), African/								
	Hospital.	Caribbean		Use of letters and tapes						
		(n=2), Other								
		(n=2)	<u> </u>							
Hack et al.,	RCT; participants	N= 628	CR of a	Perception of having been	Women who received the audiotape					
2003	randomly assigned to	Diagnosis:	primary	informed	reported receiving significantly					
(Canada)	four groups: standard	breast cancer	adjuvant	Tool: Informed	more information about the side					
	care with no audiotape	Mean age:	treatment	Communication Scale	effects of treatment compared to					

	(n=158), audiotaped but no audiotape given (n=147), audiotaped and patient given audiotape (n=174), or audiotaped and patient offered choice of whether to receive audiotape (n=149). Tertiary or community oncology clinic (N=6 sites).	56.5 years (SD, 12 years) Gender: all female	consultation	Audiotape satisfaction Tool: Audiotape Use and Satisfaction Questionnaire Audiotape use Tool: Audiotape Use and Satisfaction Questionnaire Communication satisfaction with oncologist Tool: Patient Perception Scale Psychological adjustment/mood state Tool: Profile of Mood States Cancer-specific quality of life Tool: Functional Assessment	those who did not receive the audiotape. Women rated the audiotape positively with a mean of 83.9/100 (SD 19.6). 60.7% of patients listened to at least a portion of the tape. Participants listened to the entire tape an average of 2.2 times, and the partial tape an average of 2.5 times. No significant differences between groups on communication satisfaction with oncologist, psychological adjustment/mood state, or cancer- specific quality of life.
Hack et al., 2007 (Canada)	RCT; participants randomly assigned to four groups: standard care control with no audiotape (n=113), audiotape with no audiotape given (n=98), audiotaped and audiotape given (120), or audiotape with choice to receive audiotape (n=94). Tertiary oncology clinics (N=4 sites).	N= 425 Diagnosis: prostate cancer Mean age: 67.4 years (SD, 7.7 years) Gender: all male	CR of a primary treatment consultation	Of Cancer Therapy-BreastPerception of having beeninformedTool: InformedCommunication ScaleAudiotape satisfactionTool: Audiotape Use andSatisfaction QuestionnaireAudiotape useTool: Audiotape Use andSatisfaction QuestionnaireCommunication satisfactionwith oncologistTool: Patient Perception Scale	Participants who received the audiotape reported having been provided with significantly more information overall, more information about treatment alternatives, and more treatment side effects information compared to those who did not receive the audiotape. Participants rated the audiotape positively with a mean of 83.0/100 (SD 19.2). 65.4% of patients listened to the tape. Participants listened to the entire consultation an average of 2.8 times and a portion an average of 3.0 times. No significant differences between groups on communication

Hack et al., 2007 (Canada)	RCT; participants randomly assigned to three groups: standardized audiotape (n=22), consultation audiotape (n=20), or both audiotapes (n=27). Canadian cancer centres (N= 5 sites).	N= 69 Diagnosis: breast cancer Gender: all female Race/ ethnicity: Caucasian (98.6%) and Asian (1.4%)	CR of a clinical trial consultation with an oncologist	Psychological adjustment/mood state Tool: Profile of Mood StatesCancer-specific quality of life Tool: Functional Assessment of Cancer Therapy-ProstatePerception of being informed about the clinical trial Tool: Informed Consent QuestionnaireKnowledge of information relevant to providing informed consent Tool: Informed Consent QuestionnaireSatisfaction with communication during the consultation Tool: The Patient Perception ScaleUse and satisfaction with audiotape Tool: The Audiotape Use QuestionnaireAnalysis of tape content	satisfaction with oncologist, psychological adjustment/mood state, or cancer-specific quality of life. No significant differences between groups on perception of being informed, knowledge, or satisfaction with communication. There was a marginally significant preference for a consultation recording over a standardized audiotape.
Hacking et al., 2013 (United	RCT; participants randomly assigned to usual care group	N= 113 Diagnosis: primary	Providing a CD of the consultation	Confidence about treatment decisions Tool: Decisional Self-efficacy	Significantly higher decisional self- efficacy and lower decisional regret for the intervention group at 6
Kingdom)	(n=60) or an intervention group (n=63). Diagnostic clinic.	prostate cancer Mean age: 67.2 years (control), 65.4	as part of a complex intervention	Decisional conflict Tool: Decisional Conflict Scale	months. Decisional conflict was significantly lower for intervention group patients initially but not at follow-up. No significant

	Western General Hospital.	years (intervention) Gender: all		Decisional regret Tool: Decisional Regret scale	differences between groups on measurements of mood and adjustment. 91.9% found the
		male Race/ethnicity:		Measurements of Mood and Adjustment	decision navigation intervention very helpful and 8.1% somewhat helpful Intervention group
		(100%)		Tool. Hospital Anxiety and	participants used the consultation
		()		Depression Scale and Mental	recording an average of 2.35 times.
				Adjustment to Cancer Scale	
				Rating of the navigation	
				intervention	
				Tool: 5-item navigation rating	
				scale based on the Decision	
				Preparation Measure	
Koh et al.,	RCT; participants	N=200	CR of	Recall of information	Mothers in the intervention group
2007	randomly assigned to	mothers	consultations	Tool: face-to-face telephone	were able to recall significantly
(Australia)	an intervention group	Diagnosis:	with a	interview with mothers	more information about diagnosis,
	(n=102) or a control	infant in NICU	neonatologist		treatment, and outcome compared
	group (n=98).	Gender: all		Attitudes to and Use of the	to mothers in the control group at
	Care Unit Townsville	Ternale		Tool: not described	found the tape helped them to
	Hospital			1001. not described	understand reminded them of the
	110spitai.			Satisfaction with	discussion and helped a family
				conversations	member to understand 91% of the
				Tool [·] not described	mothers listened to the tape.
					mothers listened to the tape a range
				Postnatal depression	of 1-10 times. No significant
				Tool: Edinburgh Postnatal	differences between groups on
				Depression Scale	satisfaction with conversations until
					a sub-group analysis demonstrated
				Anxiety	that mothers of babies with poor
				Tool: Spielberger State	outcomes in the intervention group
				Anxiety Inventory	were significantly more satisfied
					with the conversation. No
				Stress about parenting	significant differences between
				Tool: General Health	groups on depression, anxiety, or

				Questionnaire	stross about paranting
Liddell et al., 2004 (United Kingdom)	RCT; participants randomly assigned to an experimental group (n=95) or a control group (n=85). General practice (N=2 sites).	N= 180 Mean age: 40 years Diagnosis: varied; attending a GP appointment Gender: male (n=92) and female (n= 88)	CR of a routine consultation in a general practice setting	QuestionnaireUse of tapeAdherence to GP's adviceTool: Single item measureAnxiety about conditionTool: single item measure	stress about parenting. 64% found the CR 'useful' or 'very useful'. 61% of patients listened to the tape by 7-10 days. Participants listened to the tape an average of 1.5 times. No significant between groups on adherence to GP's advice or anxiety.
Lobb et al., 2002 (Australia)	RCT; participants randomly assigned to an intervention group (n= 98) or a control group (n= 95). Familial cancer clinics (N=10).	N= 193 Diagnosis: High risk breast cancer families Mean age: 44 years (control), 45 years (intervention) Gender: all female	CR of initial genetic counseling appointment	Psychological measuresTools: Impact of Events Scale,Hospital Anxiety andDepression Scale, and theMonitoring-Blunting StyleScaleRisk perceptionTool: perceived risk wasassessed by asking the womento choose between sevenpossible responsesSatisfaction with the geneticcounseling session and withthe information receivedTool: Satisfaction with GeneticCounseling ScaleSatisfaction with audiotapeTool: Satisfaction withaudiotape questions adaptedfrom another studyUse of the audiotape	Significant differences between groups on anxiety and depression when the analysis was repeated with those women who had listened to the tape. Unaffected women in the intervention group were significantly less likely to be accurate in their risk perception. No statistically significant differences between groups on satisfaction with the genetic counseling session or breast cancer information received. The majority of women who listened to the tape found it helpful, 20% found it satisfactory, and 35% found it very or extremely helpful. 51% listened to the tape. 31% of women listened to the audiotape once, 16% listened to it twice, and 3% listened to it three times.

5 of 73			

				Breast Cancer Knowledge Tool: Breast Cancer Knowledge Scale	
Mishra et al., 2010 (United Kingdom)	RCT; participants randomly assigned to three groups: a control group (n=29), a generic tape group which received a standardized tape about the surgery (n=25), or a consultation group (n=30) who received a tape of their consultation interview. Tertiary health care center.	N= 84 Diagnosis: elective first- time coronary artery surgery Mean age: 67 years (control & generic tape group), 66 years (CR group) Gender: 60 males and 24 females	CR of an outpatient consultation on informed consent for cardiac surgery	 Knowledge Tool: Knowledge Questionnaire Locus of Control Tool: Multidimensional Health Locus of Control Questionnaire Anxiety and Depression Tool: Hospital Anxiety and Depression Scale 	The CR group had significantly higher mean knowledge, greater locus of control, and less anxiety and depression.
Stephens et al., 2008 (United Kingdom)	RCT; participants randomly assigned to an intervention group (n= 31) or a control group (n= 27). Practice of a surgeon specializing in upper gastrointestinal surgery.	N= 58 Diagnosis: esophageal or gastric cancer Median age: 69 years (control) & 66 years (intervention) Gender: male: female ratio; 22:9 (intervention) and 21:6 (control)	CR of a consultation in which a new diagnosis of esophageal or gastric cancer was given	Information retention Tool: structured interview Psychological outcome Tool: Hospital Anxiety and Depression Questionnaire Socio-economic deprivation Tool: National Indices of Multiple Deprivation Attitudes to tape and consultation Tool: questionnaire regarding attitude to tape	Participants in the intervention group retained significantly more information about their diagnoses and treatment. No significant differences between groups on psychological outcomes. Deprivation correlated with higher HAD anxiety scores for the control group but not for the intervention group, and was not associated with information retention. 90% of the intervention group participants found the tape helpful. 29 of the 31 intervention group participants listened to the tape. Patients listened to the tape a median of 1 time (0-10 times range).

Uitdehaag et al., 2012 (The Netherland s)	RCT; participants randomly assigned to a CD group (n=10) or a no-CD group (n=7). Practice of 2 gastroenterologists, 3 otolaryngologists, and 3 surgeons (N=8 sites).	N= 17 Diagnosis: incurable or recurrent oesophageal or head and neck cancer Mean age: 68years (intervention), 62years (control) Gender: 12 males and 5 females	CR of patients being told of new diagnosis and decision to move to palliative care	Quality of LifeTools: EORTC QLQ-C15-PALOpenness to discussingcancer-related issues in thefamilyTool: Openness to DiscussCancer in the Family ScaleFeasibilityTool: researcher developedquestionnaireUse of CDTool: researcher developedquestionnaire	No significant differences between groups on quality of life or openness to discussing cancer- related issues in the family. No major technical or procedural problems occurred. 71% appreciated receiving the CR. 80% listened to the CR.
Good et al., 2016 (United Kingdom)	Quasi-experiment; participants assigned to an audio recording group (n=40) or a control group (n=27). Prostate cancer clinic of one urologist.	N= 67 Diagnosis: prostate cancer Mean age: 63.5years (control), 64.4years (intervention) Gender: all male Race/ethnicity: 96.3% Caucasian (control group); 97.5% Caucasian (intervention group)	CR of the consultation where diagnosis and management of prostate cancer were discussed	Quality of life Tools: EORTC QLQ-C30/PR 25 Decision regret Tool: Decision Regret Scale Questionnaire Patient satisfaction with CR Patient satisfaction questionnaire developed by researchers; included open- ended questions	Non-significant trend towards better global health score in the CR group. The CR group had significantly different (lower) bowel symptoms. Decision regret was significantly lower in the intervention group. In the QE study, receiving a CR was positively associated with less decision regret, and was even a stronger predictor than erectile dysfunction or incontinence. The mean score for usefulness was 1.8 on a 1-5 scale with 1 being very useful. The average amount of times patients listened to the CR was 3.

		Characteristics of C	Cross-sectional Desc	riptive Studies
Study	Study Design and Setting	Sample Characteristics	Consultation Recording	Results:
Belkora et al., 2015 (United States)	Cross-sectional descriptive study. The multidisciplinar y Breast Care Center at the University of California.	N=1812 Diagnosis: breast cancer	CR of new patients' consultations with a breast care center specialist	60% listened to the CR, 41% shared the CR with another person, and 84% would recommend CR.
Bowden et al., 2003 (United Kingdom)	Cross-sectional descriptive study. A multidisciplinar y head and neck oncology clinic.	N=50 patients Diagnosis: head and neck cancer Mean age: 66 years Gender: 24 males and 26 females	CR of new and returning patients' consultations	36 (92%) found CR useful and beneficial (30/33 of the review patients, and all 6 new patients). 88% of the review patients and 100% of the new patients listened to the tape either by themselves or with another person.
Bozic et al., 2014 (USA)	Mixed methods study with a cross-sectional descriptive component. Orthopedic practice.	N= 26 patients, 518 surgeons, and 6 employer members Diagnosis: hip or knee arthritis	Evaluation of decision and communication aids, one of which was consultation recordings	The most frequently reported benefit of CR was that it enabled the patient to review/access the consultation information (58%). 80% (19/24) of the patients reported that they would want to receive a CR of future consultations. Regarding surgeons perceptions of communication aids, 68% believed that they would improve patient satisfaction and 56% believed that CR would improve the quality of the physician-patient interaction.
Hack et al., 2013 (Canada)	Mixed methods study with a cross-sectional descriptive component. Tertiary oncology centres (N=3 sites).	N=228 patients Diagnosis: breast or prostate cancer Age: 59.8 years (SD, 15.3) Gender: 54 males and 174 females	CR of an initial postsurgical breast adjuvant treatment consultation or an initial, post- diagnostic prostate treatment consultation with an oncologist	Patients rated CR highly with 93.6% rating it between 75 and 100 out of 100, and an average rating of 93.8 (S.D. = 13.7). 68.9% of the participants listened to a portion of the CR within one week of the consultation. Patients listened to the complete recording an average of 2.0 (S.D. = 1.8) times and a portion of the recording an average of 21.7 times (SD=1.3). 58.6% of the patients shared the CR with another person with the average number of other persons being 0.8 (S.D.= 0.5).

Haslop, 2005 (United Kingdom)	Cross-sectional descriptive study. Thoracic Oncology Service, Papworth Hospital	N= 57 patients, 17 staff Diagnosis: lung cancer and other diagnoses	CR of patient consultation at a joint physician and oncologist clinic	39 (98%) reported that CR was beneficial, 89% reported that they would have future consultations recorded, and 95% said they would recommend it to others. 70% (n=40) of the patients listened to the CR, and 60% listened to the CR with a family member and one listened to it with their GP.
Knox et al., 2002 (Australia)	Cross-sectional descriptive study. Oncology outpatient appointment with one oncologist.	N= 52 patients Diagnosis: cancer Mean age: 51 years Gender: 39 females and 13 males	CR of routine follow-up oncology consultations	77% thought that the CR was useful and 86.7% listened to the CR. 26 patients listened to the CR, and 14 listened more than once. 46.5% shared the CR with another person. 57% of patients preferred to receive both the tape and letter, three preferred the tape, and seven preferred the letter. Significant predictors of wanting to keep the audiotape included being married, receiving bad news, and longer consultations.
Lipson- Smith et al., 2016 (Australia)	Cross-sectional descriptive study. Cancer hospital.	N=23, and 16 received the intervention Diagnosis: cancer Mean age: 66 years (SD, 8.6) Gender: 14 males, 9 females Language: Arabic 4.4%, Cantonese 21.7%, Greek 30.4%, or Mandarin 43.5%	Mixed intervention including cancer information sheets, a question prompt list, and CR of an oncology consultation	77% found the CR useful for helping with recall and understanding, making the doctor more attentive, and helping the family. 81% listened to the CR, with 31% listening more than once. 77% stated that family members also listened.
Masera et al., 2003 (Italy)	Cross sectional descriptive study and quality improvement project.	N= 49 sets of parents Diagnosis: leukemia Mean age of children: 5.2 years Gender of children: 35 males, 28 females	CR of initial consultation when the leukemia diagnosis, program of care, and prognosis were discussed	89% of the participants reported that they were happy to receive the CR, and 82.9% of the couples thought that the CR helped them to understand the information. Most participants (87.8%) recommended that the tape would be helpful for future families. 81.6% of the participants listened to it. Participants listened to the tape an average of 2.9 times, with a range of 1-12 times.
Purbick et al., 2006 (United Kingdom)	Cross-sectional descriptive study. Tertiary adult ocular	N= 39 patients Mean age: 63 years Gender: 12 males and 27 females	Patients provided with a CR of their consultation	74.3% of the participants rated CR as being extremely useful, 80% reported listening to it. The participants used it an average of 2.2 times and it was the most popular resource provided by the oncology centre.

	oncology centre.			
Seider et al., 2015 (USA)	Cross sectional descriptive study. Ocular Oncology Clinic, University of California.	N=13 patients Diagnosis: indeterminate or malignant ocular tumors Mean age: 56.6 years Gender: 4 males and 9 females	CR of initial consultation with an ocular oncologist	Patients found that the consultation recording was useful for understanding their diagnosis (85% very useful and 15% somewhat useful), understanding their prognosis (77% very useful and 23% somewhat useful), and understanding their treatment options and possible side- effects (85% very useful and 15% somewhat useful). Of the 13 participants, five people listed to the CR once (38%), six people (46%) listened to the CR two to three times, and two people (15%) listened to it four to six times. 54% shared the recording with someone else.
Shepherd et al., 2009 (United Kingdom)	Mixed methods study with a cross-sectional descriptive component. A specialist treatment and research center specializing in cognitive behavioral therapy for anxiety disorders.	N= 72 patients at the start of therapy and 31 at the end of therapy; 15 therapists. Diagnosis: Of the 72 patients, 78% were being treated for social anxiety, 63% for OCD, 48% for panic disorder, and 24% for PTSD	CR of cognitive behavioral sessions	80% found the CR helpful, and 90% of patients listened to the CR between therapy sessions. All therapists rated the CR as helpful (100%). Patients listened to the CR at least once (50%), sometimes (23%), more than once (20%), and not at all (7%).
Volz et al., 2015 (USA)	Cross-sectional survey. University of California Breast Care Center.	N= 82 participants Diagnosis: breast cancer Mean age: 57 years Gender: all female Race/Ethnicity: 71% were White	CR of a consultation with a breast cancer specialist	Only 20% (16/79) of patients who were prompted to make recordings did so when prompted. 53% of those who made recordings listened to it, and 40% shared it with someone else.
Wolderslund et al., 2015 (Denmark)	Cross-sectional descriptive study. Danish Regional Hospital (N= 4	N= 2784 (654 pediatric, 704 orthopedic, 737 internal medicine and 689 urological patients) Median age (years): 9	CRs of consultations.	 31% of the patients listened to the CR within 90 days of the consultation, and 19 individuals at a later date. 33.3% replayed more than once. Male patients and their relatives had a lower probability of listening to the CR together compared with female patients and their

outpatient clinics: pediatrics, urology, orthopedic internal medicine.	(pediatric), 64 (orthopaedic), 60 (internal medicine), and 69 (urological) s, and Gender (male): 55.7% (pediatric), 42.9% (orthopaedic), 44.2% (internal medicine), and 81.9% (urological)	ristics of Qualitative Studies	relatives. Relatives of male patients had a higher probability of listening to the CR on their own compared with female patients' relatives.	
Study Study Des	ign and Setting	Participants	Study Focus	
Belkora et Mixed met	hods study with a qualitative	Surgeons (5) and oncologists (9) who	Experiences of the implementation of	
al., 2008 component	t; data included semi-structured	held consultations: schedulers and	consultation planning, recording, and	
(USA) interviews	data analysis not specified.	premedical staff interns (14) who	summarizing	
Consultation	ons with breast cancer patients	facilitated the complex intervention.		
making tre	atment decisions at a breast care			
center.				
Belkora et Individual	case study; data included program	A 36 year old with stage 1 breast	Experiences of decision support	
al., 2009 records, qu	alitative survey data, and a semi-	cancer.	systems which included CR	
(USA) structured	interview; data analysis not			
specified.	Consultation with a breast cancer			
care center				
Bozic et al., Mixed met	hods study with a qualitative	Patients with hip or knee arthritis.	Experiences of decision and	
2014 (USA) component	t; data included open-ended survey	orthopedic surgeons, and healthcare	communication aids, one of which	
responses,	in-depth telephone interviews, and	purchasers. Patients (n=26) were	was consultation recordings	
field notes	; thematic data analysis.	surveyed. Of these participants, 13		
Orthopedia	e practice.	completed a telephone interview.		
Elwyn et al., Mixed met	hods study with a qualitative	17 online survey respondents who also	Motivation for patients' self-	
(2015) component	t; data included semi-structured	participated in a semi-structured	recording their clinical encounters	
(United interviews)	, thematic data analysis. Radio	interview.	(covertly or otherwise)	
Cood at al Oussi arr	priment with a qualitative	Mala nationts (N=40) undersaina	Experiences of receiving a	
2016 (United component	t data included open-ended	management of prostate cancer in the	consultation recording	
Kingdom) auestions (on a questionnaire: thematic data	IIK		
analysis P	rostate cancer clinic of one			
urologist.				
Hack et al., 2013 (Canada)	Mixed methods with a qualitative component; data included semi-structured interviews and focus groups; content	228 patients who had been newly diagnosed with breast $(n=174)$ or prostate $(n=54)$ cancer; 32 patients	Experiences of consultation recording and of implementing consultation recording	
---	---	---	--	
TT 1 ·	analysis. Tertiary care oncology centers.	oncologists.		
Hacking et al., 2014 (USA)	Qualitative study; data included semi- structured telephone interviews with patients and semi-structured face-to-face interviews with doctors; data analysis consisted of framework analysis with a priori themes and constant comparison.	Men with prostate cancer (n=6) who received decision navigation and doctors (n=4) who consulted with the patients.	Experiences of consultations using Decision Navigation, which included consultation recording	
Jackson et al., 2007 (United Kingdom)	Qualitative study; data included semi- structured telephone interviews; thematic data analysis. Inner-city medical center in a community with a high proportion of Somali residents.	58 Somali patients having a general practice consultation who needed an interpreter to accompany them.	Experiences of receiving personalized audio information from the consultation provided in one's own language.	
Leahy et al., 2005 (United Kingdom)	Mixed methods study with a qualitative component; data included semi-structured interviews; content analysis. Patients having a heart surgery consultation.	19 patients who had a heart surgery consultation (10 received a recording and 9 did not)	Experiences of audiotaping the heart surgery consultation	
Moloczij et al. (2016) (Australia)	Interpretive description; data included semi- structured interviews; thematic descriptive analysis. Metropolitan oncology departments.	20 participants which included 13 doctors and 7 hospital administrators.	Implementing CR and question prompt lists into usual care	
Pass et al., 2012 (USA)	Qualitative study; data included case notes written about each patient; modified grounded theory data analysis. Breast care center.	10 staff interns at a breast cancer clinic.	Experiences of physician engagement in the implementation of decision and communication aids (including CR)	
Shepherd et al., 2009 (United Kingdom)	Mixed methods study with a qualitative component; data included a questionnaire with open-ended questions; thematic data analysis. A treatment and research center specializing in cognitive behavioral therapy for anxiety disorders.	Patients (72 at the start of therapy and 31 at the end of therapy) and 15 therapists	Experiences of audio-recording cognitive behavioral sessions	

AMSTAR – a measurement tool to assess the methodological quality of systematic reviews.

1. Was an 'a priori' design provided?

The research question and inclusion criteria should be established before the conduct of the review.

Note: Need to refer to a protocol, ethics approval, or pre-determined/a priori published research objectives to score a "yes."

2. Was there duplicate study selection and data extraction?

There should be at least two independent data extractors and a consensus procedure for disagreements should be in place.

Note: 2 people do study selection, 2 people do data extraction, consensus process or one person checks the other's work.

3. Was a comprehensive literature search performed?

At least two electronic sources should be searched. The report must include years and databases used (e.g., Central, EMBASE, and MEDLINE). Key words and/or MESH terms must be stated and where feasible the search strategy should be provided. All searches should be supplemented by consulting current contents, reviews, textbooks, specialized registers, or experts in the particular field of study, and by reviewing the references in the studies found.

Note: If at least 2 sources + one supplementary strategy used, select "yes" (Cochrane register/Central counts as 2 sources; a grey literature search counts as supplementary).

4. Was the status of publication (i.e. grey literature) used as an inclusion criterion?

The authors should state that they searched for reports regardless of their publication type. The authors should state whether or not they excluded any reports (from the systematic review), based on their publication status, language etc.

Note: If review indicates that there was a search for "grey literature" or "unpublished literature," indicate "yes." SIGLE database, dissertations, conference proceedings, and trial registries are all considered grey for this purpose. If searching a source that contains both grey and non-grey, must specify that they were searching for grey/unpublished lit.

5. Was a list of studies (included and excluded) provided? A list of included and excluded studies should be provided.

not of moluleu and excluded studies should be provided.

Note: Acceptable if the excluded studies are referenced. If there is an electronic link to the list but the link is dead, select "no."

6. Were the characteristics of the included studies provided?

In an aggregated form such as a table, data from the original studies should be provided on the participants, interventions and outcomes. The ranges of characteristics in all the studies analyzed e.g., age, race, sex, relevant socioeconomic data, disease status, duration, severity, or other diseases should be reported.

vYes □ No □ Can't answer □ Not applicable

Can't answer

□ Not applicable

Note: Acceptable if not in table format as long as they are described as above.

Ves No Can't answer Not applicable

Not applicable

Can't answer

b Yes

n No

d√es □ No □ Can't answer □ Not applicable

ty¥es □ No □ Can't answer

res

🗆 No

□ Not applicable

1 2 3

4

5

6

7

8

9

10

11

12

13 14

15 16

17

18

19

20

21 22

23

24 25

26

27

28

29

30

31

32

33 34 35

36 37

38

39

40

41

42

43 44

45

46

47

48

49

50 51 52

53 54

55

7. Was the scientific quality of the included studies assessed and documented? 'A priori' methods of assessment should be provided (e.g., for effectiveness studies if the author(s) chose to include only randomized, double-blind, placebo controlled studies, or allocation concealment as inclusion criteria); for other types of studies alternative items will be relevant.

Note: Can include use of a quality scoring tool or checklist, e.g., Jadad scale, risk of bias, sensitivity analysis, etc., or a description of quality items, with some kind of result for EACH study ("low" or "high" is fine, as long as it is clear which studies scored "low" and which scored "high"; a summary score/range for all studies is not acceptable).

8. Was the scientific quality of the included studies used appropriately in formulating conclusions?

The results of the methodological rigor and scientific quality should be considered in the analysis and the conclusions of the review, and explicitly stated in formulating recommendations.

Note: Might say something such as "the results should be interpreted with caution due to poor quality of included studies." Cannot score "yes" for this question if scored "no" for question 7.

9. Were the methods used to combine the findings of studies appropriate?

For the pooled results, a test should be done to ensure the studies were combinable, to assess their homogeneity (i.e., Chi-squared test for homogeneity, I^2). If heterogeneity exists a random effects model should be used and/or the clinical appropriateness of combining should be taken into consideration (i.e., is it sensible to combine?).

Note: Indicate "yes" if they mention or describe heterogeneity, i.e., if they explain that they cannot pool because of heterogeneity/variability between interventions.

10. Was the likelihood of publication bias assessed?

An assessment of publication bias should include a combination of graphical aids (e.g., funnel plot, other available tests) and/or statistical tests (e.g., Egger regression test, Hedges-Olken).

Note: If no test values or funnel plot included, score "no". Score "yes" if mentions that publication bias could not be assessed because there were fewer than 10 included studies.

11. Was the conflict of interest included?

Potential sources of support should be clearly acknowledged in both the systematic review and the included studies.

Note: To get a "yes," must indicate source of funding or support for the systematic review AND for each of the included studies.

Shea et al. BMC Medical Research Methodology 2007 7:10 doi:10.1186/1471-2288-7-10

Additional notes (in italics) made by Michelle Weir, Julia Worswick, and Carolyn Wayne based on conversations with Bev Shea and/or Jeremy Grimshaw in June and October 2008 and July and September 2010.

Hes D NO Can't answer Not applicable

Pres □ No □ Can't answer □ Not applicable

d res □ No □ Can't answer □ Not applicable

□ Yes ↓No □ Can't answer □ Not applicable

d Yes □ No □ Can't answer □ Not applicable