

Central Lancashire Online Knowledge (CLoK)

Title	Effectiveness and Safety of Antibiotics for Preventing Pneumonia and Improving Outcome after Acute Stroke: Systematic Review and Meta-analysis
Type	Article
URL	https://clock.uclan.ac.uk/23967/
DOI	https://doi.org/10.1016/j.jstrokecerebrovasdis.2018.07.001
Date	2018
Citation	Badve, Monica S, Zhou, Zien, Anderson, Craig and Hackett, Maree (2018) Effectiveness and Safety of Antibiotics for Preventing Pneumonia and Improving Outcome after Acute Stroke: Systematic Review and Meta-analysis. Journal of Stroke and Cerebrovascular Diseases. ISSN 1052-3057
Creators	Badve, Monica S, Zhou, Zien, Anderson, Craig and Hackett, Maree

It is advisable to refer to the publisher's version if you intend to cite from the work.
<https://doi.org/10.1016/j.jstrokecerebrovasdis.2018.07.001>

For information about Research at UCLan please go to <http://www.uclan.ac.uk/research/>

All outputs in CLoK are protected by Intellectual Property Rights law, including Copyright law. Copyright, IPR and Moral Rights for the works on this site are retained by the individual authors and/or other copyright owners. Terms and conditions for use of this material are defined in the <http://clock.uclan.ac.uk/policies/>

Table 1. Characteristics of studies included

Study	Number of participants	Key Inclusion criteria	Baseline NIHSS median (Range)	Nasogastric tube	Intervention	Control	Time to first dose of antibiotic (Mean) hours	Length of follow-up
De Falco et al, 1998 [15]	80	acute ischemic stroke, interval between onset and hospitalization <12 hrs	NR	NR	Penicillin or penicillin derivatives	No preventive antibiotics	NR	NR
Chamorro, et al, 2005 [11] (ESPIAS)	136	ischaemic or haemorrhagic stroke, age > 18 years, onset <24 hrs, NIHSS \geq 5	11-14 (7-19)	NR	Levofloxacin 500 mg for 3 days	Placebo	14.1	12 weeks
Harms et al, 2008 [12] (PANTHERIS)	79	acute ischemic stroke (9-36 hrs after onset) in MCA territory, NIHSS score \geq 12, age \geq 18 years	15-17 (12-25)	64-67%	Moxifloxacin 400 mg iv for 5 days	Placebo	24	24 weeks

Study	Number of participants	Key Inclusion criteria	Baseline NIHSS median (Range)	Nasogastric tube	Intervention	Control	Time to first dose of antibiotic (Mean) hours	Length of follow-up
Schwarz et al, 2008 [16] (MISS)	60	age \geq 18 years, stable deficit, ischemic stroke with onset of symptoms <24 hrs, mRS score $>$ 3	15-16.5 (5-28)	NR	Mezlocillin 2g plus sulbactam 1g iv 8 hourly for 4 days	No preventive antibiotics	10.9	12 weeks
Westendorp et al, 2015 [17] (PASS)	2538	ischaemic or haemorrhagic stroke, onset<24 hrs, NIHSS score \geq 1, age \geq 18 years	5 (3-9)	NR (Dysphagia range 26-27%)	Ceftriaxone 2g iv once daily for 4 days	No preventive antibiotics	24	12 weeks
Kalra et al, 2015 [13] (STROKE-INF)	1217	age $>$ 18 years, ischaemic or haemorrhagic stroke, with onset of symptoms within 48 hrs, unsafe to swallow	14-15 (9-20)	22-27%	Amoxycillin or co-amoxyclav with clarithromycin or amoxycillin alone or amoxycillin plus metro or cephalosporins or other	No preventive antibiotics	48	12 weeks

Study	Number of participants	Key Inclusion criteria	Baseline NIHSS median (Range)	Nasogastric tube	Intervention	Control	Time to first dose of antibiotic (Mean) hours	Length of follow-up
					antibiotics for 7 days			

Abbreviations: DM, diabetes mellitus; HT, hypertension; NIHSS, National Institute of Health stroke scale; NR, not reported; mRS score, modified Rankin scale score.

*Number of participants