

## **Central Lancashire Online Knowledge (CLoK)**

| Title    | Does the Addition of Non-Approved Inclusion and Exclusion Criteria for rtPA  |
|----------|--|
|          | Impact Treatment Rates? Findings in Australia, the UK, and the USA   |
| Type     | Article  |
| URL      | https://clok.uclan.ac.uk/24195/  |
| DOI      |  |
| Date     | 2019   |
| Citation | Watkins, Caroline Leigh, Lightbody, Catherine Elizabeth, Craig, Louise, Middleton, Sandy, Hamilton, Helen, Cudlip, Fern, Swatzell, Victoria, Alexandrov, Andrei, Philip, Sheeba et al (2019) Does the Addition of Non-Approved Inclusion and Exclusion Criteria for rtPA Impact Treatment Rates? Findings in Australia, the UK, and the USA. Interventional Neurology, 8 (1). ISSN 1664-9737 |
| Creators | Watkins, Caroline Leigh, Lightbody, Catherine Elizabeth, Craig, Louise, Middleton, Sandy, Hamilton, Helen, Cudlip, Fern, Swatzell, Victoria, Alexandrov, Andrei, Philip, Sheeba, Cadilhac, Dominique, McInnes, Elizabeth, Dale, Simeon and Alexandrov, Anne  |

It is advisable to refer to the publisher's version if you intend to cite from the work.

For information about Research at UCLan please go to <a href="http://www.uclan.ac.uk/research/">http://www.uclan.ac.uk/research/</a>

All outputs in CLoK are protected by Intellectual Property Rights law, including Copyright law. Copyright, IPR and Moral Rights for the works on this site are retained by the individual authors and/or other copyright owners. Terms and conditions for use of this material are defined in the <a href="http://clok.uclan.ac.uk/policies/">http://clok.uclan.ac.uk/policies/</a>

Supplement Table A Key stroke service demographics

| Hospital setting          | AUS     | UK            | USA           |  |  |  |
|---------------------------|---------|---------------|---------------|--|--|--|
|                           | (N=63)  | (N=93)        | (N=229)       |  |  |  |
|                           | n (%)   |               |               |  |  |  |
| Tertiary                  | 29 (46) | 49 (53)       | 66 (29)       |  |  |  |
| Non-tertiary              | 30 (48) | 43 (46)       | 152 (66)      |  |  |  |
| Private                   | 4 (6.4) | 0             | 0             |  |  |  |
| Not Reported              | 0       | 1 (1)         | 11 (5)        |  |  |  |
| Stroke service            |         |               |               |  |  |  |
| Comprehensive             | 24 (38) | Not Collected | 157 (69)      |  |  |  |
| Primary                   | 31 (49) | Not Collected | 64 (28)       |  |  |  |
| General hospital          | 8 (13)  | Not Collected | 3 (1)         |  |  |  |
| Not Reported              | 0       | Not Collected | 5 (2)         |  |  |  |
| Level of telemedicine use |         |               |               |  |  |  |
| Always                    | 2 (3.2) | 6 (6.5)       | Not Collected |  |  |  |
| Often                     | 8 (13)  | 17 (18)       | Not Collected |  |  |  |
| Sometimes                 | 9 (14)  | 33 (35.5)     | Not Collected |  |  |  |
| Never                     | 43 (68) | 36 (39)       | Not Collected |  |  |  |
| Unknown                   | 1 (1.6) | 0             | Not Collected |  |  |  |
| Not Reported              | 0       | 1 (1)         | Not Collected |  |  |  |
| rt-PA protocol            |         |               |               |  |  |  |
| Yes                       | 62 (98) | 92 (99)       | NA*           |  |  |  |
| No                        | 1 (1.6) | 0 (0.0)       | NA*           |  |  |  |
| Not Reported              | 0       | 1 (1)         | NA*           |  |  |  |

<sup>\*</sup>rt-PA protocols were required to participate in the survey.

Supplement Table B Associations between service related factors and rtPA treatment rates

|  | AUS  |         |              | UK   |         |                | USA   |         |                |
|--|--|---------|--------------|--|---------|----------------|---|---------|----------------|
|  | Mean (SD)                                  | p-value | n            | Mean (SD)                                    | p-value | n              | Mean (SD)                                   | p-value | N              |
| Hospital Setting <sup>1</sup>                          |  |         |              |  |         |                |   |         |                |
| Non-Tertiary   | 7.62 (5.22)                                | 0.120   | 31           | 11.52 (4.77)                                 | 0.046   | 36             | 7.94 (5.72)                                 | 0.005   | 135            |
| Tertiary   | 9.95 (6.22)                                |         | 29           | 13.69 (4.52)                                 |         | 41             | 10.97 (7.68)                                |         | 48             |
| Average door to needle time² <45mins 46-59mins ≥60mins | 15.68 (9.21)<br>9.79 (6.21)<br>8.85 (5.09) | 0.131   | 3<br>9<br>37 | 14.88 (6.18)<br>13.99 (4.00)<br>11.90 (3.14) | 0.067   | 20<br>18<br>28 | 16.51 (7.35)<br>10.75 (7.68)<br>8.07 (5.89) | <0.001  | 7<br>24<br>151 |

<sup>&</sup>lt;sup>1</sup> Independent Sample t-test, <sup>2</sup> One-way ANOVA

## Supplement Table C Factors associated with the addition of non-standard criteria

|  | AUS                  |         | UK                 |         | USA                 |         |  |
|--|----------------------|---------|--------------------|---------|---------------------|---------|--|
|  | N=49                 |         | N=68               |         | N=184               |         |  |
| Variables entered in regression model          | Coefficients         | p-value | Coefficients       | p-value | Coefficients        | p-value |  |
| Hospital Setting                               | -0.29 (-2.84, 2.25)  | 0.817   | -0.83 (-2.39,0.72) | 0.287   | -1.72(-3.25, -0.20) | 0.03    |  |
| Service  | 0.06 (-2.35, 2.48)   | 0.958   | NA                 |         | -0.28(-1.68, 1.10)  | 0.684   |  |
| Admission volume                               | -0.00 ( -0.01, 0.01) | 0.983   | -0.01(-0.04,0.00)  | 0.434   | 0.00 (-0.00, 0.00)  | 0.187   |  |
| Average door to needle time (46mins to 59mins) | -1.33(-5.52, 2.86)   | 0.525   | 1.13(-0.86,3.12)   | 0.262   | 2.51(-1.05, 6.07)   | 0.165   |  |
| Average door to needle time (>60mins)          | 0.60 (-3.34-4.54)    | 0.760   | 1.19(-0.67,3.05)   | 0.207   | 3.57 (0.38 ,6.75)   | 0.023   |  |
| Adherence to 3 hours                           | NA                   |         | NA                 |         | -2.44(-4.30, -0.60) | 0.01    |  |