

Central Lancashire Online Knowledge (CLoK)

Title	Long term effects of gestational hypertension and pre-eclampsia on kidney
	function: Record linkage study
Туре	Article
URL	https://clok.uclan.ac.uk/25033/
DOI	https://doi.org/10.1016/j.preghy.2016.08.231
Date	2016
Citation	Ayansina, D., Black, C., Hall, S. J., Marks, Angharad, Millar, C., Prescott, Gordon, Wilde, K. and Bhattacharya, S. (2016) Long term effects of gestational hypertension and pre-eclampsia on kidney function: Record linkage study. Pregnancy Hypertension: An International Journal of Women's Cardiovascular Health, 6 (4). pp. 344-349. ISSN 2210-7789
Creators	Ayansina, D., Black, C., Hall, S. J., Marks, Angharad, Millar, C., Prescott, Gordon, Wilde, K. and Bhattacharya, S.

It is advisable to refer to the publisher's version if you intend to cite from the work. https://doi.org/10.1016/j.preghy.2016.08.231

For information about Research at UCLan please go to http://www.uclan.ac.uk/research/

All outputs in CLoK are protected by Intellectual Property Rights law, including Copyright law. Copyright, IPR and Moral Rights for the works on this site are retained by the individual authors and/or other copyright owners. Terms and conditions for use of this material are defined in the <u>http://clok.uclan.ac.uk/policies/</u>



Contents lists available at ScienceDirect

Pregnancy Hypertension: An International Journal of Women's Cardiovascular Health



journal homepage: www.elsevier.com/locate/preghy

Long term effects of gestational hypertension and pre-eclampsia on kidney function: Record linkage study



D. Ayansina^a, C. Black^b, S.J. Hall^b, A. Marks^b, C. Millar^c, G.J. Prescott^a, K. Wilde^a, S. Bhattacharya^{d,*}

^a Division of Applied Health Sciences, University of Aberdeen, United Kingdom

^b Chronic Diseases Group, Division of Applied Health Sciences, University of Aberdeen, United Kingdom

^c NHS Grampian, United Kingdom

^d Obstetric Epidemiology, Division of Applied Health Sciences, University of Aberdeen, United Kingdom

ARTICLE INFO

Article history: Received 28 June 2016 Accepted 5 August 2016 Available online 9 August 2016

Keywords: Gestational hypertension Preeclampsia Chronic kidney disease Record linkage

ABSTRACT

Objective: To assess the long term effects of hypertensive disorders of pregnancy on renal function. *Design:* Cohort study where exposure was gestational hypertension or preeclampsia in the first pregnancy. Normotensive women formed the comparison group. *Setting:* Aberdeen, Scotland. *Participants:* All women with date of birth on or before 30th June 1969 and at least their first singleton delivery recorded in the Aberdeen Maternity and Neonatal Databank.

Methods: Participants were linked to the Renal Biochemistry Register, Scottish Morbidity Records, Scottish Renal Registry and National Register for deaths.

Main outcome measures: Occurrence of chronic kidney disease (CKD) as identified from renal function tests in later life, hospital admissions or death from kidney disease or recorded as receiving renal replacement therapy.

Results: CKD was diagnosed in 7.5% and 5.2% of women who previously had GH and PE respectively compared to 3.9% in normotensive women. The unadjusted odds ratio (95% confidence interval) of having CKD in PE was 2.04 (1.53, 2.71) and that for GH was 1.37 (1.15, 1.65), while the adjusted odds ratio (95% confidence interval) of CKD was 1.93 (1.44, 2.57) and 1.36 (1.13, 1.63) in women with PE and GH respectively. Kaplan–Meier curves of survival time to development of chronic kidney disease revealed that women with preeclampsia were susceptible to kidney function impairment earliest, followed by those with gestational hypertension.

Conclusions: There was an increased subsequent risk of CKD associated with hypertensive disorders of pregnancy. Women with GH and PE were also found to have CKD earlier than normotensive women.

© 2016 The Authors. Published by Elsevier B.V. on behalf of International Society for the Study of Hypertension in Pregnancy. This is an open access article under the CC BY-NC-ND license (http://creati-vecommons.org/licenses/by-nc-nd/4.0/).

1. Introduction

Pre-eclampsia, the symptom complex of high blood pressure and proteinuria, affects 4–8% of pregnancies [1]. It is a multisystem disorder and its effects on subsequent cardiovascular health of women [2,3] are well established. Pre-eclampsia is associated with acute kidney function impairment [4,5] with recovery typically occurring within six weeks of delivery of the baby. Relatively little is known about the long term effects on the kidney either in terms of physical damage measured by albuminuria or proteinuria, function impairment measured by reduction in Glomerular Filtration Rate (GFR) or end stage kidney failure.

In their systematic review and meta-analysis, McDonald et al. [6] identified 7 papers reporting on a total of 273 patients with preeclampsia and less than 100 events of microalbuminuria. Despite the limited numbers, they found evidence of an increased risk of microalbuminuria following preeclampsia {pooled risk ratio of 4.3 (95% CI 2.7, 6.9)} but there was considerable heterogeneity among the included studies and a strong possibility of publication bias. There was, however, no difference in terms of serum creatinine levels and estimated Glomerular Filtration Rate (eGFR) between women with preeclampsia and those without. An updated search of the literature identified three additional papers [7–9] all reporting a positive association between preeclampsia

http://dx.doi.org/10.1016/j.preghy.2016.08.231

^{*} Corresponding author at: Dugald Baird Centre for Research on Women's Health, Aberdeen Maternity Hospital, Foresterhill, Aberdeen AB25 2ZL, United Kingdom.

E-mail addresses: d.ayansina@abdn.ac.uk (D. Ayansina), corri.black@abdn.ac.uk (C. Black), colin.millar@nhs.net (C. Millar), k.wilde@abdn.ac.uk (K. Wilde), sohinee. bhattacharya@abdn.ac.uk (S. Bhattacharya).

^{2210-7789/© 2016} The Authors. Published by Elsevier B.V. on behalf of International Society for the Study of Hypertension in Pregnancy. This is an open access article under the CC BY-NC-ND license (http://creativecommons.org/licenses/by-nc-nd/4.0/).

and long term kidney damage, although heterogeneity in terms of exposure and outcome measurement remained as did the possibility of publication bias. In their review, McDonald et al. [6] called for further research investigating the persistence of microalbuminuria following preeclampsia in larger population based studies, taking into consideration confounding factors such as age, BMI, smoking etc.

The overall aim of this research, was to assess the long term effects of hypertensive disorders of pregnancy on kidney function in, using a large phenotypically well-defined population based cohort of women and adjusting for a large number of confounders. In case of declining renal function, we also aimed to find out whether there was a difference in the timing of onset of any decline.

2. Methods

2.1. Data sources and record linkage

The Aberdeen Maternity and Neonatal Databank (AMND), started in 1950, is derived from the Aberdeen Maternity Hospital's obstetric records. Aberdeen Maternity Hospital is the only maternity hospital in Aberdeen and serves as both a primary as well as tertiary care unit. From the AMND, we identified a cohort of women using the following inclusion and exclusion criteria:

2.1.1. Inclusion criteria

Primiparous mothers born on or before the 30th June 1969 (aged 40 years or over at the end date of the study period with a singleton first delivery occurring at or beyond 20 weeks gestation (including still births) and resident in the Grampian region.

2.1.2. Exclusion criteria

women with pre-existing chronic hypertension, kidney disease as recorded in AMND by ICD-9 codes at the time of the first pregnancy; women who had twins or multiple foetuses in their first pregnancy; women who were temporary residents, living in Orkney or Shetland or elsewhere outside of the Grampian region and women with any delivery with a gestation of less than 20 weeks were excluded from the cohort.

The following databases were linked to the AMND to derive information on outcomes for the end of the study follow up period – 31st July 2009:

- (1) The Grampian Renal Biochemistry Dataset (GRBD) is an existing extract of NHS Grampian's Biochemistry Dataset (1999–2009). All biochemistry services (in-patient, community, private) in the Grampian region were provided by two linked National External Quality Assessment Service monitored laboratories. Data were extracted from the GRBD for women in the AMND cohort who:
 - had renal function tested at least one year after the date of delivery
 - were still living in the Grampian region after 1st July 1999
- (2) The Scottish Morbidity Record for non-obstetric and nonpsychiatric general acute inpatient and day case discharges from NHS hospitals in Scotland (SMR01) was introduced in 1961. This was used to identify any hospital admissions from kidney diseases coded according to International Classification of Diseases (10th revision) codes.
- (3) The Scottish Renal Registry (SRR) is a national registry of patients on long term renal replacement therapy. All NHS renal units contribute fully to this database. Data from Scottish renal units were available from 1960 when regular and routine renal replacement therapy (RRT) for end stage renal disease (ESRD) started in Scotland.

(4) National Records of Scotland (NRS) Data on Deaths (formerly the General Register Office for Scotland (GROS)) – Data collection in this register dates back to 1855. This register was used to identify any deaths occurring in the cohort.

The AMND and GRBD datasets were linked by deterministic matching of the Community Health Index (CHI) numbers, numbers unique to patients registered with a general practice in Scotland. CHI numbers of linked women were validated against the vital status register to identify any deaths or migrations out of Grampian region before 1st July 1999. Linkage of the AMND cohort to the Scottish Morbidity Records (SMR01), death registrations (NR Scotland) and the Scottish Renal Registry was carried out by the Information Services Division (ISD) of NHS Scotland using both deterministic matching on CHI numbers and probability matching using surname, initial of first name, date of birth and post code.

2.2. Study design

A cohort study design was used where women who had gestational hypertension (GH) or preeclampsia/ eclampsia (PE) as identified from the AMND formed the exposed cohort, while women who were normotensive (NT) in their first pregnancy constituted the comparison group.

2.3. Definition of exposure

Gestational hypertension and preeclampsia were defined according to the criteria proposed by Davey and MacGillivray [10] and used uniformly throughout the AMND. Gestational hypertension was defined as diastolic pressure greater than 90 mmHg on two occasions at least four hours apart or a single reading of >110 mmHg; from 20 weeks gestation onwards in a previously normotensive woman. Pre-eclampsia was defined as gestational hypertension plus at least one episode of proteinuria of 0.3 g/24 h. Eclampsia was defined clinically as convulsions occurring in the presence of pre-eclampsia. In this analysis, women with preeclampsia and eclampsia were considered together as one group.

2.4. Outcomes

The main outcome of interest was chronic kidney disease defined according to the NICE modified Kidney Disease Outcomes Quality Initiative (KDOQI) definitions for the classification of chronic kidney disease based on renal function measured by glomerular filtration rate (GFR) [11]. Renal function was assessed using serum creatinine values determined by the Jaffe method traceable over time to the international gold standard isotope dilution mass spectrometry (IDMS) method. It was expressed as estimated Glomerular Filtration Rate (eGFR) calculated using the modified 4 variable Modification of Diet in Renal Disease (MDRD) equation suitable for use with IDMS traceable creatinine values. According to these criteria, chronic kidney disease was present if eGFR <60 mL/min/1.73 m² sustained for \ge 90 days. Proteinuria, a measure of renal damage, was defined using urinary protein creatinine ratio (PCR) and albumin creatinine ratio (ACR) as per SIGN guidelines [12] and classified as chronic if sustained for \geq 90 days. We defined women to have CKD if they met either of these criteria in line with KDIGO stage CKD 1-5.

2.5. Statistical analysis

Baseline characteristics were examined for women with GH or PE and compared with those who were normotensive (NT) in their first pregnancy. Results were presented as means (and standard deviation) for continuous variables and frequencies (and percentages) for categorical variables. Pair-wise comparisons were done using independent samples *t*-test for normally distributed continuous variables and chi squared tests for categorical variables as appropriate. Odds ratios for the primary outcomes (having a subsequent renal function test and development of CKD) were



AMND: Aberdeen Maternity and Neonatal Databank

RBD: Renal Biochemistry Database

NRS: National Records of Scotland Data on Deaths

SMR01: Scottish Morbidity Records for General / Acute Inpatient and Day Case

SRR: Scottish Renal Registry records for end stage chronic kidney disease

CKD: Chronic Kidney Disease

calculated using logistic regression and adjusted ratios with 95% confidence intervals were calculated adjusting for age at first delivery, socio-economic status, smoking status, and body mass index (BMI) at first pregnancy.

Time to CKD was examined using survival analysis techniques. Kaplan-Meier plots and log-rank tests were employed as univariate test for between group differences (pair-wise) in survival from CKD. Cox's proportional hazards models were employed to calculate hazard ratios between groups adjusting for age at first delivery, socio-economic status, smoking status, and BMI.

2.6. Ethics Statement

The North of Scotland Research Ethics Service confirmed that formal ethical approval was not necessary to carry out analysis on anonymized data. Permissions to carry out the study were obtained from the steering committee of the Aberdeen Maternity and Neonatal Databank, the NHS Grampian Caldicott guardian, the Privacy Advisory Committee of NHS Scotland and the steering committee for the Scottish Renal Registry.

Table 1

Baseline (at time of delivery) characteristics by exposure status.

3. Results

A total of 14,851 women who had been linked to one or more of the databases recording chronic kidney disease status and had complete information regarding age, socio-economic class, smoking and body mass index (BMI) were included in a multivariate model. Of these, 811 had PE, 3583 had GH and 10,457 were normotensive in their first pregnancy (see Fig. 1).

3.1. Baseline comparisons

Table 1 presents comparison of baseline characteristics at the time of first pregnancy by exposure status. Women with GH or PE had a lower mean age when compared to normotensive women at the time of delivery. Women with GH or PE also had significantly higher mean BMI in comparison with NT women and were more likely to have chronic hypertension, kidney disease, diabetes. Those with chronic hypertension or CKD at baseline were excluded from further analysis. A lower proportion of women with GH and PE were in the higher socioeconomic classes when compared with

Characteristics	Normotensive (n = 10,457)	Gestational hypertension (n = 3583)	p value	Pre-eclampsia (n = 811)	p value
Age, mean (SD) years	25.3 (5.3)	25.0 (5.0)	< 0.001	24.8 (5.3)	<0.001
Socio-economic status, n (%)			< 0.001		0.001
Most affluent I	1088 (10.4)	344 (9.6)		69(8.5)	
II	1694 (16.2)	559 (15.6)		125(15.4)	
III	4821 (46.1)	1727 (48.2)		395 (48.7)	
IV	1955 (18.7)	659(18.4)		154 (19)	
Least affluent V	899 (8.6)	290 (8.1)		68 (8.4)	
Smoking status, n (%)			0.082		0.007
Non smoker	5887 (56.3)	2060 (57.5)		470(57.9)	
Ex smoker	398 (3.8)	129 (3.6)		21 (2.6)	
Current smoker	4172(39.9)	1394 (38.9)		320(39.5)	
BMI, mean (SD) kg/m ²	23.6 (3.6)	24.9 (4.1)	<0.001	25.3 (4.7)	<0.001

*p values represent a test for trend

Table 2

Logistic models with odds ratios for Chronic Kidney Disease; and Survival models with unadjusted and adjusted hazard ratios for Chronic Kidney Disease, hospital admissions and mortality (all cause and cause specific).

	N of events/N (%) Unadjusted	Unadjusted	95%CI for Odds ratio		p value	Adjusted*	95% CI Odds ratio		p value				
		Odds ratio	Lower	Upper		Odds ratio	Lower	Upper					
CKD, n = 14,851 Normotensive Gestational Hypertension Pre-eclampsia	405/10457 (3.9) 188/3583 (5.2) 61/811 (7.5)	1 1.37 2.02	1.15 1.53	1.64 2.67	<0.001 <0.001	1 1.36 1.92	1.14 1.45	1.63 2.56	- 0.001 <0.001				
	N of events/N (%)	Unadjusted	95%CI for Hazard ratio		95%CI for Hazard ratio		95%CI for Hazard ratio		CI for Hazard ratio p value Adjusted ^a		95% CI Hazard ratio		p value
		Hazard ratio	Lower	Upper		Hazard ratio	Lower	Upper					
CKD, n = 14,851 Normotensive Gestational Hypertension Pre-eclampsia Hospital admissions for kidney disease, n = 19,474 Normotensive Gestational Hypertension Pre-eclampsia	405/10457 (3.9) 188/3583 (5.2) 61/811 (7.5) 213/13989 (1.5) 82/4482 (1.8) 24/1003 (2.4)	1 1.31 1.70 1 1.08 1.42	1.10 1.30 0.84 0.93	1.55 2.23 1.40 2.17	- 0.002 <0.001 - 0.541 0.101	1 1.14 1.58 1 1.02 1.37	0.96 1.20 0.78 0.90	1.36 2.07 1.32 2.10	- 0.139 0.001 - 0.890 0.150				
All cause death, n = 25,692 Normotensive Gestational Hypertension Pre-eclampsia Cardiovascular deaths, n = 25,692 Normotensive Gestational Hypertension Pre-eclampsia	562/18574 (3.0) 197/5857 (3.4) 60/1261 (4.8) 52/18574 (0.3) 29/5857 (0.5) 10/1261 (0.8)	1 0.99 1.37 1 1.56 2.53	0.84 1.05 0.99 1.28	1.16 1.79 2.46 4.99	- 0.894 0.020 - 0.056 0.007	1 0.96 1.39 1 1.38 2.33	0.82 1.06 0.87 1.17	1.13 1.81 2.21 4.64	- 0.648 0.017 - 0.170 0.016				

Statistically significant p values are shown as bold.

^a Adjusted for age at delivery, BMI, socio-economic status and smoking status.

NT women. There was no significant difference in the proportion of women in the smoking categories between women with GH and NT women but women with PE had significantly lower proportions of women who had ever smoked (current and ex-smokers) compared with NT women.

3.2. Probability of developing chronic kidney disease

Table 2 presents univariate and multivariate logistic regression models for odds of developing subsequent CKD for women in the three groups with complete information for all covariates in the multivariate model (n = 14,851). Compared with normotensive women, women with GH (OR = 1.37, 95%CI 1.15, 1.64) or PE (OR = 2.02, 95%CI 1.53, 2.67) had statistically significantly higher odds of developing subsequent CKD on univariate examination. Adjusted odds ratios were 1.36 (95%CI 1.14, 1.63), and 1.92 (95% CI 1.45, 2.56), for women with GH and PE respectively (adjusted for age and year at delivery, BMI, socio-economic status and smoking status). Again, in both analyses, there was evidence of a "dose response" with OR greater in PE than GH.

3.3. Timing of onset of kidney function decline

The survival curves shown in Fig. 2 illustrate the time to development of CKD for women in the different exposure groups as first recorded in the RBD. Women were censored at death or, using the more conservative approach (as compared to the previous model), at last measurement if no further measurement of their kidney function was recorded. The log rank p values represent pair-wise comparisons between women with GH or PE and NT women. The survival curves for women with GH or PE were significantly different from that of NT women (p < 0.001 in each case).

Table 2 presents survival models with unadjusted and adjusted hazard ratios for chronic kidney disease, hospital admissions for kidney related conditions and all cause and cause specific mortality. Compared to normotensive women, women with PE had a higher risk of developing CKD {AHR 1.58 (95% CI 1.20–2.07)} of all cause mortality {AHR 1.39 (95% CI 1.06–1.81)} and death from cardiovascular disease {AHR 2.33 (95% CI 1.17–4.64)}. There was no statistically significant difference in hospital admissions from kidney disease. Women with GH did not differ in terms of CKD, hospital admissions from kidney disease and all cause and cause specific mortality from NT women.

4. Discussion

On analysis of a large linked dataset with complete information on co-variates, there appears to be a positive association between gestational hypertension as well as preeclampsia with CKD both in univariate as well as multivariate models. Further, women who had gestational hypertension or preeclampsia were at risk of developing CKD earlier than normotensive women.

The study used high quality data on a large population based cohort analysing approximately 2.5 million person years of follow up with 811 women with preeclampsia . We have also used consistent definitions of exposure and outcomes and incorporated the broadest definition of CKD using both proteinuria and eGFR. The linkage with different registers has enabled cross validation of data. We acknowledge possible selection bias arising out of restricting the study to women born before 1969. We also acknowledge that different follow-up times for the different registers meant that some women's outcome could not be adequately classified. We have already acknowledged the uncertainty in the timing of the first diagnosis of CKD for some women because the biochemistry database only started in 1999. Furthermore, we had no information on those women who remained unlinked to all registers, possibly excluding healthy women from our analysis.

Previous reports including a meta-analysis [6] have consistently found associations between PE and CKD. The few studies that have looked at GH and CKD have mostly reported positive associations [13–15] with the notable exception of Gordin [16], who did not find an increased risk of diabetic nephropathy in women with gestational hypertension. There was considerable variation in the definition of GH, PE and CKD in the literature. Most studies have used microalbuminuria as the outcome measure, both as continuous as well as a dichotomous variable. There were less consistent



Fig. 2. Kaplan-Meier plot showing survival (time to CKD) curves for three study groups.

associations observed between PE and higher levels of protein in the urine (proteinuria: >300 mg/24 h of protein detected in urine), abnormal biochemistry blood tests (elevated creatinine or reduced glomerular filtration rate (GFR)), abnormal renal biopsy or end stage renal disease (ESRD) [7–9,13]. The largest, and most recently published study [9], used record linkage of the Norwegian Birth and Renal Registries to study the association of PE with Renal Replacement Therapy (RRT). They found that PE in the first pregnancy was associated with an increased relative risk for RRT of 4.7 (95% CI 3.6, 6.1). The relative rarity of RRT meant that even a large population based study like this could only identify 477 cases, assessing only the severe end of the spectrum. McDonald's *meta*-analysis [6], identified only 273 patients with preeclampsia and less than 100 events of microalbuminuria across a total of 7 cohort studies. In contrast to the existing literature, our findings suggest that the increase in the risk of developing milder CKD in women with hypertensive disorders of pregnancy is likely to be small. In support of this, Jacquemyn [17] did not find an association between HELLP syndrome and long term renal complications.

5. Conclusions

Our analyses showed a statistically significantly increased relative risk of CKD in women with GH and PE compared to normotensive women with a possible 'dose response effect'. However the absolute risk of CKD in GH and PE was very low (0.7% and 1.3% over 2.5 million person years of follow up). Therefore, the study demonstrates an effect, a biological pathway whereby GH and PE affect renal function in the longer term but the absolute risk of this happening is very low and therefore is probably not of sufficient clinical importance to warrant any intervention or screening. If a hypothetical intervention was put in place that reduced CKD by 50% in women with PE then the numbers needed to treat (NNT) to prevent one CKD event would be around 150 (Inverse of the absolute risk difference). Similarly, for GH, the NNT would be approximately 285 women with GH. CKD occurs slightly earlier in those with GH and PE, but our study had long follow up times and large numbers of participants resulting in statistically significant effects (type 1 error) which is of interest in terms of mechanisms of disease occurrence, but probably of small clinical relevance.

Currently women with GH and PE are reviewed within 2 months postpartum to ensure that hypertension and proteinuria have resolved, but no specific advice is given regarding long term follow up of renal function [18]. There appears to be a small but significant increased risk of CKD in women with preeclampsia. This implies that alongside managing risk of cardiovascular disease in women with PE and GH, there is also a need to consider long term kidney function. The same lifestyle and risk factor modifications should apply – healthy diet, physical activity, maintaining healthy weight, smoking cessation, cholesterol reduction and blood pressure management.

Contributorship

DA analysed the data and wrote the first draft of the paper, EA reviewed the literature and helped with statistical analysis, CB, SB, GP and WCS conceived of the research idea and designed the study, SB was responsible for overall supervision. CB, AM and CM helped with the clinical interpretation of the findings, GJP supervised statistical analysis, KW was responsible for data extraction and linkage, SJH facilitated data linkage. All authors contributed to the writing of the final draft of the paper. SB is the guarantor of this paper.

Ethics statement

Permissions to carry out the study were obtained from the steering committees of the Aberdeen Maternity and Neonatal Databank (SB/SMC/AMND10 dated 08.10.2010), and the Scottish Renal Registry the NHS Grampian Caldicott guardian (dated 20.04.2011), the Privacy Advisory Committee of NHS Scotland (42/11 dated 07/10/2011). Formal ethical approval was not considered necessary by North of Scotland Research Ethics Service as only anonymised data were analysed in this study.

Funding

This research was funded by the Chief Scientists Office, Scotland (CZH/4/659).

Disclosure of interests

The authors declare that they have no conflict of interests

References

- C.B. Rudra, M.A. Williams, Monthly variation in preeclampsia prevalence: Washington State, 1987–2001, J. Matern. Fetal Neonatal Med. 18 (5) (2005) 319–324.
- [2] L. Bellamy, J. Casas, A.D. Hingorani, D.J. Williams, Pre-eclampsia and risk of cardiovascular disease and cancer in later life: systematic review and metaanalysis, Br. Med. J. 335 (7627) (2007) 974–977.
- [3] L.A. Magee, P. Von Dadelszen, Pre-eclampsia and increased cardiovascular risk, Br. Med. J. 335 (7627) (2007) 945–946.
- [4] J. Airoldi, L. Weinstein, Clinical significance of proteinuria in pregnancy, Obstet. Gynecol. Surv. 62 (2) (2007) 117–124.
- [5] A.J. Drakeley, P.A. Le Roux, J. Anthony, J. Penny, Acute renal failure complicating severe preeclampsia requiring admission to an obstetric intensive care unit, Obstet. Gynecol. 186 (2) (2002) 253–256.
- [6] S.D. McDonald, Z. Han, M.W. Walsh, H.C. Gerstein, P.J. Devereaux, Kidney disease after preeclampsia: a systematic review and meta-analysis, Am. J. Kidney Dis. 55 (6) (2010) 1026–1039.
- [7] E. Fiaschi, R. Naccarato, The histopathology of the kidney in toxaemia. Serial renal biopsies during pregnancy, puerperium and several years postpartum – light and electron microscopic and immunofluorescent studies, in: Virchows Archiv. A Pathol. Anat. 345 (4) (1968) 299–309.
- [8] S. Murakami, M. Saitoh, T. Kubo, T. Koyama, M. Kobayashi, Renal disease in women with severe preeclampsia or gestational proteinuria, Obstet. Gynecol. 96 (6) (2000) 945–949.
- [9] B.E. Vikse, L.M. Irgens, T. Leivestad, R. Skjærven, B.M. Iversen, Preeclampsia and the risk of end-stage renal disease, N. Engl. J. Med. 359 (8) (2008) 800–809.
- [10] D.A. Davey, I. MacGillivray, The classification and definition of the hypertensive disorders of pregnancy, Am. J. Obstet. Gynecol. 158 (4) (1988) 892–898.
- [11] National Collaborating Centre for Chronic Conditions, Chronic Kidney Disease: National Clinical Guideline for Early Identification and Management in Adults in Primary and Secondary Care, Royal College of Physicians, London, 2008.
- [12] Scottish Intercollegiate Guidelines Network, Diagnosis and Management of Chronic Kidney Disease, SIGN, Edinburgh, UK, 2008. Report No.: 103.
- [13] H. Nisell, H. Lintu, N.O. Lunell, G. Mollerstrom, E. Pettersson, Blood pressure and renal function seven years after pregnancy complicated by hypertension, Br. J. Obstet. Gynaecol. 102 (11) (1995) 876–881.
- [14] R.A. North, D. Simmons, D. Barnfather, M. Upjohn, What happens to women with preeclampsia? Microalbuminuria and hypertension following preeclampsia, Aust. N. Z. J. Obstet. Gynaecol. 36 (3) (1996) 233–238.
- [15] A.G. Shammas, J.F. Maayah, Hypertension and its relation to renal function 10 years after pregnancy complicated by pre-eclampsia and pregnancy induced hypertension, Saudi Med. J. 21 (2) (2000) 190–192.
- [16] D. Gordin, V. Hiilesmaa, J. Fagerudd, M. Rönnback, C. Forsblom, R. Kaaja, et al., Pre-eclampsia but not pregnancy-induced hypertension is a risk factor for diabetic nephropathy in type 1 diabetic women, Diabetologia 50 (3) (2007) 516–522.
- [17] Y. Jacquemyn, L. Jochems, E. Duiker, J. Bosmans, V. Van Hoof, C. Van Campenhout, Long-term renal function after HELLP syndrome, Gynecol. Obstet. Invest. 57 (2) (2004) 117–120.
- [18] Chronic kidney disease: early identification and management of chronic kidney disease in adults in primary and secondary care: Guideline no 73. Available at: <<u>http://guidance.nice.org.uk/CG73/NICEGuidance/pdf/English</u>>, (Accessed 03/15, 2013).