

Central Lancashire Online Knowledge (CLoK)

Title	Is Black Carbon an Unimportant Ice-Nucleating Particle in Mixed-Phase Clouds?
Type	Article
URL	https://clock.uclan.ac.uk/30829/
DOI	https://doi.org/10.1002/2017JD027831
Date	2018
Citation	Vergara-Temprado, Jesús, Holden, Mark, Orton, Thomas R., O'Sullivan, Daniel, Umo, Nsikanabasi S., Browse, Jo, Reddington, Carly, Baeza-Romero, María Teresa, Jones, Jenny M. et al (2018) Is Black Carbon an Unimportant Ice-Nucleating Particle in Mixed-Phase Clouds? Journal of Geophysical Research: Atmospheres, 123 (8). pp. 4273-4283. ISSN 2169-897X
Creators	Vergara-Temprado, Jesús, Holden, Mark, Orton, Thomas R., O'Sullivan, Daniel, Umo, Nsikanabasi S., Browse, Jo, Reddington, Carly, Baeza-Romero, María Teresa, Jones, Jenny M., Lea-Langton, Amanda, Williams, Alan, Carslaw, Ken S. and Murray, Benjamin J.

It is advisable to refer to the publisher's version if you intend to cite from the work.
<https://doi.org/10.1002/2017JD027831>

For information about Research at UCLan please go to <http://www.uclan.ac.uk/research/>

All outputs in CLoK are protected by Intellectual Property Rights law, including Copyright law. Copyright, IPR and Moral Rights for the works on this site are retained by the individual authors and/or other copyright owners. Terms and conditions for use of this material are defined in the <http://clock.uclan.ac.uk/policies/>

RESEARCH ARTICLE

10.1002/2017JD027831

J. Vergara-Temprado is the corresponding author for questions related to the modelling section and M.A. Holden for the experimental part of this study. Both authors contributed equally to this work.

Key Points:

- Soot particles generated from two different fuels do not nucleate ice above the handling blanks when immersed in water droplets
- Previous representations of the ice-nucleating ability of BC particles overestimate its importance in the atmosphere
- We use a global aerosol model to estimate the required ice-nucleating efficiency of BC for it to compete with other INP in the atmosphere

Supporting Information:

- Supporting Information S1

Correspondence to:

J. Vergara-Temprado and M. A. Holden,
jesus.vergara@env.ethz.ch;
m.a.holden@leeds.ac.uk

Citation:

Vergara-Temprado, J., Holden, M. A., Orton, T. R., O'Sullivan, D., Umo, N. S., Browse, J., et al. (2018). Is black carbon an unimportant ice-nucleating particle in mixed-phase clouds? *Journal of Geophysical Research: Atmospheres*, 123, 4273–4283. <https://doi.org/10.1002/2017JD027831>

Received 2 OCT 2017

Accepted 21 MAR 2018

Accepted article online 30 MAR 2018

Published online 26 APR 2018

Is Black Carbon an Unimportant Ice-Nucleating Particle in Mixed-Phase Clouds?

Jesús Vergara-Temprado^{1,2} , Mark A. Holden^{1,3} , Thomas R. Orton^{1,4}, Daniel O'Sullivan¹,
Nsikanabasi S. Umo^{1,5}, Jo Browse^{1,6}, Carly Reddington¹ , María Teresa Baeza-Romero⁷ ,
Jenny M. Jones⁸, Amanda Lea-Langton^{8,9} , Alan Williams⁸, Ken S. Carslaw¹ ,
and Benjamin J. Murray¹ 

¹Institute for Climate and Atmospheric Science, School of Earth and Environment, University of Leeds, Leeds, UK, ²Now at Institute for Atmospheric and Climate Science, ETH Zurich, Zurich, Switzerland, ³School of Chemistry, University of Leeds, Leeds, UK, ⁴Now at Lloyd's of London, London, UK, ⁵Now at Institute for Meteorology and Climate Research-Atmospheric Aerosol Research, Karlsruhe Institute of Technology, Eggenstein-Leopoldshafen, Germany, ⁶Now at School of Geography, University of Exeter, Penryn, Cornwall, UK, ⁷Escuela de Ingeniería Industrial de Toledo, Universidad de Castilla-La Mancha, Toledo, Spain, ⁸School of Chemical and Process Engineering, University of Leeds, Leeds, UK, ⁹Now at School of Mechanical, Aerospace and Civil Engineering, University of Manchester, Manchester, UK

Abstract It has been hypothesized that black carbon (BC) influences mixed-phase clouds by acting as an ice-nucleating particle (INP). However, the literature data for ice nucleation by BC immersed in supercooled water are extremely varied, with some studies reporting that BC is very effective at nucleating ice, whereas others report no ice-nucleating ability. Here we present new experimental results for immersion mode ice nucleation by BC from two contrasting fuels (*n*-decane and eugenol). We observe no significant heterogeneous nucleation by either sample. Using a global aerosol model, we quantify the maximum relative importance of BC for ice nucleation when compared with K-feldspar and marine organic aerosol acting as INP. Based on the upper limit from our laboratory data, we show that BC contributes at least several orders of magnitude less INP than feldspar and marine organic aerosol. Representations of its atmospheric ice-nucleating ability based on older laboratory data produce unrealistic results when compared against ambient observations of INP. Since BC is a complex material, it cannot be unambiguously ruled out as an important INP species in all locations at all times. Therefore, we use our model to estimate a range of values for the density of active sites that BC particles must have to be relevant for ice nucleation in the atmosphere. The estimated values will guide future work on BC, defining the required sensitivity of future experimental studies.

Plain Language Summary The ability of black carbon particles to trigger the formation of ice from liquid cloud droplets is assessed by combining laboratory measurement of its efficiency with a global aerosol model. The new estimates suggest that these particles are not as efficient for ice nucleation when immersed in water droplets as previously thought and therefore not as atmospherically relevant.

1. Introduction

Black carbon (BC) particles emitted from both anthropogenic and natural combustion processes are ubiquitous in the present-day atmosphere with an estimated total emission rate of 7.5 Tg/year (Bond et al., 2013). It is estimated that the anthropogenic emissions of BC have increased from ~1 Tg/year in 1850 to ~5 Tg/year in 2000 (Lee et al., 2013), which is thought to have led to a significant impact on climate (Bond et al., 2013). BC has a strong warming effect through the absorption of solar and infrared radiation, and it has been suggested that reduction in BC emissions might go some way to mitigating global warming (Bond et al., 2013). However, to accurately assess the efficacy of reducing BC emissions, it is important to quantify the impacts of BC aerosol on clouds. It is estimated that BC particles contribute substantially to global cloud condensation nuclei concentrations, and they are an important cloud condensation nuclei in industrial regions (Spracklen et al., 2011). BC therefore influences the albedo and lifetime of clouds through nucleating cloud droplets. If these immersed particles could also nucleate ice effectively, then the lifetime and albedo of supercooled clouds would be affected. This “glaciation indirect effect,” which would most likely enhance precipitation and reduce cloud lifetime, could potentially offset the aerosol effects on liquid clouds (Lohmann, 2002, 2017). However, the ice-nucleating ability of BC under conditions pertinent to supercooled clouds remains very uncertain.

©2018. The Authors.

This is an open access article under the terms of the Creative Commons Attribution License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

While it has been shown in laboratory studies that BC nucleates ice under conditions relevant for cirrus clouds (Hoose & Möhler, 2012; Kanji et al., 2011; Koehler et al., 2009; Kulkarni et al., 2016; Möhler, 2005; Ullrich et al., 2017), there are divergent results from laboratory and field studies of the ability of BC to nucleate ice under water-saturated conditions, which are relevant for mixed-phase clouds (DeMott, 1990; Diehl & Mitra, 1998; Hoose & Möhler, 2012; Schill et al., 2016; Ullrich et al., 2017). For example, a strong correlation between BC and the ice crystal concentration in a mixed-phase orographic mountain wave cloud suggested that BC might nucleate ice (Twohy et al., 2010). However, in the same field campaign BC was not enhanced significantly in the ice crystal residues over the background air (Pratt et al., 2009). In one study of mixed-phase clouds at a high-altitude observatory in the Alps, soot particles only made up 5% of the submicron aerosol particles, but 27% of ice crystal residues (Cozic et al., 2008). In contrast, several studies found that BC accounted for only a minor fraction of ice crystal residues, but mineral dust was clearly enhanced (Baustian et al., 2012; Kamphus et al., 2010; Kupiszewski et al., 2016; Schmidt et al., 2017). More recently, analysis of ice-nucleating particle (INP) chemical composition in air influenced by biomass burning events using electron microscopy showed that between 0% and 64% of INPs were BC particles and suggested that biomass burning particles could be an important regional source of INP, especially during periods when other INPs such as desert dust are absent (Mccluskey et al., 2014). Some other studies have observed an enhancement in the INP concentration in biomass burning aerosols (Levin et al., 2016; Petters et al., 2009). For cases suggesting a role of BC INPs, it is not clear if it is the BC component of these aerosols that causes nucleation or some other components mixed with it. For example, Levin et al. (2016) removed refractory BC via laser-induced incandescence observing a decrease in the concentration of INPs for some of their samples. However, the ice-nucleating ability of kaolinite internally mixed with BC was affected when the refractory BC part was removed by this method, which suggest that the ice-nucleating ability of other aerosol components mixed with BC could also be affected by the removal of the refractory BC component. Measurements of the ice-nucleating efficiencies of BC particles from diesel engines (Schill et al., 2016) found that fresh and photochemically aged BC particles did not nucleate ice effectively above their limit of detection, defined by the number of background nucleation events in their instrument. Overall, the field results do not clarify whether BC particles are consistently playing a role as INPs in the atmosphere, and they suggest that BC (or compounds generated and transported along with BC) might be playing a sporadic role in nucleating ice under certain atmospheric conditions.

The available data from laboratory studies also leave open the question of whether or not BC is an efficient INP. Some studies show evidence that BC can nucleate ice in supercooled droplets (Brooks et al., 2014; DeMott, 1990; Diehl & Mitra, 1998; Gorbunov et al., 2001; Popovicheva et al., 2008; Wright et al., 2013). Based on the available literature data (DeMott, 1990; Diehl & Mitra, 1998), a parameterization of the density of ice-nucleating active sites (n_s) was derived and, in combination with BC concentrations in the atmosphere, it was suggested that BC is a very important INP type (Murray et al., 2012). However, more recent studies could not reproduce similar values of n_s for BC, and the upper limits estimated from the limits of detection of the instruments were orders of magnitude lower (Schill et al., 2016; Ullrich et al., 2017). The large variation in BC ice nucleation activity reported in these studies means that the contribution of BC to a possible anthropogenic glaciation effect has remained poorly quantified, since modeling results will depend strongly on the represented ability of BC for nucleating ice under mixed-phase conditions (Fan et al., 2012; Hoose, Kristjánsson, Chen, & Hazra, 2010; Hoose, Kristjánsson, & Burrows, 2010; Phillips et al., 2008, 2013; Savre & Ekman, 2015; Wang et al., 2014; Yun & Penner, 2012).

2. Results and Discussion

2.1. Laboratory Study of Ice Nucleation by Soot Samples

Given the large variation in ice-nucleating activities reported for the relatively few experimental studies of the ice-nucleating ability of BC, we have made new laboratory measurements. We have taken great care in these experiments to characterize the background INP, which inevitably contaminate experiments such as these, but which can lead to a false ice nucleation signal. We have also taken care to generate BC samples in a reproducible and well-characterized way.

For these experiments, we have generated BC particles from the incomplete combustion of liquid fuels. By definition (Petzold et al., 2013), these laboratory-generated BC samples should be referred to as soot. Much of the BC in the atmosphere originates from incomplete combustion, but on transport through the




Figure 1. Transmission electron microscopy images of eugenol and *n*-decane soot that were generated in the laboratory for this study. Eugenol soot images are shown on the top plates (a and b), while the plates below (c and d) are *n*-decane soot images. The scale of each image is labeled at the bottom left-hand corner of the image.

atmosphere it is expected to evolve through the adsorption of other chemical species, reactions with gas phase constituents, and aggregation with other aerosol particles; hence, this atmospheric material is then generally termed as BC. In these experiments, we consider fresh soot particles generated in our laboratory as a proxy for atmospheric BC.

The fuels used to generate the soots for this study were a proxy for hydrocarbon combustion (*n*-decane, $C_{10}H_{22}$), and a proxy for biomass burning (eugenol, $C_{10}H_{12}O_2$). Eugenol is used as a proxy for the combustion of lignin, which constitutes 20% of pinewood (Fitzpatrick et al., 2008). Lignin has previously been shown to contribute to soot production in biomass burning, alongside cellulose (Fitzpatrick et al., 2008; J. M. Wilson et al., 2013). Consequently, soot from eugenol is similar in composition to that from pine combustion (Baeza-Romero et al., 2010). We used the same methods to produce soot from *n*-decane and eugenol, a wick diffusion burner with filtered air, as described in previous studies where the soot was characterized by mass spectrometry (Baeza-Romero et al., 2010; J. M. Wilson et al., 2013; the methodology is described in detail in supporting information S1). For our experiments, soot was collected on glass slides at the top of a glass chimney and both fuels led to soot spherules with the classic fractal soot morphology (see transmission electron microscopy images, Figure 1). While they are morphologically similar, the soot from these fuels differ in several ways. For example, soot from eugenol contains larger oxygenated polyaromatic hydrocarbons, has a greater oxygen content and has a lower elemental carbon:total carbon ratio than soot from *n*-decane (Baeza-Romero et al., 2010; Fitzpatrick et al., 2008). Hence, we produced two contrasting examples of soot, both of which are thought to be relevant for the atmosphere.

Water suspensions of soot were prepared at 10^{-3} wt%. This is a lower mass ratio compared to similar micro-litre experiments performed on mineral dusts and was selected to avoid significant aggregation of the soot, since this would lead to poor dispersion in suspension and introduce additional uncertainties into the ice nucleation measurements. Laser diffraction using a Malvern Mastersizer 2000E instrument was used to




Figure 2. (a) Fraction frozen curves of our experiments for the two different BC samples generated from eugenol and *n*-decane with the baseline of our experiments defined by the representative handling blanks (dark blue). The results shown are for the experiments with 10⁻³ wt%. (b) Upper limit of density of active sites that the studied BC particles can have. The errors have been estimated as the Poisson counting errors following the method presented in Harrison et al. (2016). Other parameterizations from the literature are shown for comparison. The upper limit is parameterized with the following equation: $n_s \text{ (cm}^{-2}\text{)} = \exp(-6.608 - 0.419 \times T(\text{°C}))$ valid in the temperature range -30 to -12°C.

assess the particle size distributions and aggregation, the results of which are shown in the supporting information. The mastersizer records the angular intensity of scatter laser light and then estimates the particle size distribution in terms of volume equivalent spheres using Mie Theory. At 10⁻¹ wt%, about 50% of the soot surface area is associated with particles >1 μm in diameter, compared with 0–20% of the surface area at the concentrations used in this study. For even more hydrophobic soots, such as acetylene burner soot, it was not possible to produce suspensions. Microlitre droplets of these suspensions were cooled down to test their ice-nucleating efficiency. Detailed discussion of the technique can be found in Whale et al. (2014), so only a short description is presented here. Briefly, this droplet freezing assay consists of an array of droplets (between 30 and 44 droplets per experiment), pipetted on to a silanized slide (Hampton Scientific), which is cooled down at a ramp rate of -1°C/min using a stirling engine chiller. Here the active site density is used (Connolly et al., 2009), which is a singular time-independent description of ice nucleation. For a population of N droplets, the fraction of frozen droplets (ff) is calculated as $ff(T) = n(T)/N$ where $n(T)$ is number of droplets frozen at a temperature T . The density of active sites n_s can be calculated from the fraction frozen as

$$n_s(T) = -\frac{\ln(1 - ff(T))}{A} \quad (1)$$

where A is the surface area of BC particles per droplet determined by Brunauer-Emmett-Teller (BET) multilayer adsorption. For eugenol, the specific surface area was $49.43 \pm 0.89 \text{ m}^2/\text{g}$, while for *n*-decane soot it was $65.47 \pm 0.81 \text{ m}^2/\text{g}$.

The resulting fraction of droplets frozen as a function of temperature are shown in Figure 2a for experiments with and without soot in the droplets. In all the experiments conducted, we did not measure any significant increase in ice nucleation activity above the handling blanks when soot particles were present. In these experiments the handling blanks did not freeze at the homogeneous limit but instead froze heterogeneously. This is caused by the contact between the water and the hydrophobic glass slides, or by trace contaminants. The handling blanks were MilliQ water samples run alongside the soot suspensions, reproducing every process, including sonication and stirring, that the soot suspensions were exposed to. These handling blanks froze over a broader range than standard MilliQ blanks and included freezing at warmer temperatures caused by the introduction of impurities. Given that there is no significant difference in freezing temperatures between the soot samples and handling blanks, it is not possible to ascribe the freezing temperatures measured to the influence of soot alone; instead, the freezing could either be entirely unrelated to the soot or a convolution of the soot and other contaminants acting as INPs. These results are similar to the observations of Schill et al. (2016) and Ullrich et al. (2017), where no significant ice nucleation ability of BC was observed. However, while no significant activity of these soots has been measured, they do define a

limiting freezing efficiency, which we can use to draw conclusions about the potential of BC to contribute to the population of atmospheric INP.

The upper limit of the density of active sites that our soot samples can be estimated from our fraction frozen results by assuming that all the nucleation events that we observed were produced by our soot particles. The real value of n_s is likely to be lower, as the freezing events observed cannot be distinguished from the handling blanks. The density of active sites is then calculated using equation (1) with the BET-specific surface areas of dry soot. A simple parameterization of this upper limit calculated by fitting the values to a line in logarithmic scale is given in the figure caption (see Figure 2b). The upper limit defined here (NEW-UPL) gives smaller values of the upper limits of n_s than previously reported by Ullrich et al. (2017), but similar to that defined by Schill et al. (2016). All three of the recent upper limit parameterizations (this study, Schill et al., 2016; Ullrich et al., 2017) are at least 3 orders of magnitude smaller than the old parameterization defined by Murray et al. (2012) on the basis of literature data from DeMott (1990) and Diehl and Mitra (1998). DeMott (1990) used an expansion chamber to study the ice-nucleating ability of soots generated by a acetylene burner, whereas Diehl and Mitra (1998) froze droplets containing soot particles from kerosene burner exhaust by injecting them into a wind tunnel at various temperatures.

Given the discrepancy in the observed values of n_s from different studies (Figure 2b), one cannot readily conclude what the typical ice-nucleating activity of BC particles is in the atmosphere. These observed differences in BC n_s could be due to structural differences in various types of BC from different fuels or be related to processes affecting the BC ice-nucleating ability such as other materials being adsorbed to the soot during production, such as organic carbon species/polyaromatic hydrocarbons. Atmospheric aging could also change the ice-nucleating properties of BC particles as organic or sulfate coatings could potentially affect its ice-nucleating ability (Kulkarni et al., 2016). Therefore, soot particles might have different ice-nucleating abilities depending on their properties and sources, so assuming a single distribution of n_s values at all times and locations might misrepresent its ability as an INP in the atmosphere. However, an exploratory study of its atmospheric potential as INP can be done by comparing its potential contribution to global INP concentrations with that of other well-characterized INP species.

2.2. Modeling the Contribution of BC INP to the Global Atmospheric Burden on INP

To estimate the possible contribution of BC particles to the global distribution of INP, we use the global aerosol model GLObal Model of Aerosol Processes (GLOMAP) mode, as used in Vergara-Temprado et al. (2017, hereafter VT17). We estimate the concentrations in the atmosphere of two well-known ice-nucleating aerosol species, K-feldspar (Atkinson et al., 2013) and marine organic aerosols (T. W. Wilson et al., 2015) and compare these to the predicted contribution of BC INP. We can consider that for any aerosol species to be relevant in the atmosphere as an INP, it will have to produce similar or greater concentrations to the simulated INP concentrations of K-feldspar and marine organics (Figure 3). Below these concentrations, it is unlikely to be an important INP; however, higher concentrations would only indicate that this is potentially more important than the two species modeled (in some locations other INP types may also be important). In our model, BC is emitted from wildfires that vary seasonally (Van der Werf et al., 2003), fossil fuel, and biofuel emissions as described in Mann et al. (2010). The annual-mean fluxes are defined by Bond (2004). BC is emitted internally mixed with organic carbon into the insoluble Aitken mode, and then it is moved to the soluble modes by atmospheric aging. The transport, mixing, and scavenging of BC particles are driven by the meteorology of the year 2001 as used in VT17. A more detailed description of the model is given in the supporting information (see Text S2). The model simulated BC concentrations are evaluated using several data sets from the Global Aerosol Synthesis and Science Project repository presented in Reddington et al. (2017; see Text S2; Barth et al., 2015; Harris et al., 2017; Leon et al., 2016; Liu et al., 2015; Matsui et al., 2011; McMeeking et al., 2010, 2011; Metcalf et al., 2012; O'Shea et al., 2014; Oshima et al., 2012; Perring et al., 2017; Ryerson et al., 2013; Schutgens et al., 2016, 2017; Schwarz et al., 2008, 2010; Spackman et al., 2010; Subramanian et al., 2010; Taylor et al., 2014).

With the BC concentrations simulated, we can calculate the BC INP concentration for a given $n_s(T)$ following the method shown in VT17. As the reported values of n_s range by several orders of magnitude, we define two limiting parameterizations, one using the upper limit presented in this study (NEW-UPL) and another using




Figure 3. Ice-nucleating particles (INPs) concentrations from black carbon (BC) particles and the simulated concentrations in VT17 (using feldspar and marine organic aerosols). The contour lines show $[INP]_{-25}$ from VT17, and the color maps show the same values simulated when using BC INP calculated with (a) NEW-UPL and (b) OLD. A similar figure for an activation temperature of -30°C is given in the supporting information.

the maximum observed values reported in literature data corresponding to the parameterization shown in Murray et al. (2012; OLD; Figure 2), which was based on data from DeMott (1990) and Diehl and Mitra (1998).

The INP distribution at the Earth's surface is shown at an activation temperature of -25°C (i.e., $[INP]_{-25}$, where square brackets indicate concentration); that is, the number of particles that would nucleate ice if exposed to this temperature in a mixed-phase cloud (Figure 3). These surface-level plots are useful for assessing the distribution and makeup of the INP population around the globe, but they do not tell us where INPs can nucleate ice and influence clouds. To show this, we calculated the INP concentration throughout the atmosphere using local ambient temperatures and particle concentrations. The annual mean $[INP]_{\text{ambient}}$ plotted in Figure 4 was calculated by averaging the daily $[INP]_{\text{ambient}}$ values as the daily variations in temperature can substantially affect the simulated mean concentrations. At temperatures below the minimum temperature limit of each parameterization, we use the value of n_s for the lowest experimental temperature reported to avoid extrapolating the parameterizations.

In both the $[INP]_{-25}$ (Figure 3) and $[INP]_{\text{ambient}}$ plots (Figure 4), when the NEW-UPL n_s values are used, the concentrations of INP from BC are several orders of magnitude smaller than simulated assuming K-feldspar and marine organic aerosols, which suggests that BC is unlikely to play an influential role as an INP on global scales if the soot we generated is representative of atmospheric BC. However, when the OLD parameterization for n_s is used, BC completely dominates the global INP distribution both for $[INP]_{\text{ambient}}$ and $[INP]_{-25}$.

Given this large difference, we estimate the n_s values required for BC to be an important INP type given present-day BC emissions by calculating the percentage of surface grid boxes that would be dominated by BC particles if they were to have a particular n_s value (Figure 5) over the mixed-phase temperature range. This is done by calculating BC $[INP]$ from the simulated surface area distribution of BC for a range of n_s values (from 10^{-5} to 10^9 cm^{-2}). We then calculate at each temperature the fraction of grid boxes in our model where the BC $[INP]$ exceeds the INP concentrations simulated by VT17, weighting each grid box by the geographic surface area that it represents. This is done for surface level aerosol concentrations (Figure 5a) and for a pressure level of 600 hPa (Figure 5b). Figure 5b predicts the range where BC would start to compete with the other species to be narrower than at surface level because of the larger homogeneity of aerosol




Figure 4. Zonal mean profiles of the ratio between the number of black carbon (BC) ice-nucleating particles (INPs) at local ambient temperature ($[INP]_{\text{ambient}}$) and the concentrations as simulated in VT17 for marine organics and K-feldspar. (a) Using NEW-UPL for calculating BC $[INP]_{\text{ambient}}$ and (b) using OLD.

concentrations once they are transported and mixed in the atmosphere. This approach helps us to place the other literature data for ice nucleation by BC in context and will similarly help place any future measurements of the ice nucleation ability of BC in context. The OLD parameterization and those presented by Phillips et al. (2008, 2013) fall in the range of values where BC would dominate the surface INP concentrations by orders of magnitude. On the other hand, the NEW-UPL produces values that are orders of magnitude lower than the minimum required to influence surface concentrations. Previously reported upper limits are also lower than the required n_s values necessary to influence surface concentrations. We note that the model presented in VT17 is subject to low biases at high temperatures ($< -15^\circ\text{C}$) in continental regions, produced probably by the absence of other terrestrial ice-nucleating aerosols in the model, so the results at these temperatures should be interpreted accordingly.

We then test the realism of the OLD and NEW-UPL parameterizations against ambient INP observations by evaluating the simulated INP concentrations against two contrasting atmospheric INP data sets, one based on remote marine locations in the Southern Ocean (Bigg, 1973) and another in a relatively higher BC loaded environment from several places in China (Yin et al., 2012; Figures 6a and 6b). The simulated concentrations of INP are calculated by applying either the NEW-UPL or the OLD parameterization to the annual mean BC concentrations simulated at the locations where the campaigns took place. We also show for comparison the prediction with VT17. When the NEW-UPL is used, BC alone cannot explain the observed INP concentrations and underrepresents the atmospheric concentrations by more than an order of magnitude, suggesting that other species are responsible for producing these INP concentrations. On the other hand, when the OLD parameterization is used, the measured concentrations in both environments are overestimated by more




Figure 5. (a) Percentage of the globe surface area that would be dominated by black carbon (BC) particles at each temperature as a function of BC n_s , when compared with the sum of ice-nucleating particle (INP) produced from marine organics and K-feldspar (from VT17). This plot is for INP at the surface. (b) Same as (a) but for a pressure level of 600 hPa. The parameterizations labeled Phillips et al. (2008) and Phillips et al. (2013) were derived using the approach given by Hoose and Möhler (2012).




Figure 6. (a) Comparison between the simulated values of black carbon ice-nucleating particles (INPs) when using NEW-UPL and OLD parameterization, and observed INP concentrations from different places in China (Yin et al., 2012). (b) Same as (a) but for the Southern Ocean (Bigg, 1973). The comparison with the sum of marine organics and K-feldspar (from VT17) is also shown for comparison in both panels.

than 2 orders of magnitude, suggesting that if we consider all atmospheric BC particles to act with this efficiency in the atmosphere, we will likely overestimate their influence as INPs. This conclusion is in agreement with many field observations, which suggest that mineral dust is the dominant aerosol found in ice crystals residues (Baustian et al., 2012; Kamphus et al., 2010; Pratt et al., 2009; Schmidt et al., 2017) although others did show that BC aerosols might contribute to the observed INP concentration (Cozic et al., 2008; Mccluskey et al., 2014). K-feldspar and marine organic aerosols can explain these atmospheric concentrations within an order of magnitude, as shown previously in Vergara-Temprado et al. (2017). Hence, we conclude that the OLD parameterization is probably unrealistic and that it is not possible that all atmospheric BC has such a high ice-nucleating efficiency. This conclusion has important implications for modeling studies that have previously treated BC as INP in the immersion mode (Fan et al., 2012; Hoose, Kristjánsson, Chen, & Hazra, 2010; Hoose, Kristjánsson, & Burrows, 2010; Phillips et al., 2008, 2013; Savre & Ekman, 2015; Wang et al., 2014; Yun & Penner, 2012). However, we cannot completely dismiss the potential influence that BC produced from different fuels, or exposed to different conditions, might have on INPs regionally, or during exceptional events such as large biomass burning events.

3. Conclusions

Our modeling estimates suggest that if all BC particles in the atmosphere behave as reported in this study, and by several other recent studies (Schill et al., 2016; Ullrich et al., 2017), BC is unlikely to play a substantial atmospheric role as INPs through the immersion mode in mixed-phase clouds. We also conclude that a representation of BC INPs from Murray et al. (2012), which was based on the studies of DeMott (1990) and Diehl and Mitra (1998), results in an overestimation of surface-level concentrations of INPs in remote and polluted environments by more than 2 orders of magnitude compared to observations.

The question of whether the ice-nucleating ability of these studied BC particles is representative of the ice-nucleating ability of atmospheric BC particles globally at all times remains open, since the discrepancies between various studies cannot be currently explained. Furthermore, we cannot rule out that atmospheric BC particles could be affected by processes enhancing their ice-nucleating ability to levels that could make them relevant regionally or sporadically. Hence, we suggest that more studies to clarify the sources of discrepancies in the laboratory datasets are necessary to either quantify the effect of BC as INPs in the atmosphere or rule out its relevance completely. Specifically, experiments with contemporary techniques where special attention is paid to characterizing and controlling impurities need to be done with the specific BC types used in previous studies where soot was found to be an effective ice-nucleating material. Nevertheless, we recommend that the old parameterizations, such as Murray et al. (2012), should not be used to describe the ice-nucleating ability of all soot in the atmosphere. Overall, the available evidence suggests that BC is at most, of second-order importance when compared to other ice-nucleating aerosol types such as mineral dust or marine organics.

Acknowledgments

This study has been funded by the European Union's Seventh Framework Programme (FP7/2007-797 2013) under grant agreement 603445 (BACCHUS), the European Research Council (ERC, 240449 ICE and 648661 Marinelce), and the Engineering and Physical Sciences Research Council (EPSRC, EP/M003027/1). The global model simulations were performed on the ARCHER UK National Supercomputing Service. Ken S. Carslaw is a Royal Society Wolfson Merit Award holder. Nsikanabasi Umo acknowledges funding from the Niger Delta Development Commission (NDDC) in Nigeria (NDDC/DEHSS/ 2010PGFS/AK/ 011) and the Alexander von Humboldt Foundation, Germany (1188375). All the data used in this manuscript can be accessed at <https://doi.org/10.5518/310>.

References

- Atkinson, J. D., Murray, B. J., Woodhouse, M. T., Whale, T. F., Baustian, K. J., Carslaw, K. S., et al. (2013). The importance of feldspar for ice nucleation by mineral dust in mixed-phase clouds. *Nature*, 498(7454), 355–358. <https://doi.org/10.1038/nature12278>
- Baeza-Romero, M. T., Wilson, J. M., Fitzpatrick, E. M., Jones, J. M., & Williams, A. (2010). In situ study of soot from the combustion of a biomass pyrolysis intermediate—Eugenol—and n-decane using aerosol time of flight mass spectrometry. *Energy & Fuels*, 24(1), 439–445. <https://doi.org/10.1021/ef9008746>
- Barth, M. C., Cantrell, C. A., Brune, W. H., Rutledge, S. A., Crawford, J. H., Huntrieser, H., et al. (2015). The deep convective clouds and chemistry (DC3) field campaign. *Bulletin of the American Meteorological Society*, 96(8), 1281–1309. <https://doi.org/10.1175/BAMS-D-13-00290.1>
- Baustian, K. J., Cziczo, D. J., Wise, M. E., Pratt, K. A., Kulkarni, G., Hallar, A. G., & Tolbert, M. A. (2012). Importance of aerosol composition, mixing state, and morphology for heterogeneous ice nucleation: A combined field and laboratory approach. *Journal of Geophysical Research*, 117, D06217. <https://doi.org/10.1029/2011JD016784>
- Bigg, E. K. (1973). Ice nucleus concentrations in remote areas. *Journal of the Atmospheric Sciences*, 30(6), 1153–1157. [https://doi.org/10.1175/1520-0469\(1973\)030%3C1153:INCIRA%3E2.0.CO;2](https://doi.org/10.1175/1520-0469(1973)030%3C1153:INCIRA%3E2.0.CO;2)
- Bond, T. C. (2004). A technology-based global inventory of black and organic carbon emissions from combustion. *Journal of Geophysical Research*, 109, D14203. <https://doi.org/10.1029/2003JD003697>
- Bond, T. C., Doherty, S. J., Fahey, D. W., Forster, P. M., Berntsen, T., DeAngelo, B. J., et al. (2013). Bounding the role of black carbon in the climate system: A scientific assessment. *Journal of Geophysical Research: Atmospheres*, 118, 5380–5552. <https://doi.org/10.1002/jgrd.50171>
- Brooks, S. D., Suter, K., & Olivarez, L. (2014). Effects of chemical aging on the ice nucleation activity of soot and polycyclic aromatic hydrocarbon aerosols. *The Journal of Physical Chemistry A*, 118(43), 10,036–10,047. <https://doi.org/10.1021/jp508809y>
- Connolly, P. J., Möhler, O., Field, P. R., Saathoff, H., Burgess, R., Choularton, T., & Gallagher, M. (2009). Studies of heterogeneous freezing by three different desert dust samples. *Atmospheric Chemistry and Physics*, 9(8), 2805–2824. <https://doi.org/10.5194/acp-9-2805-2009>
- Cozic, J., Mertes, S., Verheggen, B., Cziczo, D. J., Gallavardin, S. J., Walter, S., et al. (2008). Black carbon enrichment in atmospheric ice particle residuals observed in lower tropospheric mixed phase clouds. *Journal of Geophysical Research*, 113, D15209. <https://doi.org/10.1029/2007JD009266>
- DeMott, P. J. (1990). An exploratory study of ice nucleation by soot aerosols. *Journal of Applied Meteorology*, 29(10), 1072–1079. [https://doi.org/10.1175/1520-0450\(1990\)029%3C1072:AESOIN%3E2.0.CO;2](https://doi.org/10.1175/1520-0450(1990)029%3C1072:AESOIN%3E2.0.CO;2)
- Diehl, K., & Mitra, S. K. (1998). A laboratory study of the effects of a kerosene-burner exhaust on ice nucleation and the evaporation rate of ice crystals. *Atmospheric Environment*, 32(18), 3145–3151. [https://doi.org/10.1016/S1352-2310\(97\)00467-6](https://doi.org/10.1016/S1352-2310(97)00467-6)
- Fan, S. M., Schwarz, J. P., Liu, J., Fahey, D. W., Ginoux, P., Horowitz, L. W., et al. (2012). Inferring ice formation processes from global-scale black carbon profiles observed in the remote atmosphere and model simulations. *Journal of Geophysical Research*, 117, D23205. <https://doi.org/10.1029/2012JD018126>
- Fitzpatrick, E. M., Jones, J. A., Pourkashanian, M., Ross, A. B., Williams, A., & Bartle, K. D. (2008). Mechanistic aspects of soot formation from the combustion of pine wood. *Energy & Fuels*, 22(6), 3771–3778. <https://doi.org/10.1021/ef800456k>
- Gorbunov, B., Baklanov, A., Kakutkina, N., Windsor, H., & Toumi, R. (2001). Ice nucleation on soot particles. *Journal of Aerosol Science*, 32(2), 199–215. [https://doi.org/10.1016/S0021-8502\(00\)00077-X](https://doi.org/10.1016/S0021-8502(00)00077-X)
- Harris, N. R. P., Carpenter, L. J., Lee, J. D., Vaughan, G., Filus, M. T., Jones, R. L., et al. (2017). Coordinated airborne studies in the tropics (CAST). *Bulletin of the American Meteorological Society*, 98(1), 145–162. <https://doi.org/10.1175/BAMS-D-14-00290.1>
- Harrison, A. D., Whale, T. F., Carpenter, M. A., Holden, M. A., Neve, L., O'Sullivan, D., et al. (2016). Not all feldspar is equal: A survey of ice nucleating properties across the feldspar group of minerals. *Atmospheric Chemistry and Physics Discussions*, 1–26. <https://doi.org/10.5194/acp-2016-136>
- Hoose, C., Kristjánsson, J. E., & Burrows, S. M. (2010). How important is biological ice nucleation in clouds on a global scale? *Environmental Research Letters*, 5(2), 024009. <https://doi.org/10.1088/1748-9326/5/2/024009>
- Hoose, C., Kristjánsson, J. E., Chen, J.-P., & Hazra, A. (2010). A classical-theory-based parameterization of heterogeneous ice nucleation by mineral dust, soot, and biological particles in a global climate model. *Journal of the Atmospheric Sciences*, 67(8), 2483–2503. <https://doi.org/10.1175/2010JAS3425.1>
- Hoose, C., & Möhler, O. (2012). Heterogeneous ice nucleation on atmospheric aerosols: A review of results from laboratory experiments. *Atmospheric Chemistry and Physics*, 12(20), 9817–9854. <https://doi.org/10.5194/acp-12-9817-2012>
- Kamphus, M., Ettner-Mahl, M., Klimach, T., Drewnick, F., Keller, L., Cziczo, D. J., et al. (2010). Chemical composition of ambient aerosol, ice residues and cloud droplet residues in mixed-phase clouds: Single particle analysis during the cloud and aerosol characterization experiment (CLACE 6). *Atmospheric Chemistry and Physics*, 10(16), 8077–8095. <https://doi.org/10.5194/acp-10-8077-2010>
- Kanji, Z. A., DeMott, P. J., Möhler, O., & Abbatt, J. P. D. (2011). Results from the University of Toronto continuous flow diffusion chamber at ICIS 2007: Instrument intercomparison and ice onsets for different aerosol types. *Atmospheric Chemistry and Physics*, 11(1), 31–41. <https://doi.org/10.5194/acp-11-31-2011>
- Koehler, K. A., DeMott, P. J., Kreidenweis, S. M., Popovicheva, O. B., Petters, M. D., Carrico, C. M., et al. (2009). Cloud condensation nuclei and ice nucleation activity of hydrophobic and hydrophilic soot particles. *Physical Chemistry Chemical Physics*, 11(36), 7906–7920. <https://doi.org/10.1039/b905334b>
- Kulkarni, G., China, S., Liu, S., Nandasiri, M., Sharma, N., Wilson, J., et al. (2016). Ice nucleation activity of diesel soot particles at cirrus relevant temperature conditions: Effects of hydration, secondary organics coating, soot morphology, and coagulation. *Geophysical Research Letters*, 43, 3580–3588. <https://doi.org/10.1002/2016GL068707>
- Kupiszewski, P., Zanatta, M., Mertes, S., Vochezer, P., Lloyd, G., Schneider, J., et al. (2016). Ice residual properties in mixed-phase clouds at the high-alpine Jungfraujoch site. *Journal of Geophysical Research: Atmospheres*, 121, 12,343–12,362. <https://doi.org/10.1002/2016JD024894>
- Lee, Y. H., Lamarque, J. F., Flanner, M. G., Jiao, C., Shindell, D. T., Berntsen, T., et al. (2013). Evaluation of preindustrial to present-day black carbon and its albedo forcing from Atmospheric Chemistry and Climate Model Intercomparison Project (ACCMIP). *Atmospheric Chemistry and Physics*, 13(5), 2607–2634. <https://doi.org/10.5194/acp-13-2607-2013>
- Leon, D. C., French, J. R., Lasher-Trapp, S., Blyth, A. M., Abel, S. J., Ballard, S., et al. (2016). The Convective Precipitation Experiment (COPE): Investigating the origins of heavy precipitation in the southwestern United Kingdom. *Bulletin of the American Meteorological Society*, 97(6), 1003–1020. <https://doi.org/10.1175/BAMS-D-14-00157.1>
- Levin, E. J. T., McMeeking, G. R., DeMott, P. J., McCluskey, C. S., Carrico, C. M., Nakao, S., et al. (2016). Ice-nucleating particle emissions from biomass combustion and the potential importance of soot aerosol. *Journal of Geophysical Research: Atmospheres*, 121, 5888–5903. <https://doi.org/10.1002/2016JD024879>

- Liu, D., Quennehen, B., Darbyshire, E., Allan, J. D., Williams, P. I., Taylor, J. W., et al. (2015). The importance of Asia as a source of black carbon to the European Arctic during springtime 2013. *Atmospheric Chemistry and Physics*, 15(20), 11,537–11,555. <https://doi.org/10.5194/acp-15-11537-2015>
- Lohmann, U. (2002). A glaciation indirect aerosol effect caused by soot aerosols. *Geophysical Research Letters*, 29(4), 1052. <https://doi.org/10.1029/2001GL014357>
- Lohmann, U. (2017). Anthropogenic aerosol influences on mixed-phase clouds. *Current Climate Change Reports*, 3(1), 32–44. <https://doi.org/10.1007/s40641-017-0059-9>
- Mann, G. W., Carslaw, K. S., Spracklen, D. V., Ridley, D. A., Manktelow, P. T., Chipperfield, M. P., et al. (2010). Description and evaluation of GLOMAP-mode: A modal global aerosol microphysics model for the UKCA composition-climate model. *Geoscientific Model Development*, 3(2), 519–551. <https://doi.org/10.5194/gmd-3-519-2010>
- Matsui, H., Kondo, Y., Moteki, N., Takegawa, N., Sahu, L. K., Zhao, Y., et al. (2011). Seasonal variation of the transport of black carbon aerosol from the Asian continent to the Arctic during the ARCTAS aircraft campaign. *Journal of Geophysical Research*, 116, D05202. <https://doi.org/10.1029/2010JD015067>
- Mccluskey, C. S., Demott, P. J., Prenni, A. J., Levin, E. J. T., Mcmeeking, G. R., Sullivan, A. P., et al. (2014). Characteristics of atmospheric ice nucleating particles associated with biomass burning in the US: Prescribed burns and wildfire. *Journal of Geophysical Research: Atmospheres*, 119, 10,458–10,470. <https://doi.org/10.1002/2014JD021980>
- McMeeking, G. R., Hamburger, T., Liu, D., Flynn, M., Morgan, W. T., Northway, M., et al. (2010). Black carbon measurements in the boundary layer over western and northern Europe. *Atmospheric Chemistry and Physics*, 10(19), 9393–9414. <https://doi.org/10.5194/acp-10-9393-2010>
- McMeeking, G. R., Morgan, W. T., Flynn, M., Highwood, E. J., Turnbull, K., Haywood, J., & Coe, H. (2011). Black carbon aerosol mixing state, organic aerosols and aerosol optical properties over the United Kingdom. *Atmospheric Chemistry and Physics*, 11(17), 9037–9052. <https://doi.org/10.5194/acp-11-9037-2011>
- Metcalfe, A. R., Craven, J. S., Ensberg, J. J., Brioude, J., Angevine, W., Sorooshian, A., et al. (2012). Black carbon aerosol over the Los Angeles Basin during CalNex. *Journal of Geophysical Research*, 117, D00V13. <https://doi.org/10.1029/2011JD017255>
- Möhler, O. (2005). Effect of sulfuric acid coating on heterogeneous ice nucleation by soot aerosol particles. *Journal of Geophysical Research*, 110, D11210. <https://doi.org/10.1029/2004JD005169>
- Murray, B. J., O'Sullivan, D., Atkinson, J. D., & Webb, M. E. (2012). Ice nucleation by particles immersed in supercooled cloud droplets. *Chemical Society Reviews*, 41(19), 6519–6554. <https://doi.org/10.1039/c2cs35200a>
- O'Shea, S. J., Allen, G., Gallagher, M. W., Bower, K., Illingworth, S. M., Muller, J. B. A., et al. (2014). Methane and carbon dioxide fluxes and their regional scalability for the European Arctic wetlands during the MAMM project in summer 2012. *Atmospheric Chemistry and Physics*, 14(23), 13,159–13,174. <https://doi.org/10.5194/acp-14-13159-2014>
- Oshima, N., Kondo, Y., Moteki, N., Takegawa, N., Koike, M., Kita, K., et al. (2012). Wet removal of black carbon in Asian outflow: Aerosol Radiative Forcing in East Asia (A-FORCE) aircraft campaign. *Journal of Geophysical Research*, 117, D03204. <https://doi.org/10.1029/2011JD016552>
- Perring, A. E., Schwarz, J. P., Markovic, M. Z., Fahey, D. W., Jimenez, J. L., Campuzano-Jost, P., et al. (2017). In situ measurements of water uptake by black carbon-containing aerosol in wildfire plumes. *Journal of Geophysical Research: Atmospheres*, 122, 1086–1097. <https://doi.org/10.1002/2016JD025688>
- Petters, M. D., Parsons, M. T., Prenni, A. J., DeMott, P. J., Kreidenweis, S. M., Carrico, C. M., et al. (2009). Ice nuclei emissions from biomass burning. *Journal of Geophysical Research*, 114, D07209. <https://doi.org/10.1029/2008JD011532>
- Petzold, A., Ogren, J. A., Fiebig, M., Laj, P., Li, S. M., Baltensperger, U., et al. (2013). Recommendations for reporting black carbon measurements. *Atmospheric Chemistry and Physics*, 13(16), 8365–8379. <https://doi.org/10.5194/acp-13-8365-2013>
- Phillips, V. T. J., DeMott, P. J., & Andronache, C. (2008). An empirical parameterization of heterogeneous ice nucleation for multiple chemical species of aerosol. *Journal of the Atmospheric Sciences*, 65(9), 2757–2783. <https://doi.org/10.1175/2007JAS2546.1>
- Phillips, V. T. J., Demott, P. J., Andronache, C., Pratt, K. A., Prather, K. A., Subramanian, R., & Twohy, C. (2013). Improvements to an empirical parameterization of heterogeneous ice nucleation and its comparison with observations. *Journal of the Atmospheric Sciences*, 70(2), 378–409. <https://doi.org/10.1175/JAS-D-12-080.1>
- Popovicheva, O., Kireeva, E., Persiantseva, N., Khokhlova, T., Shonija, N., Tishkova, V., & Demirdjian, B. (2008). Effect of soot on immersion freezing of water and possible atmospheric implications. *Atmospheric Research*, 90(2–4), 326–337. <https://doi.org/10.1016/j.atmosres.2008.08.004>
- Pratt, K. A., DeMott, P. J., French, J. R., Wang, Z., Westphal, D. L., Heymsfield, A. J., et al. (2009). In situ detection of biological particles in cloud ice-crystals. *Nature Geoscience*, 2(6), 398–401. <https://doi.org/10.1038/ngeo521>
- Reddington, C. L., Carslaw, K. S., Stier, P., Schutgens, N., Coe, H., Liu, D., et al. (2017). The Global Aerosol Synthesis And Science Project (GASSP): Measurements and modelling to reduce uncertainty. *Bulletin of the American Meteorological Society*, 98(9), 1857–1877. <https://doi.org/10.1175/BAMS-D-15-00317.1>
- Ryerson, T. B., Andrews, A. E., Angevine, W. M., Bates, T. S., Brock, C. A., Cairns, B., et al. (2013). The 2010 California research at the Nexus of air quality and climate change (CalNex) field study. *Journal of Geophysical Research: Atmospheres*, 118, 5830–5866. <https://doi.org/10.1002/jgrd.50331>
- Savre, J., & Ekman, A. M. L. (2015). A theory-based parameterization for heterogeneous ice nucleation and implications for the simulation of ice processes in atmospheric models. *Journal of Geophysical Research: Atmospheres*, 120, 4937–4961. <https://doi.org/10.1002/2014JD023000>
- Schill, G. P., Jathar, S. H., Kodros, J. K., Levin, E. J. T., Galang, A. M., Friedman, B., et al. (2016). Ice nucleating particle emissions from photochemically-aged diesel and biodiesel exhaust. *Geophysical Research Letters*, 43, 5524–5531. <https://doi.org/10.1002/2016GL069529>
- Schmidt, S., Schneider, J., Klimach, T., Mertes, S., Schenk, L. P., Kupiszewski, P., et al. (2017). Online single particle analysis of ice particle residuals from mountain-top mixed-phase clouds using laboratory derived particle type assignment. *Atmospheric Chemistry and Physics*, 17(1), 575–594. <https://doi.org/10.5194/acp-17-575-2017>
- Schutgens, N., Tsyro, S., Gryspeerdt, E., Goto, D., Weigum, N., Schulz, M., & Stier, P. (2017). On the spatio-temporal representativeness of observations. *Atmospheric Chemistry and Physics*, 17(16), 9761–9780. <https://doi.org/10.5194/acp-17-9761-2017>
- Schutgens, N. A. J., Gryspeerdt, E., Weigum, N., Tsyro, S., Goto, D., Schulz, M., & Stier, P. (2016). Will a perfect model agree with perfect observations? The impact of spatial sampling. *Atmospheric Chemistry and Physics*, 16(10), 6335–6353. <https://doi.org/10.5194/acp-16-6335-2016>
- Schwarz, J. P., Gao, R. S., Spackman, J. R., Watts, L. A., Thomson, D. S., Fahey, D. W., et al. (2008). Measurement of the mixing state, mass, and optical size of individual black carbon particles in urban and biomass burning emissions. *Geophysical Research Letters*, 35, L13810. <https://doi.org/10.1029/2008GL033968>

- Schwarz, J. P., Spackman, J. R., Gao, R. S., Watts, L. A., Stier, P., Schulz, M., et al. (2010). Global-scale black carbon profiles observed in the remote atmosphere and compared to models. *Geophysical Research Letters*, 37, L18812. <https://doi.org/10.1029/2010GL044372>
- Spackman, J. R., Gao, R. S., Neff, W. D., Schwarz, J. P., Watts, L. A., Fahey, D. W., et al. (2010). Aircraft observations of enhancement and depletion of black carbon mass in the springtime Arctic. *Atmospheric Chemistry and Physics*, 10(19), 9667–9680. <https://doi.org/10.5194/acp-10-9667-2010>
- Spracklen, D. V., Carslaw, K. S., Pöschl, U., Rap, A., & Forster, P. M. (2011). Global cloud condensation nuclei influenced by carbonaceous combustion aerosol. *Atmospheric Chemistry and Physics*, 11(17), 9067–9087. <https://doi.org/10.5194/acp-11-9067-2011>
- Subramanian, R., Kok, G. L., Baumgardner, D., Clarke, A., Shinozuka, Y., Campos, T. L., et al. (2010). Black carbon over Mexico: The effect of atmospheric transport on mixing state, mass absorption cross-section, and BC/CO ratios. *Atmospheric Chemistry and Physics*, 10(1), 219–237. <https://doi.org/10.5194/acp-10-219-2010>
- Taylor, J. W., Allan, J. D., Allen, G., Coe, H., Williams, P. I., Flynn, M. J., et al. (2014). Size-dependent wet removal of black carbon in Canadian biomass burning plumes. *Atmospheric Chemistry and Physics*, 14(24), 13,755–13,771. <https://doi.org/10.5194/acp-14-13755-2014>
- Twohy, C. H., DeMott, P. J., Pratt, K. A., Subramanian, R., Kok, G. L., Murphy, S. M., et al. (2010). Relationships of biomass-burning aerosols to ice in orographic wave clouds. *Journal of the Atmospheric Sciences*, 67(8), 2437–2450. <https://doi.org/10.1175/2010JAS3310.1>
- Ullrich, R., Hoose, C., Möhler, O., Niemand, M., Wagner, R., Höhler, K., et al. (2017). A new ice nucleation active site parameterization for desert dust and soot. *Journal of the Atmospheric Sciences*, 74(3), 699–717. <https://doi.org/10.1175/JAS-D-16-0074.1>
- Van der Werf, G. R., Randerson, J. T., Collatz, G. J., & Giglio, L. (2003). Carbon emissions from fires in tropical and subtropical ecosystems. *Global Change Biology*, 9(4), 547–562. <https://doi.org/10.1046/j.1365-2486.2003.00604.x>
- Vergara-Temprado, J., Murray, B. J., Wilson, T. W., O'Sullivan, D., Browse, J., Pringle, K. J., et al. (2017). Contribution of feldspar and marine organic aerosols to global ice nucleating particle concentrations. *Atmospheric Chemistry and Physics*, 17(5), 3637–3658. <https://doi.org/10.5194/acp-17-3637-2017>
- Wang, Y., Liu, X., Hoose, C., & Wang, B. (2014). Different contact angle distributions for heterogeneous ice nucleation in the Community Atmospheric Model version 5. *Atmospheric Chemistry and Physics*, 14(19), 10,411–10,430. <https://doi.org/10.5194/acp-14-10411-2014>
- Whale, T. F., Murray, B. J., O'Sullivan, D., Umo, N. S., Baustian, K. J., Atkinson, J. D., & Morris, G. J. (2014). A technique for quantifying heterogeneous ice nucleation in microlitre supercooled water droplets. *Atmospheric Measurement Techniques Discussions*, 7(9), 9509–9536. <https://doi.org/10.5194/amtd-7-9509-2014>
- Wilson, J. M., Baeza-Romero, M. T., Jones, J. M., Pourkashanian, M., Williams, A., Lea-Langton, A. R., et al. (2013). Soot formation from the combustion of biomass pyrolysis products and a hydrocarbon fuel, n-decane: An Aerosol Time Of Flight Mass Spectrometer (ATOFMS) study. *Energy & Fuels*, 27(3), 1668–1678. <https://doi.org/10.1021/ef3019386>
- Wilson, T. W., Ladino, L. A., Alpert, P. A., Breckels, M. N., Brooks, I. M., Browse, J., et al. (2015). A marine biogenic source of atmospheric ice-nucleating particles. *Nature*, 525(7568), 234–238. <https://doi.org/10.1038/nature14986>
- Wright, T. P., Petters, M. D., Hader, J. D., Morton, T., & Holder, A. L. (2013). Minimal cooling rate dependence of ice nuclei activity in the immersion mode. *Journal of Geophysical Research: Atmospheres*, 118, 10,535–10,543. <https://doi.org/10.1002/jgrd.50810>
- Yin, J., Wang, D., & Zhai, G. (2012). An evaluation of ice nuclei characteristics from the long-term measurement data over North China. *Asia-Pacific Journal of Atmospheric Sciences*, 48(2), 197–204. <https://doi.org/10.1007/s13143-012-0020-8>
- Yun, Y., & Penner, J. E. (2012). Global model comparison of heterogeneous ice nucleation parameterizations in mixed phase clouds. *Journal of Geophysical Research*, 117, D07203. <https://doi.org/10.1029/2011JD016506>