

Central Lancashire Online Knowledge (CLOK)

Title	Apposition to endometrial epithelial cells activates mouse blastocysts for implantation.
Type	Article
URL	https://clock.uclan.ac.uk/31553/
DOI	https://doi.org/10.1093/molehr/gax043
Date	2017
Citation	Ruane, Peter T., Berneau, Stephane, Koeck, Rebekka, Watts, Jessica, Kimber, Susan J., Brison, Daniel R., Westwood, Melissa and Aplin, John D. (2017) Apposition to endometrial epithelial cells activates mouse blastocysts for implantation. <i>Molecular Human Reproduction</i> , 23 (9). pp. 617-627. ISSN 1360-9947
Creators	Ruane, Peter T., Berneau, Stephane, Koeck, Rebekka, Watts, Jessica, Kimber, Susan J., Brison, Daniel R., Westwood, Melissa and Aplin, John D.

It is advisable to refer to the publisher's version if you intend to cite from the work.
<https://doi.org/10.1093/molehr/gax043>

For information about Research at UCLan please go to <http://www.uclan.ac.uk/research/>

All outputs in CLOK are protected by Intellectual Property Rights law, including Copyright law. Copyright, IPR and Moral Rights for the works on this site are retained by the individual authors and/or other copyright owners. Terms and conditions for use of this material are defined in the <http://clock.uclan.ac.uk/policies/>



Apposition to endometrial epithelial cells activates mouse blastocysts for implantation

DOI:
[10.1093/molehr/gax043](https://doi.org/10.1093/molehr/gax043)

Document Version
Accepted author manuscript

[Link to publication record in Manchester Research Explorer](#)

Citation for published version (APA):

Ruane, P., Berneau, S., Koeck, R., Kimber, S., Brison, D., Westwood, M., & Aplin, J. (2017). Apposition to endometrial epithelial cells activates mouse blastocysts for implantation. *Molecular Human Reproduction*, 23(9), 617-627. <https://doi.org/10.1093/molehr/gax043>

Published in:
Molecular Human Reproduction

Citing this paper

Please note that where the full-text provided on Manchester Research Explorer is the Author Accepted Manuscript or Proof version this may differ from the final Published version. If citing, it is advised that you check and use the publisher's definitive version.

General rights

Copyright and moral rights for the publications made accessible in the Research Explorer are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

Takedown policy

If you believe that this document breaches copyright please refer to the University of Manchester's Takedown Procedures [<http://man.ac.uk/04Y6Bo>] or contact uml.scholarlycommunications@manchester.ac.uk providing relevant details, so we can investigate your claim.



Draft Manuscript For Review. Reviewers should submit their review at
<http://mc.manuscriptcentral.com/molehr>

Apposition to endometrial epithelial cells activates mouse blastocysts for implantation

Journal:	<i>Molecular Human Reproduction</i>
Manuscript ID	MHR-17-0084.R1
Manuscript Type:	Original Research
Date Submitted by the Author:	19-Jun-2017
Complete List of Authors:	Ruane, Peter; University of Manchester, Maternal and Fetal Health Berneau, Stephane; University of Manchester, Maternal and Fetal Health Koeck, Rebekka; University of Manchester, Maternal and Fetal Health Watts, Jessica; University of Manchester, Maternal and Fetal Health Kimber, Susan J; University of Manchester, Division of Cell Matrix Biology and Regenerative Medicine Brison, Daniel; University of Manchester, Maternal and Fetal Health; Central Manchester University Hospitals NHS Foundation Trust, Department of Reproductive Medicine Westwood, Melissa; University of Manchester, Maternal and Fetal Health Aplin, John; University of Manchester, Maternal and Fetal Health
Key Words:	implantation, endometrium, embryo development, trophoblast, transcription factors
Note: The following files were submitted by the author for peer review, but cannot be converted to PDF. You must view these files (e.g. movies) online.	
14fps 36 frames 100um scale bar 8bit.avi	

Table 1

Gene	Primer sequences (5'-3')
<i>Cdx2</i>	CAAGGACGTGAGCATGTATCC GTAACCACCGTAGTCCGGGTA
<i>Gata3</i>	CTCGGCCATTCTGTACATGGAA GGATACCTCTGCACCGTAGC
<i>Eomes</i>	GCGCATGTTTCCTTTCTTGAG GGTCGGCCAGAACCACTTC
<i>Hand1</i>	CTACCAGTTACATCGCCTACTTG ACCACCATCCGTCTTTTTGAG
<i>Gata2</i>	CACCCCGCCGTATTGAATG CCTGCGAGTCGAGATGGTTG
<i>Hes1</i>	CCAGCCAGTGTCAACACGA AATGCCGGGAGCTATCTTTCT
<i>Gapdh</i>	AGGTCGGTGTGAACGGATTTG GGGGTCGTTGATGGCAACA
<i>Ywhaz</i>	TTGATCCCCAATGCTTCGCAA CAGCAACCTCGGCCAAGTAA


Figure 1

Figure 1 A Schematic to illustrate the criteria for the 4-point grading of the stability of mouse blastocyst attachment to Ishikawa cells; [1] = not attached, [2] = weakly attached, [3] = intermediately attached, [4] = stably attached. B Hatched E4.5 mouse blastocysts were co-cultured with Ishikawa cells for 48h and the stability of attachment was scored (1-4) every 4h from 0-12h, 24-36h, and at 48h. The mean attachment stability scores, from 4 independent experiments using 102 blastocysts, are presented. Error bars +/- standard error of the mean (SEM). C The mean attachment stability score was calculated from the data in B. Error bars +/- SEM. D Phase contrast images of the mouse blastocyst-Ishikawa cell co-culture at 4h, 8h, 12h, 28h, 32h and 36h. Blastocysts are indicated by arrows and the attachment score for each blastocyst is displayed. Scale bars 200µm. E The cumulative frequency of the time at which the final position of attachment was reached was plotted alongside the percentage of blastocysts scored as intermediately or stably attached (mean +/- SEM from 3 independent experiments using 84 blastocysts). F Ishikawa cells were pre-treated with 10nM 17β-estradiol 48h prior to co-culture, then with 10nM 17β-estradiol and 1µM progesterone 24h prior to co-culture and during co-culture with hatched E4.5 mouse blastocysts. The

attachment stability of mouse blastocysts was scored every 4h from 0-12h, 24-36h, and at 48h. Blastocysts scored as intermediately and stably attached were added together and the mean from 3 independent experiments using 72 blastocysts was plotted +/- SEM.

201x296mm (300 x 300 DPI)

For Review Only


Figure 2

Figure 2 Attached embryos not breaching (A) and breaching (B) Ishikawa cell layers were optically sectioned after PFA fixation and incubation with phalloidin to label actin fibres (red) and DAPI to label nuclei (blue).

Invasive cells of breaching embryos exhibit binuclearity (closed arrow), prominent nucleosomes (arrowhead), and actin stress fibres (open arrow). Upper/lower section refers to depth of the optical section in the Z-axis. Scale bars 20 μm . C The size of invasive embryonic nuclei in the lower optical section was measured and compared to the size of outer, peripheral embryonic nuclei, presumed to be TE and not inner cell mass, in the corresponding upper section. 309 nuclei were measured in 12 embryos and mean nuclear size was plotted $\pm$ SEM. *** t-test $p < 0.001$. Breaching embryos were fixed and labelled with anti-CDX2 antibody (D), rabbit isotype control antibody (D inset), anti-GATA3 antibody (E), or mouse isotype control antibody (E inset), as well as phalloidin and DAPI. Upper/lower section refers to depth of the optical section in the Z-axis. Scale bars 20 μm . F The intensity of CDX2 and GATA3 fluorescence labelling was measured in invasive embryonic nuclei in the lower optical section and compared to that of outer embryonic nuclei, presumed to be TE and not inner cell mass, in the corresponding upper section. 158 nuclei were measured in 5 embryos labelled with anti-CDX2 and 226 nuclei were measured in 5 embryos labelled with anti-GATA3. The mean ratio of invasive nuclei to upper TE nuclei fluorescence intensity was plotted $\pm$ SEM. ** t-test

$p < 0.01$, NS not significant.

208x239mm (300 x 300 DPI)

For Review Only


Figure 3

Figure 3 A Schematic representing embryo-Ishikawa cell co-culture experimental design. B Hatched E4.5 mouse blastocysts were either co-cultured with Ishikawa cells to E6.5 or cultured alone in co-culture media to E5.5 before co-culture with Ishikawa cells to E6.5 or E7.5. The stability of attachment was scored every 4h from E5.5-6 and at E6.5. Mean percentage blastocysts scored as intermediately or stably attached from 6 independent experiments using 149 blastocysts was plotted $\pm$ SEM. C Attached embryos co-cultured from E4.5-6.5, E5.5-6.5 and E5.5-7.5, were fixed, labelled with phalloidin and DAPI and optically sectioned to determine whether embryonic cells had breached the Ishikawa cells. The mean proportion of attached embryos to have breached the Ishikawa cells was plotted $\pm$ SEM. ** ANOVA $p < 0.01$, *** ANOVA $p < 0.001$, demonstrating significant difference from co-culture E4.5-6.5.

209x234mm (300 x 300 DPI)

Figure 4




Figure 4 A Schematic representing embryo-Ishikawa cell co-culture experimental design. B Hatched E4.5 mouse blastocysts were co-cultured directly with Ishikawa cells from E4.5-6.5, co-cultured directly with Ishikawa cells to E5.5 before transfer to fresh Ishikawa cells and co-cultured to E6.5, or co-cultured in a transwell above Ishikawa cells to E5.5 before transfer to fresh Ishikawa cells and co-cultured to E6.5. The stability of attachment was scored every 4h from E5.5-6 and at E6.5. Mean percentage blastocysts scored as intermediately or stably attached from 5 independent experiments using 157 blastocysts was plotted +/- SEM. C Attached embryos were fixed, labelled with phalloidin and DAPI and optically sectioned to determine whether embryonic cells have breached the Ishikawa cells. The mean proportion of attached embryos to have breached the Ishikawa cells +/- SEM. ** ANOVA $p < 0.01$, demonstrating significant difference from direct co-culture E4.5-6.5.

254x250mm (300 x 300 DPI)

Figure 5




Figure 5 Hatched E4.5 mouse blastocysts were cultured alone, co-cultured directly with Ishikawa cells, co-cultured in a transwell above Ishikawa cells or co-cultured directly with human fibroblasts as a control cell type, to E5.5 before retrieval and lysis for RNA isolation and gene expression analysis by qPCR. Data is presented as expression relative to reference genes (2-ct, relative to geometric mean of Gapdh and Ywhaz) of A *Cdx2*, B *Gata3*, C *Eomes*, D *Hand1*, E *Gata2*, and F *Hes1* in embryos cultured in indicated conditions (mean $\pm$ SEM of 5 independent experiments using 200 embryos). * ANOVA $p < 0.05$, demonstrating significant difference from no co-culture E4.5-5.5.

207x198mm (300 x 300 DPI)

**Apposition to endometrial epithelial cells activates mouse blastocysts for
implantation**

Running title: Blastocyst apposition to endometrial cells

Peter T Ruane^{1,2,*}, Stéphane C Berneau^{1,2}, Rebekka Koeck^{1,2}, Jessica Watts^{1,2}, Susan J
Kimber³, Daniel R Brison^{1,2,4}, Melissa Westwood^{1,2} and John D Aplin^{1,2}

¹Maternal and Fetal Health Research Centre, Division of Developmental Biology and
Medicine, School of Medical Sciences, Faculty of Biology, Medicine and Health, University of
Manchester, Manchester Academic Health Sciences Centre, St. Mary's Hospital,
Manchester, M13 9WL.

²Maternal and Fetal Health Research Centre, St. Mary's Hospital, Central Manchester
University Hospitals NHS Foundation Trust, Manchester Academic Health Sciences Centre,
Manchester, M13 9WL.

³Division of Cell Matrix Biology and Regenerative Medicine, School of Biological Sciences,
Faculty of Biology Medicine and Health, University of Manchester, Michael Smith Building,
Manchester, M13 9PT.

⁴Department of Reproductive Medicine, Old St Mary's Hospital, Central Manchester
University Hospitals NHS Foundation Trust, Manchester Academic Health Science Centre,
Oxford Road, Manchester M13 9WL.

* Corresponding author peter.ruane@manchester.ac.uk

20 Abstract

21 **Study question:** How do interactions between blastocyst-stage embryos and endometrial
22 epithelial cells regulate the early stages of implantation in an in-vitro model?

23 **Summary Answer:** Mouse blastocyst apposition with human endometrial epithelial cells
24 initiates trophectoderm differentiation to trophoblast, which goes on to breach the
25 endometrial epithelium.

26 **What is known already:** In-vitro models using mouse blastocysts and human endometrial
27 cell lines have proven invaluable in the molecular characterisation of embryo attachment to
28 endometrial epithelium at the onset of implantation. Genes involved in embryonic breaching
29 of the endometrial epithelium have not been investigated in such in-vitro models.

30 **Study design, size, duration:** This study used an established in-vitro model of implantation
31 to examine cellular and molecular interactions during blastocyst attachment to endometrial
32 epithelial cells.

33 **Participants/materials, setting, methods:** Mouse blastocysts developed from embryonic
34 day (E) 1.5 in vitro were hatched and co-cultured with confluent human endometrial
35 adenocarcinoma-derived Ishikawa cells in serum-free medium. A scale of attachment
36 stability based on blastocyst oscillation upon agitation was devised. Blastocysts were
37 monitored for 48h to establish the kinetics of implantation, and optical sectioning using
38 fluorescence microscopy revealed attachment and invasion interfaces. Quantitative PCR
39 was used to determine blastocyst gene expression. Data from a total of 680 mouse
40 blastocysts are reported, with 3-6 experimental replicates. T-test and ANOVA analyses
41 established statistical significance at $p < 0.05$, $p < 0.01$ and $p < 0.001$.

42 **Main results and the role of chance:** Hatched E4.5 mouse blastocysts exhibited weak
43 attachment to confluent Ishikawa cells over the first 24h of co-culture, with intermediate and
44 stable attachment occurring from 28h (E5.5+4h) in a hormone-independent manner.

Attached embryos fixed after 48h (E6.5) frequently exhibited outgrowths, characterised morphologically and with antibody markers as trophoblast giant cells (TGCs), which had breached the Ishikawa cell layer. Beginning co-culture at E5.5 also resulted in intermediate and stable attachment from E5.5+4h, however these embryos did not go on to breach the Ishikawa cell layer, even when co-culture was extended to E7.5 ($p<0.01$). Blastocysts cultured from E4.5 in permeable transwell inserts above Ishikawa cells before transfer to direct co-culture at E5.5 went on to attach but failed to breach the Ishikawa cell layer by E6.5 ($p<0.01$). Gene expression analysis at E5.5 demonstrated that direct co-culture with Ishikawa cells from E4.5 resulted in downregulation of trophoblast transcription factors *Cdx2* ($p<0.05$) and *Gata3* ($p<0.05$) and upregulation of the trophoblast giant cell transcription factor *Hand1* ($p<0.05$). Co-culture with non-endometrial human fibroblasts did not alter the expression of these genes.

Large scale data: N/A

Limitations, reasons for caution: The in-vitro model used here combines human carcinoma-derived endometrial cells with mouse embryos, in which the cellular interactions observed may not fully recapitulate those in-vivo. The data gleaned from such models can be regarded as hypothesis-generating, and research is now needed to develop more sophisticated models of human implantation combining multiple primary endometrial cell types with surrogate and real human embryos.

Wider implications of the findings: This study implicates blastocyst apposition to endometrial epithelial cells as a critical step in trophoblast differentiation required for implantation. Understanding this maternal regulation of the embryonic developmental programme may lead to novel treatments for infertility.

Study funding and competing interest(s): This work was supported by funds from the charities Wellbeing of Women (RG1442) and Diabetes UK (15/0005207), and studentship support for SCB from the Anatomical Society. No conflict of interest is declared.

71 **Keywords: implantation, endometrium, embryo development, trophoblast,**
72 **transcription factors**

73 **Introduction**

74 Implantation begins with attachment of the trophectoderm (TE) of a blastocyst-stage embryo
75 to the hormonally-responsive receptive endometrial luminal epithelium (LE), followed by
76 breaching of this barrier and invasion of TE-derived trophoblast into the underlying decidua
77 (Aplin & Kimber 2004, Aplin & Ruane 2017). Implantation failure remains a bottleneck in
78 human assisted reproduction treatment, with only ~25% of treatment cycles resulting in a
79 live birth (Ferraretti *et al.* 2012), despite efforts to select developmentally-competent
80 embryos and receptive endometrium (Brison *et al.* 2004, Aplin 2006, Dominguez *et al.* 2008,
81 Glujovsky *et al.* 2012, Ruiz-Alonso *et al.* 2013, Salamonsen *et al.* 2013, Armstrong *et al.*
82 2015, Harbottle *et al.* 2015, Simon *et al.* 2015). A comprehensive understanding of this
83 founding stage of pregnancy is necessary to improve treatments for infertility.

84 Initial maternal-embryo interactions are mediated through the trophic and paracrine action of
85 uterine fluid, secreted by uterine glands, which promotes embryo development to blastocyst
86 and subsequent implantation (Filant & Spencer 2014). There is now considerable evidence
87 that bidirectional signalling between human blastocysts and endometrium elicits responses
88 which modulate implantation receptivity (Evans *et al.* 2016). For example, blastocyst-
89 dependent local reduction in the anti-adhesive glycocalyx of the LE (Meseguer *et al.* 2001),
90 and LE microRNA regulation of TE gene expression (Vilella *et al.* 2015), are suggested to
91 promote embryo attachment. Molecular dialogue has been observed upon blastocyst
92 apposition to the endometrium in mice, with both paracrine and juxtacrine activation of
93 ErbB4 by soluble and membrane-bound HB-EGF, respectively, promoting expression of
94 integrins at the surface of LE and TE, leading to blastocyst attachment and invasion (Wang
95 *et al.* 2000, Wang *et al.* 2002). In humans, the homophilic receptor, trophinin, may feed into
96 the HB-EGF axis at apposition to promote TE proliferation in readiness for invasion while

97 triggering LE apoptosis to allow epithelial breaching (Sugihara *et al.* 2007, Tamura *et al.*
98 2011).

99 After blastocyst attachment to the LE in humans and rodents, embryonic invasion results in
100 haemochorial placentation (Rossant & Cross 2001). The initial invasive cell type in the
101 mouse embryo is the trophoblast giant cell (TGC) (Dickson 1963), whereas in human and
102 macaque this is thought to be primary syncytium (Hertig *et al.* 1956, Enders 2007).
103 Significant progress has been made in understanding the regulatory networks governing the
104 formation of these cell types (Knott & Paul 2014). Recently trophoblast lineages have been
105 shown to arise from embryos attaching to culture surfaces in the absence of maternal cells
106 (Bedzhov & Zernicka-Goetz 2014, Deglincerti *et al.* 2016, Shahbazi *et al.* 2016), however the
107 contribution of endometrial-embryo interactions to the development of the pioneering
108 invasive embryonic cells remains unknown.

109 Characterisation of this early stage of implantation is particularly tractable in vitro. The
110 human endometrial adenocarcinoma Ishikawa cell line (Nishida *et al.* 1985), offers a model
111 epithelial system for investigating interactions with rodent and human embryos (Singh *et al.*
112 2010, Kaneko *et al.* 2011, Kang *et al.* 2014). Ishikawa cells exhibit moderate epithelial
113 polarisation and surface glycoprotein composition comparable with LE in vivo (Heneweer *et al.*
114 2005, Singh & Aplin 2014, Buck *et al.* 2015). They also mount transcriptional responses
115 to estrogen and progesterone, but do not require steroid hormones for receptivity to mouse
116 embryos (Lessey *et al.* 1996, Castelbaum *et al.* 1997, Singh *et al.* 2010, Tamm-Rosenstein
117 *et al.* 2013). Ishikawa cell-rodent embryo co-cultures are recognised as a useful model to
118 investigate molecular pathways of attachment (Kaneko *et al.* 2011, Kaneko *et al.* 2012, Kang
119 *et al.* 2014, Green *et al.* 2015, Kang *et al.* 2015), however a thorough dissection of
120 attachment and invasion in this model system has not been performed.

121 Here, we have characterised the kinetics of attachment and invasion of mouse embryos on
122 Ishikawa cells. We show that prior to stable attachment, apposition is required for

subsequent TGC invasion. Moreover, apposition leads to changes in embryonic gene expression consistent with TE differentiation to invasive TGCs. Our data suggests that the differentiation of trophoblast required for implantation of mouse blastocysts is maternally regulated, and implicates a conserved system in human LE.

Materials and Methods

Cell culture

Ishikawa cells (ECACC 99040201) and primary human foreskin fibroblasts were maintained at 37°C, 95% air and 5% CO₂ in growth medium (1:1 Dulbecco's modified Eagle's medium:Ham's-F12 (Sigma) containing 10% fetal bovine serum (Sigma) supplemented with 2mM L-glutamine, 100µg/ml streptomycin and 100IU/ml penicillin (Sigma)). Cells were cultured to confluency in 24-well plates (Greiner) on 13mm glass coverslips coated with 2% growth factor-reduced Matrigel (Sigma).

Mouse embryos

All experiments were conducted and licensed under the Animal Act, 1986, and had local ethical approval for care and use of laboratory animals and standards of humane animal care. CD1 strain mice (Charles River) were maintained by the Biological Services Unit at the University of Manchester and kept under standard environmental conditions of 12h light and 12h dark at 20–22°C and 40–60% humidity with food and water provided *ad libitum*. 8-10-week old female mice were superovulated by intraperitoneal injection with 5IU pregnant mare serum gonadotrophin (Intervet), followed by 5IU human chorionic gonadotrophin (Intervet) 46h later, then housed overnight with ≤9-month-old stud males for mating. Midday the following day was designated embryonic day (E) 0.5. Embryos were collected at E1.5 by flushing dissected oviducts with M2 medium (Millipore) containing 0.4% w/v BSA (Sigma). All embryo manipulation was performed using a Flexipet with 140µm and 300µm pipettes (Cook). E1.5 embryos were cultured for 72h in KSOM medium (Millipore) containing 0.4%

148 BSA at 37°C, 95% air and 5% CO₂. E4.5 blastocysts were hatched from the zona pellucida
149 (ZP) using acid Tyrode's (pH 2.5) (Sigma).

150 **In-vitro implantation model**

151 Ishikawa cells were grown to full confluence in 24-well plates, washed and replenished with
152 serum-free co-culture medium (1:1 DMEM:F12 supplemented with 2mM L-glutamine, 100
153 µg/ml streptomycin and 100IU/ml penicillin) 24h before transfer of three hatched mouse
154 blastocysts per well and co-culture at 37°C, 95% air and 5% CO₂. The stability of mouse
155 blastocyst attachment to Ishikawa cells was assessed using a four-point scale of blastocyst
156 behaviour upon agitation of the sample; translocation (not attached), major oscillation about
157 an attachment point (weakly attached), minor oscillation (intermediately attached), and no
158 oscillation (stably attached) (Figure 1A, Movie 1) (Kang *et al.* 2014, Kang *et al.* 2015). The
159 stability of blastocyst attachment was assessed at 4h intervals from 0-12h, 24-36h and 48h
160 using an inverted phase contrast microscope (Evos). At termination, co-cultures were
161 washed in phosphate-buffered saline (PBS) and fixed with 4% paraformaldehyde (PFA) in
162 PBS for 20 minutes. For hormone treatments, Ishikawa cells were maintained in growth
163 medium containing 10nM 17β-estradiol (Sigma) for 24h, then co-culture medium containing
164 10nM 17β-estradiol and 1µM progesterone (Sigma) for a further 24h, before addition of
165 embryos. To separate embryos from Ishikawa cells during co-culture, 3µm transwell filters
166 (Costar) were used. Blastocysts were collected from co-cultures and transwells using a
167 300µm Flexipet.

168 **Fluorescence staining and microscopy**

169 Fixed attached embryo samples were washed with PBS and quenched with 50mM
170 ammonium chloride solution before permeabilisation with 0.5% Triton-X100 PBS. Samples
171 were incubated with primary antibody (rabbit anti-CDX2 or mouse anti-GATA3, both Cell
172 Signalling Technologies) in PBS for 2h or overnight, followed by alexa568-phalloidin (Life
173 Technologies), 4',6-diamidino-2-phenylindole (DAPI) (Sigma), and alexa488 secondary

antibodies (Life Technologies) for 1h before mounting in a chamber of 3% 1,4-diazabicyclo[2.2.2]octane (Sigma) in PBS. Fluorescence microscopy was carried out using a Zeiss Axiophot microscope equipped with an Apotome module for optical sectioning. Images were analysed and processed using Zeiss Zen software and ImageJ.

Blastocyst RNA extraction and quantitative polymerase chain reaction

RNA from 10 blastocysts per treatment was isolated using the RNeasy Micro Kit (Qiagen), according to the manufacturer's instructions. Samples of 25ng RNA were added to reverse transcription (RT) reactions with random 9mer primers (Agilent) using the Sensiscript RT kit (Qiagen), according to the manufacturer's instructions. Quantitative polymerase chain reactions (qPCR) were carried out using the RT reactions along with 0.25µM primers (Table 1) and QuantiTect SYBR green PCR kit (Qiagen). qPCRs were run on a Stratagene Mx3000p machine with thermocycle parameters according to QuantiTect instructions (35 cycles using 58°C annealing temperature for all primers), and analysed with Stratagene MxPro to yield cycle threshold (Ct) values. RT reactions without sample RNA and without reverse transcriptase enzyme were used as controls in qPCR reactions with all primer pairs, and all yielded no Ct value. Dissociation curves were run with each sample to rule out the presence of non-specific PCR products.

Statistical Analysis

Independent t-test and ANOVA with Dunnett's post-hoc test were performed using SPSS (IBM), with significance at $p < 0.05$.

Results

E4.5 mouse blastocysts weakly and reversibly attach to Ishikawa cells over 24h before stably attaching in a hormone-independent manner

E4.5 mouse blastocysts hatched from the ZP barrier were introduced into co-culture with confluent Ishikawa cells in serum-free medium and monitored at 4h intervals from 0-12h, 24-

36h, and at 48h, to characterise the kinetics of attachment (Figure 1B). Only weak attachment was observed during the first 24h of co-culture (E4.5-5.5), with intermediate and stable attachment increasing from 28-48h (E5.5+4h-E6.5), as demonstrated by plotting the average stability of attachment at each time point (Figure 1C). Weak attachment was reversible since blastocysts cycled between weakly attached and not attached, and positional analysis demonstrated that weak attachment could occur at multiple successive sites (Figure 1D). Later attached blastocysts did not change position over time (Figure 1D), and cumulative frequency of the final position of blastocyst attachment correlated with intermediate and stable attachment scores (Figure 1E). These scores of attachment stability therefore represent irreversible attachment, with the lag between final position of attachment and intermediate-stable attachment likely reflecting a weak attachment phase prior to more stable attachment.

This process of prolonged, reversible weak attachment followed by initiation of stable attachment after 28h co-culture appears to mirror the apposition and attachment stages, respectively, of implantation in vivo (Enders & Schlafke 1969). Thus, the summation of intermediate and stable attachments was used for all subsequent analyses. Additionally, the kinetics of attachment were not altered by treating Ishikawa cells with estradiol and progesterone (Figure 1F), therefore further experiments were performed in the absence of steroid hormones.

Mouse blastocysts produce trophoblast giant cells to breach Ishikawa cell layers

Blastocysts attached to Ishikawa cells were fixed after 48h co-culture, fluorescently labelled and optically sectioned to visualise the interface. Although embryos that had attached to the apical surface of the Ishikawa cells but not breached the cell layer were observed (Figure 2A), outgrowths from attached embryos that had breached the Ishikawa cell layer were frequently seen (Figure 2B).

These outgrowths contained spread cells with large nuclei, clear nucleoli (arrowheads) and prominent actin stress fibres (thin arrows), and were occasionally bi- and tri-nucleate (thick arrows), reminiscent of trophoblast giant cells (TGCs) (Figure 2B). When compared to the TE nuclei above the plane of the Ishikawa cells (nuclei on the periphery of the embryo), the nuclei of embryonic cells invading into the Ishikawa cell layer were significantly larger (Figure 2C). Moreover, antibodies to the TE transcription factors CDX2 and GATA3, which are downregulated upon trophoblast differentiation to TGCs (Knott & Paul 2014), appeared to preferentially label the upper, non-invasive TE nuclei (Figure 2D, E), although only CDX2 labelling exhibited significantly lower intensity in the invasive nuclei (Figure 2F). Altogether, these observations demonstrate that mouse embryos can breach and invade Ishikawa cell layers and implicate TGCs in this process.

Co-culture from E4.5-5.5 is required for embryos to progress beyond attachment to breach the Ishikawa cell layer

We asked whether early weak attachments influence later embryo-Ishikawa interactions by comparing the attachment kinetics of blastocysts cultured with or without Ishikawa cells during E4.5-5.5 (Figure 3A). Embryos without prior co-culture were still able to stably attach to Ishikawa cells during E5.5-E6.5 (Figure 3B), though such embryos exhibited very few breaching events (Figure 3C). Notably, continued co-culture of these embryos to E7.5 still did not yield the levels of breaching observed for embryos co-cultured from E4.5-6.5 (Figure 3C). This suggests that E4.5-5.5 blastocysts require a dialogue with Ishikawa cells in order to breach the Ishikawa cell layer at a later embryonic stage.

Physical apposition with Ishikawa cells from E4.5-5.5 promotes subsequent embryonic breaching

To characterise the dialogue between embryos and Ishikawa cells, we employed a separated co-culture system. Blastocysts were incubated in permeable transwell inserts above Ishikawa cells to allow paracrine crosstalk but prevent direct contact during E4.5-5.5.

The capacity of these embryos to attach and breach after transfer to fresh Ishikawa cells at E5.5 was compared to embryos which were co-cultured directly with Ishikawa cells from E4.5 before collection (all embryos, non-attached and attached, were collected) and transfer to fresh Ishikawa cells at E5.5 (Figure 4A). The E5.5-6.5 attachment kinetics of the two groups were indistinguishable and did not differ from a control group of embryos co-cultured directly with Ishikawa cells from E4.5-6.5 without transfer to fresh Ishikawa cells at E5.5 (Figure 4B). Conversely, compared to embryos in direct co-culture during E4.5-5.5, significantly fewer embryos co-cultured in transwells from E4.5-5.5 went on to breach Ishikawa cells at E6.5 (Figure 4C). These data rule out priming of Ishikawa cells by embryos from E4.5-5.5 as a mechanism contributing embryonic invasion, and suggest that physical apposition with Ishikawa cells from E4.5-5.5 activates blastocysts for invasion.

Apposition from E4.5-5.5 induces a TGC differentiation program in the embryo

The cells forming the embryonic outgrowths at Ishikawa cell breach sites bore hallmarks of TGCs (Figure 2), leading us to hypothesise that apposition during E4.5-5.5 initiates a TGC differentiation program in the blastocyst TE. Therefore, we assessed the expression of transcription factors associated with TGC differentiation (Simmons & Cross 2005, Liu *et al.* 2009, Knott & Paul 2014), in E5.5 embryos that had not been co-cultured, had been directly or indirectly co-cultured with Ishikawa cells for 24h, or had been directly co-cultured with human foreskin fibroblasts as an alternative cell type. We found significant changes in the expression of the TE transcription factors *Cdx2* and *Gata3* (0.36- and 0.55-fold downregulation, respectively) and the TGC transcription factor *Hand1* (upregulated 3.17-fold), but only in blastocysts that had been directly co-cultured with Ishikawa cells from E4.5-5.5 (Figure 5A, B, D, respectively). No change in the expression of other TE (*Eomes*) or TGC transcription factors (*Gata2* and *Hes1*) was detected (Figure 5C, E, F, respectively). Other TGC markers, such as *Bhlhe40*, *Mdfi* and *Prf3d1*, were not detected at this stage of embryonic development (data not shown).

Discussion

Dialogue between the embryo and endometrium is necessary to orchestrate implantation in both mouse and human (Wang & Dey 2006). Here, observations in an established in-vitro model of implantation lead to the proposal that blastocyst apposition with LE initiates changes in embryonic gene expression which result in the differentiation of TE to invasive trophoblast. Such maternal regulation of embryonic differentiation constitutes a novel and potentially critical stage in embryo development; a detailed mechanistic understanding of this process could offer targets for the treatment of infertility.

Combining a categorical scale of mouse blastocyst attachment stability with microscopic analysis to determine embryonic breaching of the Ishikawa cell layer has revealed progressive stages of weak reversible attachment, stable irreversible attachment and subsequent breaching of the epithelium. The rates of stable attachment and breaching in our in-vitro model correlate with in-vitro-matured blastocyst implantation rates in CD1 mice invivo (Schwarzer *et al.*, 2012; Hemkemeyer *et al.*, 2014). Kinetic analysis demonstrated that in-vitro-matured, chemically hatched mouse blastocysts are competent for stable attachment to Ishikawa cells between E5.5 and E6.5, independent of prior co-culture. These findings are consistent with those of a previous study which found that in-vivo-matured, naturally hatched mouse blastocysts co-cultured with Ishikawa cells from E4.5-6.5 attached only after E5.5 and did not require prior co-culture (attachment was scored using a binary scale in the face of medium aspiration by pipette, likely to identify only intermediate and stably attached embryos) (Green *et al.* 2015). Our data also imply that Ishikawa cells are constitutively receptive to mouse blastocyst attachment, as neither co-culture with blastocysts prior to E5.5 nor hormonal stimulation was required to induce an adhesive Ishikawa cell phenotype. Increased mouse blastocyst stable attachment to Ishikawa cells in response to estrogen and progesterone has been reported (Singh *et al.* 2010), however subsequent studies have not used hormones to induce receptivity to rodent and human embryos (Kaneko *et al.* 2011, Kang *et al.* 2014, Green *et al.* 2015). The steroid hormone-mediated shift from non-receptive

303 to receptive endometrium seen in vivo is not replicated by Ishikawa cells in these culture
304 conditions, despite expression of estrogen and progesterone receptors and responsiveness
305 to steroid hormones at the level of gene expression (Tamm-Rosenstein *et al.* 2013).

306 The reversible weak attachment we observed from E4.5-5.5 may mimic the apposition stage
307 of mouse implantation in vivo, which is thought to occur from E4-5 and entail embryo-
308 maternal dialogue leading to locally adhesive LE (Cha & Dey 2014, Aplin & Ruane 2017).
309 Our finding that apposition is required for embryonic breaching of Ishikawa cells suggests
310 that juxtacrine signalling prior to attachment promotes trophoblast penetration of the LE. In
311 contrast, apposition from E4.5-5.5 is not necessary for stable attachment to be achieved in
312 the period from E5.5-6.5. This also suggests that initiating TE differentiation to invasive
313 trophoblast is not required for blastocyst attachment. It is possible that LE-derived signals,
314 such as HB-EGF, trophinin and microRNA hsa-miR-30d (Wang *et al.* 2002, Sugihara *et al.*
315 2007, Vilella *et al.* 2015), promote stable attachment just prior to and during the process.
316 Importantly, this model shows that the attachment competence of mouse blastocysts is
317 regulated during the window of attachment but progress to invasion requires maternal input
318 during the developmental window prior to attachment.

319 We characterised the invasive embryonic cells in our model as TGCs, as they were
320 occasionally bi-/tri-nucleate, exhibited large nuclei with prominent nucleoli, contained
321 pronounced actin stress fibres, and had reduced CDX2 levels (Simmons *et al.* 2007). This
322 mimics the pioneering invasive trophoblast observed in vivo (Dickson 1963), indicating that
323 mouse embryos respond to human Ishikawa cells in ways that resemble implantation in vivo.
324 Moreover, the finding that Ishikawa cells, but not human fibroblasts, can induce specific
325 changes in embryo gene expression suggests that human and mouse LE may share some
326 phenotypic properties, and that Ishikawa cells retain receptive LE-like functions despite
327 being derived from an endometrial adenocarcinoma (Nishida *et al.* 1985).

The mechanism of breaching the Ishikawa cells is not yet clear, but our observations are suggestive of an initial narrow penetration, possibly between cells, followed by trophoblast outgrowth into the surrounding cell layer. LE apoptosis at the embryo attachment site has been proposed as a mechanism of LE penetration in mouse and human (Parr *et al.* 1987, Galan *et al.* 2000, Tu *et al.* 2015), and a recent in-vivo study in mice suggested entosis of LE cells by TE as the mechanism of penetration (Li *et al.* 2015). We did not detect clear morphological signs of either process at sites of embryonic breaching, however induction of TE differentiation to TGC by apposition with LE is not incompatible with these mechanisms. Furthermore, LE induction of TE differentiation to invasive trophoblast recalls human LE-TE interactions from in-vitro studies, whereby homophilic trophinin engagement promotes TE invasion by concomitantly driving LE apoptosis and TE proliferation (Sugihara *et al.* 2007, Tamura *et al.* 2011).

Our investigation implicates the TE differentiation to TGC induced by apposition to LE as a critical step in mouse embryo implantation. Rodent trophoblast differentiation has been widely studied, particularly using blastocyst-derived trophoblast stem cells (TSCs), and transcription factor hierarchies that lead to distinct trophoblast lineages have been determined (Simmons & Cross 2005). *Cdx2*, *Gata3* and *Eomes* are essential to the first lineage allocation to TE, and are associated with TSC-like states, as their downregulation is required for differentiation into all trophoblast lineages (Guzman-Ayala *et al.* 2004, Ralston *et al.* 2010). TGC differentiation from TE/TSCs also requires upregulation of *Hand1*, *Bhlhe40* and *Mdfr* (Cross *et al.* 1995, Kraut *et al.* 1998, Scott *et al.* 2000, Hughes *et al.* 2004). We detected downregulation of both *Cdx2* and *Gata3*, and upregulation of *Hand1* in E5.5 mouse blastocysts specifically after apposition with Ishikawa cells. In addition, anti-CDX2 antibody staining was significantly reduced in embryonic cells breaching the Ishikawa cell layer. We propose that a maternally-derived juxtacrine signal impinges on TE gene expression during apposition to downregulate TSC-like transcription factors and upregulate *Hand1*. We could

not detect *Bhlhe40* or *Mdfi*, suggesting that *Hand1* is an early, maternally-regulated TGC transcription factor.

In humans, the pioneering invasive trophoblast at implantation is thought to be primary syncytium (Hertig *et al.* 1956, Aplin & Ruane 2017), the formation of which is controlled by the expression of the syncytium regulator, transcription factor GCM1 (Yu *et al.* 2002, Liang *et al.* 2010). The existence of an Ishikawa cell-derived signal that promotes mouse TE differentiation in this in-vitro model implicates maternally-regulated induction of trophoblast differentiation as an important mechanism in human embryo implantation. Understanding such events may lead to the development of novel treatments for implantation failure in human ART.

Acknowledgements

We acknowledge staff at the Biological Services Unit, University of Manchester, for their support with animal care. We also acknowledge Berenika Plusa for her advice on mouse embryo culture.

Authors' roles

PTR, SJK, DRB, MW and JDA designed the study, and SJK, DRB, MW and JDA obtained funding. PTR, SCB, RK and JW carried out the experimental work. PTR wrote the paper, which was edited by SJK, DRB, MW and JDA.

Funding

This work was supported by funds from the charities Wellbeing of Women (RG1442) and Diabetes UK (15/0005207), and studentship support for SCB from the Anatomical Society.

Conflict of interest

We declare no conflicts of interest.

References

- Aplin JD** 2006 Embryo implantation: the molecular mechanism remains elusive. *Reprod Biomed Online* **13** 833-839.
- Aplin JD & Kimber SJ** 2004 Trophoblast-uterine interactions at implantation. *Reprod Biol Endocrinol* **2** 48.
- Aplin JD & Ruane PT** 2017 Embryo-epithelium interactions during implantation at a glance. *J Cell Sci* **130** 15-22.
- Armstrong S, Vail A, Mastenbroek S, Jordan V & Farquhar C** 2015 Time-lapse in the IVF-lab: how should we assess potential benefit? *Hum Reprod* **30** 3-8.
- Bedzhov I & Zernicka-Goetz M** 2014 Self-organizing properties of mouse pluripotent cells initiate morphogenesis upon implantation. *Cell* **156** 1032-1044.
- Brison DR, Houghton FD, Falconer D, Roberts SA, Hawkhead J, Humpherson PG, Lieberman BA & Leese HJ** 2004 Identification of viable embryos in IVF by non-invasive measurement of amino acid turnover. *Hum Reprod* **19** 2319-2324.
- Buck VU, Gellersen B, Leube RE & Classen-Linke I** 2015 Interaction of human trophoblast cells with gland-like endometrial spheroids: a model system for trophoblast invasion. *Hum Reprod* **30** 906-916.
- Castelbaum AJ, Ying L, Somkuti SG, Sun J, Ilesanmi AO & Lessey BA** 1997 Characterization of integrin expression in a well differentiated endometrial adenocarcinoma cell line (Ishikawa). *J Clin Endocrinol Metab* **82** 136-142.
- Cha J & Dey SK** 2014 Cadence of procreation: orchestrating embryo-uterine interactions. *Semin Cell Dev Biol* **34** 56-64.

- 400 **Cross JC, Flannery ML, Blonar MA, Steingrimsson E, Jenkins NA, Copeland NG,**
401 **Rutter WJ & Werb Z** 1995 Hxt encodes a basic helix-loop-helix transcription factor that
402 regulates trophoblast cell development. *Development* **121** 2513-2523.
- 403 **Deglincerti A, Croft GF, Pietila LN, Zernicka-Goetz M, Siggia ED & Brivanlou AH** 2016
404 Self-organization of the in vitro attached human embryo. *Nature* **533** 251-254.
- 405 **Dickson AD** 1963 Trophoblastic Giant Cell Transformation of Mouse Blastocysts. *J Reprod*
406 *Fertil* **6** 465-466.
- 407 **Dominguez F, Gadea B, Esteban FJ, Horcajadas JA, Pellicer A & Simon C** 2008
408 Comparative protein-profile analysis of implanted versus non-implanted human blastocysts.
409 *Hum Reprod* **23** 1993-2000.
- 410 **Enders AC** 2007 Implantation in the macaque: expansion of the implantation site during the
411 first week of implantation. *Placenta* **28** 794-802.
- 412 **Enders AC & Schlafke S** 1969 Cytological aspects of trophoblast-uterine interaction in early
413 implantation. *Am J Anat* **125** 1-29.
- 414 **Evans J, Salamonsen LA, Winship A, Menkhorst E, Nie G, Gargett CE & Dimitriadis E**
415 2016 Fertile ground: human endometrial programming and lessons in health and disease.
416 *Nat Rev Endocrinol* **12** 654-667.
- 417 **Ferraretti AP, Goossens V, de Mouzon J, Bhattacharya S, Castilla JA, Korsak V, Kupka**
418 **M, Nygren KG, Nyboe Andersen A, European IVFm, Consortium for European Society**
419 **of Human R & Embryology** 2012 Assisted reproductive technology in Europe, 2008: results
420 generated from European registers by ESHRE. *Hum Reprod* **27** 2571-2584.
- 421 **Filant J & Spencer TE** 2014 Uterine glands: biological roles in conceptus implantation,
422 uterine receptivity and decidualization. *Int J Dev Biol* **58** 107-116.

- 423 **Galan A, Herrer R, Remohi J, Pellicer A & Simon C** 2000 Embryonic regulation of
424 endometrial epithelial apoptosis during human implantation. *Hum Reprod* **15 Suppl 6** 74-80.
- 425 **Glujovsky D, Blake D, Farquhar C & Bardach A** 2012 Cleavage stage versus blastocyst
426 stage embryo transfer in assisted reproductive technology. *Cochrane Database Syst Rev* **7**
427 CD002118.
- 428 **Green CJ, Fraser ST & Day ML** 2015 Insulin-like growth factor 1 increases apical
429 fibronectin in blastocysts to increase blastocyst attachment to endometrial epithelial cells in
430 vitro. *Hum Reprod* **30** 284-298.
- 431 **Guzman-Ayala M, Ben-Haim N, Beck S & Constam DB** 2004 Nodal protein processing
432 and fibroblast growth factor 4 synergize to maintain a trophoblast stem cell
433 microenvironment. *Proc Natl Acad Sci U S A* **101** 15656-15660.
- 434 **Harbottle S, Hughes C, Cutting R, Roberts S, Brison D, Association Of Clinical E &**
435 **The British Fertility S** 2015 Elective Single Embryo Transfer: an update to UK Best
436 Practice Guidelines. *Hum Fertil (Camb)* **18** 165-183.
- 437 **Hemkemeyer SA, Schwarzer C, Boiani M, Ehmcke J, Le Gac S, Schlatt S & Nordhoff V**
438 2014 Effects of embryo culture media do not persist after implantation: a histological study in
439 mice. *Hum Reprod* **29** 220-233.
- 440 **Heneweer C, Schmidt M, Denker HW & Thie M** 2005 Molecular mechanisms in uterine
441 epithelium during trophoblast binding: the role of small GTPase RhoA in human uterine
442 Ishikawa cells. *J Exp Clin Assist Reprod* **2** 4.
- 443 **Hertig AT, Rock J & Adams EC** 1956 A description of 34 human ova within the first 17
444 days of development. *Am J Anat* **98** 435-493.

- 445 **Hughes M, Dobric N, Scott IC, Su L, Starovic M, St-Pierre B, Egan SE, Kingdom JC &**
446 **Cross JC** 2004 The Hand1, Stra13 and Gcm1 transcription factors override FGF signaling to
447 promote terminal differentiation of trophoblast stem cells. *Dev Biol* **271** 26-37.
- 448 **Kaneko Y, Day ML & Murphy CR** 2011 Integrin beta3 in rat blastocysts and epithelial cells
449 is essential for implantation in vitro: studies with Ishikawa cells and small interfering RNA
450 transfection. *Hum Reprod* **26** 1665-1674.
- 451 **Kaneko Y, Murphy CR & Day ML** 2012 Extracellular matrix proteins secreted from both the
452 endometrium and the embryo are required for attachment: a study using a co-culture model
453 of rat blastocysts and Ishikawa cells. *J Morphol* **274** 63-72.
- 454 **Kang YJ, Forbes K, Carver J & Aplin JD** 2014 The role of the osteopontin-integrin
455 alphavbeta3 interaction at implantation: functional analysis using three different in vitro
456 models. *Hum Reprod* **29** 739-749.
- 457 **Kang YJ, Lees M, Matthews LC, Kimber SJ, Forbes K & Aplin JD** 2015 miR-145
458 suppresses embryo-epithelial juxtacrine communication at implantation by modulating
459 maternal IGF1R. *J Cell Sci* **128** 804-814.
- 460 **Knott JG & Paul S** 2014 Transcriptional regulators of the trophoblast lineage in mammals
461 with hemochorial placentation. *Reproduction* **148** R121-136.
- 462 **Kraut N, Snider L, Chen CM, Tapscott SJ & Groudine M** 1998 Requirement of the mouse
463 I-mfa gene for placental development and skeletal patterning. *EMBO J* **17** 6276-6288.
- 464 **Lessey BA, Ilesanmi AO, Castelbaum AJ, Yuan L, Somkuti SG, Chwalisz K &**
465 **Satyaswaroop PG** 1996 Characterization of the functional progesterone receptor in an
466 endometrial adenocarcinoma cell line (Ishikawa): progesterone-induced expression of the
467 alpha1 integrin. *J Steroid Biochem Mol Biol* **59** 31-39.

- 468 **Li Y, Sun X & Dey SK** 2015 Entosis Allows Timely Elimination of the Luminal Epithelial
469 Barrier for Embryo Implantation. *Cell Rep* **11** 358-365.
- 470 **Liang CY, Wang LJ, Chen CP, Chen LF, Chen YH & Chen H** 2010 GCM1 regulation of the
471 expression of syncytin 2 and its cognate receptor MFSD2A in human placenta. *Biol Reprod*
472 **83** 387-395.
- 473 **Liu J, Xu W, Sun T, Wang F, Puschek E, Brigstock D, Wang QT, Davis R & Rappolee**
474 **DA** 2009 Hyperosmolar stress induces global mRNA responses in placental trophoblast
475 stem cells that emulate early post-implantation differentiation. *Placenta* **30** 66-73.
- 476 **Meseguer M, Aplin JD, Caballero-Campo P, O'Connor JE, Martin JC, Remohi J,**
477 **Pellicer A & Simon C** 2001 Human endometrial mucin MUC1 is up-regulated by
478 progesterone and down-regulated in vitro by the human blastocyst. *Biol Reprod* **64** 590-601.
- 479 **Nishida M, Kasahara K, Kaneko M, Iwasaki H & Hayashi K** 1985 [Establishment of a new
480 human endometrial adenocarcinoma cell line, Ishikawa cells, containing estrogen and
481 progesterone receptors]. *Nihon Sanka Fujinka Gakkai Zasshi* **37** 1103-1111.
- 482 **Parr EL, Tung HN & Parr MB** 1987 Apoptosis as the mode of uterine epithelial cell death
483 during embryo implantation in mice and rats. *Biol Reprod* **36** 211-225.
- 484 **Ralston A, Cox BJ, Nishioka N, Sasaki H, Chea E, Rugg-Gunn P, Guo G, Robson P,**
485 **Draper JS & Rossant J** 2010 Gata3 regulates trophoblast development downstream of
486 Tead4 and in parallel to Cdx2. *Development* **137** 395-403.
- 487 **Rossant J & Cross JC** 2001 Placental development: lessons from mouse mutants. *Nat Rev*
488 *Genet* **2** 538-548.
- 489 **Ruiz-Alonso M, Blesa D, Diaz-Gimeno P, Gomez E, Fernandez-Sanchez M, Carranza F,**
490 **Carrera J, Vilella F, Pellicer A & Simon C** 2013 The endometrial receptivity array for

- 491 diagnosis and personalized embryo transfer as a treatment for patients with repeated
492 implantation failure. *Fertil Steril* **100** 818-824.
- 493 **Salamonsen LA, Edgell T, Rombauts LJ, Stephens AN, Robertson DM, Rainczuk A,**
494 **Nie G & Hannan NJ** 2013 Proteomics of the human endometrium and uterine fluid: a
495 pathway to biomarker discovery. *Fertil Steril* **99** 1086-1092.
- 496 **Schwarzer C, Esteves TC, Arauzo-Bravo MJ, Le Gac S, Nordhoff V, Schlatt S & Boiani**
497 **M** 2012 ART culture conditions change the probability of mouse embryo gestation through
498 defined cellular and molecular responses. *Hum Reprod* **27** 2627-2640.
- 499 **Scott IC, Anson-Cartwright L, Riley P, Reda D & Cross JC** 2000 The HAND1 basic helix-
500 loop-helix transcription factor regulates trophoblast differentiation via multiple mechanisms.
501 *Mol Cell Biol* **20** 530-541.
- 502 **Shahbazi MN, Jedrusik A, Vuoristo S, Recher G, Hupalowska A, Bolton V, Fogarty NM,**
503 **Campbell A, Devito LG, Ilic D, Khalaf Y, Niakan KK, Fishel S & Zernicka-Goetz M** 2016
504 Self-organization of the human embryo in the absence of maternal tissues. *Nat Cell Biol* **18**
505 700-708.
- 506 **Simmons DG & Cross JC** 2005 Determinants of trophoblast lineage and cell subtype
507 specification in the mouse placenta. *Dev Biol* **284** 12-24.
- 508 **Simmons DG, Fortier AL & Cross JC** 2007 Diverse subtypes and developmental origins of
509 trophoblast giant cells in the mouse placenta. *Dev Biol* **304** 567-578.
- 510 **Simon C, Sakkas D, Gardner DK & Critchley HO** 2015 Biomarkers in reproductive
511 medicine: the quest for new answers. *Hum Reprod Update* **21** 695-697.
- 512 **Singh H & Aplin JD** 2014 Endometrial apical glycoproteomic analysis reveals roles for
513 cadherin 6, desmoglein-2 and plexin b2 in epithelial integrity. *Mol Hum Reprod* **21** 81-94.

- 514 **Singh H, Nardo L, Kimber SJ & Aplin JD** 2010 Early stages of implantation as revealed by
515 an in vitro model. *Reproduction* **139** 905-914.
- 516 **Spandidos A, Wang X, Wang H & Seed B** 2010 PrimerBank: a resource of human and
517 mouse PCR primer pairs for gene expression detection and quantification. *Nucleic Acids*
518 *Res* **38** D792-799.
- 519 **Sugihara K, Sugiyama D, Byrne J, Wolf DP, Lowitz KP, Kobayashi Y, Kabir-Salmani M,**
520 **Nadano D, Aoki D, Nozawa S, Nakayama J, Mustelin T, Ruoslahti E, Yamaguchi N &**
521 **Fukuda MN** 2007 Trophoblast cell activation by trophinin ligation is implicated in human
522 embryo implantation. *Proc Natl Acad Sci U S A* **104** 3799-3804.
- 523 **Tamm-Rosenstein K, Simm J, Suhorutshenko M, Salumets A & Metsis M** 2013
524 Changes in the transcriptome of the human endometrial Ishikawa cancer cell line induced by
525 estrogen, progesterone, tamoxifen, and mifepristone (RU486) as detected by RNA-
526 sequencing. *PLoS One* **8** e68907.
- 527 **Tamura N, Sugihara K, Akama TO & Fukuda MN** 2011 Trophinin-mediated cell adhesion
528 induces apoptosis of human endometrial epithelial cells through PKC-delta. *Cell Cycle* **10**
529 135-143.
- 530 **Tu Z, Wang Q, Cui T, Wang J, Ran H, Bao H, Lu J, Wang B, Lydon JP, DeMayo F,**
531 **Zhang S, Kong S, Wu X & Wang H** 2015 Uterine RAC1 via Pak1-ERM signaling directs
532 normal luminal epithelial integrity conducive to on-time embryo implantation in mice. *Cell*
533 *Death Differ* **23** 169-181.
- 534 **Vilella F, Moreno-Moya JM, Balaguer N, Grasso A, Herrero M, Martinez S, Marcilla A &**
535 **Simon C** 2015 Hsa-miR-30d, secreted by the human endometrium, is taken up by the pre-
536 implantation embryo and might modify its transcriptome. *Development* **142** 3210-3221.
- 537 **Wang H & Dey SK** 2006 Roadmap to embryo implantation: clues from mouse models. *Nat*
538 *Rev Genet* **7** 185-199.

539 **Wang J, Mayernik L & Armant DR** 2002 Integrin signaling regulates blastocyst adhesion to
540 fibronectin at implantation: intracellular calcium transients and vesicle trafficking in primary
541 trophoblast cells. *Dev Biol* **245** 270-279.

542 **Wang J, Mayernik L, Schultz JF & Armant DR** 2000 Acceleration of trophoblast
543 differentiation by heparin-binding EGF-like growth factor is dependent on the stage-specific
544 activation of calcium influx by ErbB receptors in developing mouse blastocysts. *Development*
545 **127** 33-44.

546 **Yu C, Shen K, Lin M, Chen P, Lin C, Chang GD & Chen H** 2002 GCMA regulates the
547 syncytin-mediated trophoblastic fusion. *J Biol Chem* **277** 50062-50068.

548 **Table, movie and figure legends**

549 **Table 1** Mouse gene names and corresponding forward and reverse primer sequences.
550 Primer sequences were obtained from PrimerBank (Spandidos et al. 2010), with the
551 following references: *Cdx2* 31560722a1; *Gata3* 6679951a1; *Eomes* 26354683a1; *Hand1*
552 118130896c3; *Gata2* 226530725c1; *Hes1* 6680205a1; *Gapdh* 126012538c1; *Ywhaz*
553 359385697c3.

554 **Movie 1** Differential interference contrast imaging at 14 frames per second of E5.5+4h
555 mouse embryos after 28h co-culture with Ishikawa cells exhibiting, upon agitation, weak
556 attachment (major oscillation about an attachment point, left embryo), intermediate
557 attachment (minor oscillation, right embryo) and stable attachment (no oscillation, middle
558 embryo) to Ishikawa cells. Scale bar 100µm.

559 **Figure 1 A** Schematic to illustrate the criteria for the 4-point grading of the stability of mouse
560 blastocyst attachment to Ishikawa cells; [1] = not attached, [2] = weakly attached, [3] =
561 intermediately attached, [4] = stably attached. **B** Hatched E4.5 mouse blastocysts were co-
562 cultured with Ishikawa cells for 48h and the stability of attachment was scored (1-4) every 4h
563 from 0-12h, 24-36h, and at 48h. The mean attachment stability scores, from four

independent experiments using 102 blastocysts, are presented. Error bars +/- standard error of the mean (SEM). **C** The mean attachment stability score was calculated from the data in B. Error bars +/- SEM. **D** Phase contrast images of the mouse blastocyst-Ishikawa cell co-culture at 4h, 8h, 12h, 28h, 32h and 36h. Blastocysts are indicated by arrows and the attachment score for each blastocyst is displayed. Scale bars 200µm. **E** The cumulative frequency of the time at which the final position of attachment was reached was plotted alongside the percentage of blastocysts scored as intermediately or stably attached (mean +/- SEM from three independent experiments using 84 blastocysts). **F** Ishikawa cells were pre-treated with 10nM 17β-estradiol 48h prior to co-culture, then with 10nM 17β-estradiol and 1µM progesterone 24h prior to co-culture and during co-culture with hatched E4.5 mouse blastocysts. The attachment stability of mouse blastocysts was scored every 4h from 0-12h, 24-36h, and at 48h. Blastocysts scored as intermediately and stably attached were added together and the mean from three independent experiments using 72 blastocysts was plotted +/- SEM.

Figure 2 Attached embryos not breaching (**A**) and breaching (**B**) Ishikawa cell layers were optically sectioned after PFA fixation and incubation with phalloidin to label actin fibres (red) and DAPI to label nuclei (blue). Invasive cells of breaching embryos exhibit binuclearity (thick arrow), prominent nucleosomes (arrowhead), and actin stress fibres (thin arrow). Upper/lower section refers to depth of the optical section in the Z-axis. Scale bars 20µm. **C** The size of invasive embryonic nuclei in the lower optical section was measured and compared to the size of outer, peripheral embryonic nuclei, presumed to be TE and not inner cell mass, in the corresponding upper section; 309 nuclei were measured in 12 embryos and mean nuclear size was plotted +/- SEM. *** t-test $p < 0.001$. Breaching embryos were fixed and labelled with anti-CDX2 antibody (**D**), rabbit isotype control antibody (**D** inset), anti-GATA3 antibody (**E**), or mouse isotype control antibody (**E** inset), as well as phalloidin and DAPI. Upper/lower section refers to depth of the optical section in the Z-axis. Scale bars 20µm. **F** The intensity of CDX2 and GATA3 fluorescence labelling was measured in invasive

embryonic nuclei in the lower optical section and compared to that of outer embryonic nuclei, presumed to be TE and not inner cell mass, in the corresponding upper section; 158 nuclei were measured in five embryos labelled with anti-CDX2 and 226 nuclei were measured in five embryos labelled with anti-GATA3. The mean ratio of invasive nuclei to upper TE nuclei fluorescence intensity was plotted $\pm$ SEM. ** t-test $p < 0.01$, NS not significant.

Figure 3 A Schematic representing embryo-Ishikawa cell co-culture experimental design. **B** Hatched E4.5 mouse blastocysts were either co-cultured with Ishikawa cells to E6.5 or cultured alone in co-culture media to E5.5 before co-culture with Ishikawa cells to E6.5 or E7.5. The stability of attachment was scored every 4h from E5.5-6 and at E6.5. Mean percentage blastocysts scored as intermediately or stably attached from six independent experiments using 149 blastocysts was plotted $\pm$ SEM. **C** Attached embryos co-cultured from E4.5-6.5, E5.5-6.5 and E5.5-7.5, were fixed, labelled with phalloidin and DAPI and optically sectioned to determine whether embryonic cells had breached the Ishikawa cells. The mean proportion of attached embryos to have breached the Ishikawa cells was plotted $\pm$ SEM. ** ANOVA $p < 0.01$, *** ANOVA $p < 0.001$, demonstrating significant difference from co-culture E4.5-6.5.

Figure 4 A Schematic representing embryo-Ishikawa cell co-culture experimental design. **B** Hatched E4.5 mouse blastocysts were co-cultured directly with Ishikawa cells from E4.5-6.5, co-cultured directly with Ishikawa cells to E5.5 before transfer to fresh Ishikawa cells and co-culture to E6.5, or co-cultured in a transwell above Ishikawa cells to E5.5 before transfer to fresh Ishikawa cells and co-culture to E6.5. The stability of attachment was scored every 4h from E5.5-6 and at E6.5. Mean percentage blastocysts scored as intermediately or stably attached from five independent experiments using 157 blastocysts was plotted $\pm$ SEM. **C** Attached embryos were fixed, labelled with phalloidin and DAPI and optically sectioned to determine whether embryonic cells have breached the Ishikawa cells. The mean proportion of attached embryos to have breached the Ishikawa cells $\pm$ SEM. ** ANOVA $p < 0.01$, demonstrating significant difference from direct co-culture E4.5-6.5.

618 **Figure 5** Hatched E4.5 mouse blastocysts were cultured alone, co-cultured directly with
619 Ishikawa cells, co-cultured in a transwell above Ishikawa cells or co-cultured directly with
620 human fibroblasts as a control cell type, to E5.5 before retrieval and lysis for RNA isolation
621 and gene expression analysis by qPCR. Data is presented as expression relative to
622 reference genes ($2^{-\text{ct}}$, relative to geometric mean of *Gapdh* and *Ywhaz*) of **A** *Cdx2*, **B** *Gata3*,
623 **C** *Eomes*, **D** *Hand1*, **E** *Gata2*, and **F** *Hes1* in embryos cultured in the indicated conditions
624 (mean $\pm$ SEM of five independent experiments using 200 embryos). * ANOVA $p < 0.05$,
625 demonstrating significant difference from no co-culture E4.5-5.5.