

## Central Lancashire Online Knowledge (CLOK)

Title	Facilitators and barriers to participation in mental well-being programs by older Australians with vision impairment: community and stakeholder perspectives
Type	Article
URL	<a href="https://clock.uclan.ac.uk/33654/">https://clock.uclan.ac.uk/33654/</a>
DOI	<a href="https://doi.org/10.1038/s41433-020-0992-z">https://doi.org/10.1038/s41433-020-0992-z</a>
Date	2020
Citation	Dillon, Lisa, Tang, Diana, Liew, Gerald, Hackett, Maree, Craig, Ashley, Gopinath, Bamini and Keay, Lisa (2020) Facilitators and barriers to participation in mental well-being programs by older Australians with vision impairment: community and stakeholder perspectives. Eye, 34 (7). pp. 1287-1295. ISSN 0950-222X
Creators	Dillon, Lisa, Tang, Diana, Liew, Gerald, Hackett, Maree, Craig, Ashley, Gopinath, Bamini and Keay, Lisa

It is advisable to refer to the publisher's version if you intend to cite from the work.  
<https://doi.org/10.1038/s41433-020-0992-z>

For information about Research at UCLan please go to <http://www.uclan.ac.uk/research/>

All outputs in CLOK are protected by Intellectual Property Rights law, including Copyright law. Copyright, IPR and Moral Rights for the works on this site are retained by the individual authors and/or other copyright owners. Terms and conditions for use of this material are defined in the <http://clock.uclan.ac.uk/policies/>

1   **Title**

2   Facilitators and barriers to participation in mental wellbeing programs by older Australians  
3   with vision impairment: community and stakeholder perspectives

4   **Running title**

5   Vision impairment and mental wellbeing programs

6   **Authors**

7   Lisa Dillon<sup>a b\*</sup> PhD, lisa.dillon1@unsw.edu.au

8   Diana Tang<sup>c</sup> MND, diana.tang@sydney.edu.au

9   Gerald Liew<sup>c</sup> PhD MBBS MMED, gerald.liew@sydney.edu.au

10   Maree Hackett<sup>b d</sup> PhD, mhackett@georgeinstitute.org.au

11   Ashley Craig<sup>e</sup> PhD, a.craig@sydney.edu.au

12   Bamini Gopinath<sup>c</sup> PhD, bamini.gopinath@sydney.edu.au

13   Lisa Keay<sup>a b</sup> PhD, l.keay@unsw.edu.au

14   a School of Optometry and Vision Science, Faculty of Science, UNSW Sydney, Postal:  
15   School of Optometry and Vision Science, The University of New South Wales, Sydney  
16   UNSW NSW 2052, Australia

17   b The George Institute for Global Health, Faculty of Medicine, UNSW Sydney, Postal: PO  
18   Box M201 Missenden Rd NSW 2050, Australia

19   c Centre for Vision Research, Department of Ophthalmology, The Westmead Institute for  
20   Medical Research, The University of Sydney, Sydney, Australia, Postal: Centre for Vision  
21   Research, 176 Hawkesbury Road, Westmead NSW 2145

22   d Faculty of Health and Wellbeing, University of Central Lancashire, Preston, United  
23   Kingdom, Postal: University of Central Lancashire, Preston, Lancashire PR1 2HE

24 e John Walsh Centre for Rehabilitation Research, Northern Clinical School, Faculty of  
25 Medicine and Health, The University of Sydney, Postal: Kolling Institute of Medical  
26 Research, RNSH, St. Leonards, NSW, 2650

27 \*corresponding author

## 28 **Conflict of interest**

29 The authors declare no conflict of interest.

## 30 **Funding**

31 This research was funded by the UNSW Sydney-University of Sydney: Mental Health and  
32 Wellbeing – Early Intervention and Prevention (Older People) Seed Funding Scheme

## Abstract

**Objective:** Older adults with vision impairment experience high rates of mental health problems, but very few access psychological support. We investigated community and stakeholder perspectives of the barriers and facilitators to participation in mental wellbeing programs for older adults with vision impairment.

**Methods:** Adults aged  $\geq 50$  years with vision impairment (community) were recruited from the client database, and low vision rehabilitation (LVR) professionals (stakeholders) from staff of a LVR provider. Participants completed one-on-one semi-structured interviews which were designed and analysed using behaviour change theory.

**Results:** Twenty-nine participants were interviewed; 16 community members and 13 stakeholders. Both groups cited mental health problems as a major concern, with many stakeholders reporting the grief and distress associated with vision loss experienced by their clients as having a negative impact on their mental and physical health. Major barriers to participation in mental wellbeing programs included a lack of awareness and difficulties accessing such programs, with stakeholders adding that their clients' lack of insight into their own mental health problems may reduce motivation to participate. Facilitators to participation in programs included the appeal of social interaction and inspirational speakers. An appropriate intervention could overcome these barriers, or enhance participation through education, persuasion, incentivisation, modeling, environmental restructuring, training, and enablement.

**Conclusions:** While barriers were discussed more than facilitators to participation, there was general support for mental wellbeing programs. This study provides guidance from stakeholders for the development of mental wellbeing programs to address mental health problems in the growing number of older adults with vision impairment.

## Introduction

Globally, it is estimated that the number of people living with vision impairment (encompassing low vision or blindness that cannot be corrected with glasses or surgery) will rise from 38.5 million in 2020 to 115 million people by 2050; most of which (78%) are aged 50 years or older [1]. Older age and vision impairment are associated with an increased risk of developing a mental health condition, particularly depression [2]. An estimated 69,519 non-Indigenous Australians aged 50 years or above and 4,282 Indigenous Australians aged 40 years or above were living with irreversible vision impairment in 2016 [3]; all at consequent risk of mental health problems.

Older adults with vision impairment are approximately three times more likely to experience significant depressive symptoms, compared with those without vision impairment [2, 4]. This is attributed to the impacts of vision impairment on functional capacity and activities of daily living [5-8]. Research suggests that low vision rehabilitation (LVR) services, and psychological interventions involving self-management and problem-solving, may be effective in reducing depressive symptoms and depressive disorders among adults with vision impairment [9-12]. However, although 70% of adults with vision impairment report wanting psychological support, only 9% report receiving it [13], and it remains unclear why uptake of support is so low.

One study of LVR professionals' (hereafter stakeholders) perspectives of a mental wellbeing program found a number of barriers to the problem-solving treatment for primary care (PST-PC) being delivered in a LVR setting, that could be overcome through professional training, support and improved screening [14]. The only other study reported on the reasons older adults with vision impairment withdrew early from PST-PC, and found responses ranging from the program being perceived as not relevant, to the program goals being achieved early [10]. Given this scant literature, the aim of this qualitative study is to investigate the barriers and facilitators to participation in a mental wellbeing programs, from the perspectives of

community-dwelling older adults with vision impairment (community members) and stakeholders.

## Subjects and Methods

This qualitative study involved semi-structured, one-on-one telephone or face-to-face interviews with community members and stakeholders, including orientation and mobility specialists and optometrists. This study was conducted within New South Wales (NSW) and the Australian Capital Territory (ACT), Australia.

Community members and stakeholders were recruited using purposive sampling through invitation letters and follow-up phone calls between August and October 2019. To address the study aim, we expected to recruit up to 20 community members and up to 15 stakeholders for interview, or until data was saturated in each group, which was assessed by researchers (LD, LK) at regular meetings. Community members were recruited from the client database of an Australian vision rehabilitation organisation, Guide Dogs NSW/ACT. Clients ~~that~~ who had consented to be contacted regarding research were sent an email and invited to contact the research team directly to express interest in the study. Participants were required to be aged 50 years and older and speak conversational English. Stakeholder participants were employed in a client-facing capacity by Guide Dogs NSW/ACT, and were similarly sent an email and invited to contact the research team directly to express interest in the study.

The behaviour change wheel [15] was used as a framework to design the interview guides (Supplementary File 1 and 2), and consequent analysis of the data. The behaviour of interest was participation in a mental wellbeing program. Specifically, the wheel uses the COM-B model to explain an individual's propensity to adopt a behaviour (see the inner wheel of Figure 1). Behaviour can be in turn targeted by nine intervention functions: coercion, education, enablement, environmental restructuring, incentivisation, modelling, persuasion,

restriction, training [15]. The wheel also includes policy categories which can impact the intervention functions; however, these were considered outside the scope of this study.

Semi-structured interviews, lasting 15-45 minutes, were conducted by two Master of Clinical Optometry students (AW, RM) and a Faculty of Medicine and Health PhD student (DT). The students were given initial training and ongoing support by two experienced public health qualitative researchers (LD, LK). Interviews were audio recorded using digital recorders, transcribed verbatim, and analysed using NVivo software, using deductive analysis [16], following the COM-B and intervention functions of the behaviour change wheel [15]. We took an iterative approach to data analysis [17], whereby data were revisited, coded, and themes discussed many times to ensure analytic reflexivity. Transcripts were coded separately by two of the students (AW, RM). Initial coding was then presented and critiqued in a meeting including all three students, chaired by LD. Two of the students (AW, RM) then collaborated to establish themes under each subset of the COM-B model, which were discussed and agreed upon at regular fortnightly meetings with LD and LK. Themes were then coded using intervention functions, and categorised as barriers or facilitators to participation in a mental wellbeing program by LD and DT.

#### **Patient and public involvement**

No patient under medical care or members of the general public were involved in the design, recruitment or conduct of the study. However, there will be formal engagement with the LVR provider's advisory panel to discuss the next stages of developing and implementing a mental wellbeing program. Those participants who requested feedback will be informed via email or telephone call about the results.

#### **Ethics approval**

Ethics approval was granted by the University of New South Wales Human Research Ethics Committee (HC190356). A participation information statement was sent to interested participants, and oral informed consent was gained before commencement of the interview.

The study is reported in line with the COREQ statement[18], supporting transparency in reporting of qualitative research.

## Results

Of the 35 community members and stakeholders invited, 29 (83%, 16 community members and 13 stakeholders) completed an interview. The 13 stakeholders included 12 orientation and mobility specialists and one optometrist. All stakeholder interviews were conducted over the phone, while seven of the community participants requested face-to-face interviews in their homes due to difficulties with hearing and/or accessing a phone.

We identified twelve themes (Figure 1); eight represented barriers, and four facilitators to mental wellbeing program participation. Illustrative quotes associated with each theme are presented below, with intervention functions in brackets next to the COM-B component.

Participants are identified as C for community members and S for stakeholders, followed by an identification number.

Both participant groups showed a high level of interest through long and engaging discussions around the mental wellbeing of people with vision impairment. Stakeholders expressed that many of their clients have concerns regarding their mental wellbeing, indicating the importance of this topic and need for such programs:

*"I'm finding most of the clients that I...work with, express that they have anxiety, depression, or have had nervous breakdowns, or are currently having mental health issues." (S002)*

## Barriers

### ***Physical capability (Enablement, Training)***

#### *Mobility impacts participation*

Participants reported difficulties in mobility as a result of their vision loss which prevented them from engaging with their community and/or programs:



159       *"Because you can't go out. That's the biggest problem..." (C008)*

160       *"So, you don't tread on little kids or get tangled up in dog leads...that's always in the*  
161       *back of your mind. I think I'll just stay at home." (C004)*

162   In particular, there were concerns regarding loss of the ability to drive:

163       *"You're not going to get them in a group scenario, but also they can't drive so they can't*  
164       *get to anywhere." (S008)*

165   Some participants also found that their mobility limitations and location of residence  
166   interfered with their ability to connect with other individuals with vision impairment and  
167   contributed to feelings of isolation:

168       *"I would love to have a group of other people who were also vision impaired, but they*  
169       *seem to be scattered all over the country." (C020)*

170   *Vision loss impacts everyday activities*

171   The inability to accomplish activities of daily living deterred participation in programs and  
172   other activities:

173       *"Because of their vision impairment, I don't think they integrate as much with others in*  
174       *their community." (S009)*

175       *"I can't watch TV and I do like TV actually. I can't read anything anymore and I used to*  
176       *love a newspaper." (C020)*

177   ***Psychological capability (Education, Training, Enablement)***

178   *Ripple effects*

179   Stakeholders expressed concerns about mental health problems extending into other  
180   aspects of their client's lives and preventing participation in mental wellbeing programs:

181       *"Stress and mental health, the physiological changes to the body... impacts on people's*  
182       *functional vision. There's all these ripple effects if the mental health component is not*  
183       *addressed."* (S006)

184       *"I'd be the only person they'd see that week and maybe they'd have a cry...need to work*  
185       *through the issues they have before they can start doing routes and getting out in their*  
186       *community."* (S015)

#### 187   *Self-perception of mental health*

188   The acknowledgement of having a mental health problem varied among participants:

189       *"One very good friend who's also legally blind...when I bring up the subject with him he*  
190       *says, Oh, better not even to think about it. And he's not really getting all the services*  
191       *that he should be getting as a blind person."* (C004)

192       *"Has it affected me? I lost my licence. I'm pretty much housebound. My doctor wants*  
193       *me to see... a psychologist. And I said, No, I'm not that bad."* (C008)

194       *"I try to prepare myself for the future, but I don't think it's affected my mental health, as*  
195       *far as others are concerned anyway."* (C009)

#### 196   ***Physical opportunity (Environmental restructuring, Enablement)***

##### 197   *Facilitation of services*

198   Participants expressed concerns about their ability to access mental wellbeing programs  
199   following their vision loss:

200       *"When I lost my sight I had to really scabble and... call for information."* (C007)

201       *"If there is support, they [support provider] could call our clients ...[so] the client doesn't*  
202       *have to initiate [seeking services] themselves...because I have a lot of clients who once*  
203       *they lose their vision, they're not able to navigate a phone."* (S003)

204 Stakeholders also expressed their concerns about how they can best assist their clients to  
205 receive support for their mental health problems:

206 *"If we had a script or a proper format that once a client has identified that they've got*  
207 *mental health issues that we should say because you've mentioned that to us can we*  
208 *write some information down and pass it on to someone to suggest help or at least to*  
209 *refer."* (S013)

210 *Lack of awareness of services*

211 Community and stakeholders were unaware of current mental wellbeing programs targeted  
212 at Australians with vision impairment:

213 *"I'm not aware of any program for... people with vision impairment."* (S012)

214 *"In a major Sydney hospital and they didn't ever say to me that I needed to be referred*  
215 *to Vision Australia or Guide Dogs or anywhere like that. And I just said, Okay, well, I can*  
216 *do this. But the trouble is I couldn't see."* (C005)

217 ***Social opportunity (Environmental restructuring, Enablement, Modelling)***

218 *Stigma about mental health problems*

219 Participants described the negative stigma surrounding mental health issues as a barrier to  
220 participation:

221 *"...it doesn't seem to be talked about a lot."* (C015)

222 *"People have a perception, there's still a stigma around mental illness, so a lot of them*  
223 *might not disclose."* (S005)

224 *"I would say don't call it a mental health program."* (C005)

225 ***Reflective motivation (Education, Persuasion, Incentivisation)***

226 *Grief associated with vision loss*

227 Feelings of grief and beliefs regarding vision loss can make participation and engagement  
228 unmotivating:

229 *"If a person is depressed and feels like they can't do anything, then it's likely they're not*  
230 *going to go into orientation or mobility training with high expectations, and a good*  
231 *learning frame of mind."* (S012)

## 232 **Facilitators**

### 233 ***Social opportunity (Environmental restructuring, Enablement, Modelling)***

#### 234 *Community connections*

235 The desire for a connection to the community particularly with those experiencing similar  
236 vision loss was described by participants:

237 *"...having a group that comes together that has a focus on adapting to change and*  
238 *understanding... I think is fairly powerful."* (C015)

239 *"Although technology and online stuff is cool and innovative...it sort of takes away from*  
240 *that human connection that you have when you sit in the same room as somebody."*  
241 *(S015)*

242 *"There's others there to talk to and pass the time of day with. It's not a discussion group,*  
243 *it's just to pass the time. You know, with the social aspect of it, but they're getting to*  
244 *know me."* (C003)

245 *"For so long I just thought our family was a family of freaks because not knowing of*  
246 *anyone else that had it...30 years ago I joined up with the Nepean Blind Sports Club...I*  
247 *met a couple of people with the same condition and... went to social events out there."*  
248 *(C012)*

249 *Who can facilitate mental wellbeing programs*

250 Participants also expressed their opinions about who they think would be best suited to  
251 deliver low vision mental wellbeing programs. Different levels of expertise were articulated  
252 ranging from lay-facilitators to mental health professionals like psychologists:

253 *“Anybody who has got group-based skills and some level of working with groups.”*  
254 *(C005)*

255 *“I always feel that if someone’s been through a situation they’re the ones I think are the*  
256 *better ones.” (C012)*

257 *A consultant psychologist with a good knowledge of grief... it would be great to be able*  
258 *to run scenarios by that person. And say, this is what I’m dealing with, or this is what I’ve*  
259 *done. It would be lovely to be able to say to clients If you want to talk more about this,*  
260 *we have a psychologist on staff that might be able to give you a ring.” (S004)*

261 *“It would be good if we had one person that they could make a call to that has more*  
262 *specific information and then can put them in touch with people in their own area... or*  
263 *what organisations are close to them.” (S013)*

#### 264 **Automatic motivation (Modelling, Enablement)**

265 *Inspiration/role model*

266 Some participants stated how a role model is an inspiring motivation for participating in  
267 mental wellbeing programs:

268 *“Those have been some of our better meetings when we’ve been inspired by others in*  
269 *other words.” (C009)*

270 *“...hearing people’s stories about how they might have done it is really powerful for*  
271 *people who might be going through that part of the process.” (C005)*

#### 272 **Reflective motivation (Education, Persuasion, Incentivisation)**

273 *Self-efficacy*

274 Many participants believed they were self-sufficient and capable of handling their own  
275 issues:

276 *"I can't change it, I've just got to learn to adjust to it and that's just going to be my life."*  
277 (C011)

278 *"I don't feel sorry for myself, there's no point. I've got two legs, I can walk, so life's*  
279 *good."* (C022)

280

## 281 **Discussion**

282 Consistent with evidence of the high burden of mental health problems in older adults with  
283 vision impairment [2, 4], we found strong interest in developing mental wellbeing programs  
284 among community members and stakeholders. The interviews revealed that both groups  
285 acknowledged the substantial negative impacts of mental health problems. However, both  
286 groups were unaware of any mental wellbeing programs specific to people with vision  
287 impairment, thus, confirming the importance of developing a mental wellbeing program  
288 tailored to this population group.

289 We identified two times more barriers than facilitators (i.e. eight vs four) to effective  
290 participation in a mental wellbeing program. The intervention function *Enablement* was  
291 relevant to all three facilitators, and the majority of barriers. Enablement refers to increasing  
292 means, and reducing barriers [15], and is key to designing a program for this population. A  
293 major concern related to Enablement, along with Environmental Restructuring (changing the  
294 physical or social context [15]), was functional disability. Many community members felt that  
295 vision loss limited their capability and opportunity to access programs due to difficulty  
296 reading or finding information, and travelling to where programs were held. Travel concerns  
297 are a common barrier in this population [19], and is influenced by the severity of vision  
298 impairment, distance to program location, and availability of support people.

299 Recommendations to overcome these barriers, as they relate to intervention functions [15],

include: producing material in larger, easy-to-read print (*Enablement*); LVR professionals telling community members what options are available (*Education*; increasing knowledge or understanding); and program organisers making community transport available (*Environmental Restructuring*) or provide orientation and mobility services (*Training*; imparting skills), if travel is required.

In addition to physical limitations, the acknowledgement of having a mental health problem influenced participation and varied among participants. Some mentioning conditions like depression and anxiety, others downplaying any concerns and associating it with complaining, and the remainder stating that their vision loss did not have any impact on their mental health. The intervention functions of *Modelling* (an example for people to aspire to or imitate) and *Education* are particularly relevant to alleviate these concerns and key to tackling the barriers of social stigma and self-perception. Nyman et al. [20] identified that vision impairment can impact psychosocial well-being through social isolation; while acceptance of vision loss, and social support were facilitators of adjustment. We also identified self-efficacy as a facilitator, which has been shown to lead to good mental health outcomes in other contexts [21]. Given there was general consensus among both groups of the benefits of social connection and positive role models, it is recommended interventions include this in their design.

A pilot feasibility study in Australia recently investigated the delivery of PST-PC by LVR professionals to older adults with vision impairment who met the criteria for mild symptoms of depression [10]. PST-PC, based on Cognitive Behavioural Therapy (CBT) principles, is a low-intensity approach designed to assist with functional adjustment, resilience building and generalised well-being. PST-PC can be delivered by non-mental health professionals, face-to-face, as well as over the phone, or over video conference, significantly increasing its reach. Holloway et al. [10] found that those that who stayed in the study had significant reductions in depressive symptoms, and improvements in health-related quality-of-life and problem-focused coping. Those who withdrew from the study early were typically older, with

complex needs, as has been found in other studies [22-24]. Thus, it is critical that any future interventions must consider methods to retain adherence of older participants with competing health priorities.

Varying levels of mental health problems and requirements for support were reported by community members as well as stakeholders about their clients. A stepped-care approach may be the most efficient in this context, as not all clients require the same type or intensity of intervention [25]. Stepped-care comprises different intervention components, with the idea that if the first, less intensive step does not lead to a reduction in symptoms, then a person moves to the next step, consisting of more intensive and potentially more expensive interventions [9]. Several randomised controlled trials conducted outside the field of low vision have found that a stepped-care approach can be effective in minimising depression and/or anxiety [26, 27], and has been endorsed to address depression in older adults in clinical guidelines, such as the UK NICE guidelines [28]. The ideas for interventions identified in this study could be integrated within a stepped-care approach, including social groups, psychologist referral and PST-PC.

## **Strengths and limitations**

This is the first study to explore the perspectives of older adults with vision impairment and client-facing professionals involved in LVR regarding mental health problems. The inclusion of client-facing professionals provides essential insight needed to design a stakeholder-driven intervention. A second strength is the semi-structured nature of the interviews which allowed for tailored discussions to explore each participant's personal outlook and interpretation of the impacts of mental health problems and vision impairment. However, this also resulted in interviews of varying length, and occasional tangents in conversation. Despite compelling findings, the authors acknowledge that the study is limited to one community organisation and therefore the presented results are only a preliminary indicator of the mental health perspectives in this population group. Moreover, community member participants were recruited through purposive sampling from the client base of one



Australian LVR provider, and specifically only those clients who had agreed to be contacted  
about participating in research, and had also agreed to participate in this particular study.  
This potential selection bias may have influenced our results, and may limit the  
generalisability of findings to those not associated with an Australian LVR provider, or those  
not interested in participating in research or this particular study. Similarly, stakeholder  
participants were orientation and mobility specialists and one optometrist from the one  
Australian LVR organisation, which may limit the generalisability of findings to these LVR  
providers in other organisations, or other professions involved in LVR, such as occupational  
therapists, orthoptists, assistive technology specialists and social workers. Nonetheless, as  
data saturation was achieved, we are confident the range of themes and corresponding  
intervention functions suitably address the study aim. ~~Moreover, as participants were~~  
~~recruited through purposive sampling, there may be influences of selection bias influencing~~  
~~results and perhaps likely underestimating the mental health problems of individuals not~~  
~~associated with an LVR provider.~~ Furthermore, information on visual acuity, cause of vision  
loss and presence of comorbidities was not available for collection. This information may  
have been valuable in terms of interpreting the qualitative data.

This study has several key implications for clinical practice. First, improving the mental  
wellbeing of visually impaired older adults is a priority. We postulate this may enhance their  
participation in other programs, such as orientation and mobility, and this in turn could  
improve their physical health, community connections, and overall well-being. Second, LVR  
professionals, such as orientation and mobility specialists, and optometrists, are in a unique  
position to identify those at risk of, or already experiencing mental health problems and can  
refer as necessary, if provided with the necessary training. Third, older adults with vision  
impairment may benefit from appropriate education and training to better support them as  
they adjust to living with a vision impairment. We suggest this may be in the form of  
evidence-based education regarding their vision diagnosis and prognosis, or psychological  
support from an appropriately trained mental health practitioner. Fourth, providing older

381 adults with vision impairment the opportunity to connect with others in similar situations, as  
382 well as those who are living well with vision impairment, is likely to improve their community  
383 connections and social engagement. As a next step, the suggestions for future interventions  
384 collected here will be presented through round tables with older adults with vision  
385 impairment and service providers. This will ensure that any developed strategies are  
386 practical and acceptable to the community and stakeholders, with the aim to improve the  
387 ultimate adoption and scalability of a mental wellbeing program.

## 388    **Acknowledgments**

389    We would like to acknowledge students Rachel Muyco and Aimee Wassermann for  
390    conducting many of the semi-structured interviews along with Diana Tang. We would also  
391    like to acknowledge Kerrie Ren for her assistance with study administration, as well as  
392    Guide Dogs NSW/ACT.

## 393    **Conflict of Interest**

394    The authors declare that they have no conflict of interest.

## References

1. Bourne RRA, Flaxman SR, Braithwaite T, Cicinelli MV, Das A, Jonas JB *et al.* Magnitude, temporal trends, and projections of the global prevalence of blindness and distance and near vision impairment: a systematic review and meta-analysis. *Lancet Glob Health* 2017; **5**: e888-e897.
2. Court H, McLean G, Guthrie B, Mercer SW, Smith DJ. Visual impairment is associated with physical and mental comorbidities in older adults: a cross-sectional study. *BMC Med* 2014; **12**: 181.
3. Foreman J, Keel S, Xie J, cvan Wijngaarden P, Crowston J, Taylor HR *et al.* 2016. National Eye Health Survey. <https://www.vision2020australia.org.au/resources/national-eye-health-survey-report/>
4. Evans JR, Fletcher AE, Wormald RP. Depression and anxiety in visually impaired older people. *Ophthalmology* 2007; **114**: 283-288.
5. Nyman SR. Psychosocial issues in engaging older people with physical activity interventions for the prevention of falls. *Can J Aging* 2011; **30**: 45-55.
6. Brody BL, Gamst AC, Williams RA, Smith AR, Lau PW, Dolnak D *et al.* Depression, visual acuity, comorbidity, and disability associated with age-related macular degeneration. *Ophthalmology* 2001; **108**: 1893-1900.
7. Rovner BW, Casten RJ, Hegel MT, Massof RW, Leiby BE, Ho AC *et al.* Improving function in age-related macular degeneration: a randomized clinical trial. *Ophthalmology* 2013; **120**: 1649-1655.
8. Casten R, Rovner BW, Leiby BE, Tasman W. Depression despite anti-vascular endothelial growth factor treatment of age-related macular degeneration. *Arch Ophthalmol* 2010; **128**: 506-508.

- 419 9. van der Aa HP, van Rens GH, Comijs HC, Bosmans JE, Margrain TH, van Nispen  
420 RM. Stepped-care to prevent depression and anxiety in visually impaired older  
421 adults--design of a randomised controlled trial. *BMC Psychiatry* 2013; **13**: 209.
- 422 10. Holloway E, Sturrock B, Lamoureux E, Hegel M, Casten R, Mellor D *et al.* Delivering  
423 problem-solving treatment in low-vision rehabilitation: A pilot feasibility study. *Rehabil*  
424 *Psychol* 2018; **63**: 349-356.
- 425 11. van der Aa HP, Margrain TH, van Rens GH, Heymans MW, van Nispen RM.  
426 Psychosocial interventions to improve mental health in adults with vision impairment:  
427 systematic review and meta-analysis. *Ophthalmic Physiol Opt* 2016; **36**: 584-606.
- 428 12. Kamga H, McCusker J, Yaffe M, Sewitch M, Sussman T, Strumpf E *et al.* Self-care  
429 tools to treat depressive symptoms in patients with age-related eye disease: a  
430 randomized controlled clinical trial. *Clin Exp Ophthalmol* 2017; **45**: 371-378.
- 431 13. Sturrock B. 2018. Treating depression in people with vision impairment.  
432 [https://www.psychology.org.au/for-members/publications/inpsych/2018/feb/Treating-](https://www.psychology.org.au/for-members/publications/inpsych/2018/feb/Treating-depression-in-people-with-vision-impairm)  
433 [depression-in-people-with-vision-impairm](https://www.psychology.org.au/for-members/publications/inpsych/2018/feb/Treating-depression-in-people-with-vision-impairm)
- 434 14. Holloway E, Sturrock B, Lamoureux E, Keeffe J, Hegel M, Casten R *et al.* Can we  
435 address depression in vision rehabilitation settings? Professionals' perspectives on  
436 the barriers to integrating problem-solving treatment. *Disabil Rehabil* 2018; **40**: 287-  
437 295.
- 438 15. Michie S, van Stralen MM, West R. The behaviour change wheel: a new method for  
439 characterising and designing behaviour change interventions. *Implement Sci* 2011; **6**:  
440 42.
- 441 16. Vaismoradi M, Turunen H, Bondas T. Content analysis and thematic analysis:  
442 Implications for conducting a qualitative descriptive study. *Nurs Health Sci* 2013; **15**:  
443 398-405.

- 444 17. Srivastava P, Hopwood N. A Practical Iterative Framework for Qualitative Data  
445 Analysis. *International Journal of Qualitative Methods* 2009; **8**: 76-84.
- 446 18. Tong A, Sainsbury P, Craig J. Consolidated criteria for reporting qualitative research  
447 (COREQ): a 32-item checklist for interviews and focus groups. *Int J Qual Health Care*  
448 2007; **19**: 349-357.
- 449 19. Dillon L, Duffy P, Tiedemann A, Keay L. Acceptability of fall prevention strategies for  
450 older people with vision impairment. *International Journal of Orientation & Mobility*  
451 2018; **9**: 1-9.
- 452 20. Nyman SR, Dibb B, Victor CR, Gosney MA. Emotional well-being and adjustment to  
453 vision loss in later life: a meta-synthesis of qualitative studies. *Disabil Rehabil* 2012;  
454 **34**: 971-981.
- 455 21. Craig A, Nicholson Perry K, Guest R, Tran Y, Middleton J. Adjustment following  
456 chronic spinal cord injury: Determining factors that contribute to social participation.  
457 *Br J Health Psychol* 2015; **20**: 807-823.
- 458 22. Nollett CL, Bray N, Bunce C, Casten RJ, Edwards RT, Hegel MT *et al.* High  
459 Prevalence of Untreated Depression in Patients Accessing Low-Vision Services.  
460 *Ophthalmology* 2016; **123**: 440-441.
- 461 23. Boeschoten RE, Dekker J, Uitdehaag BM, Polman CH, Collette EH, Cuijpers P *et al.*  
462 Internet-based self-help treatment for depression in multiple sclerosis: study protocol  
463 of a randomized controlled trial. *BMC Psychiatry* 2012; **12**: 137.
- 464 24. Reinhardt JP, Horowitz A, Cimarolli VR, Eimicke JP, Teresi JA. Addressing  
465 depression in a long-term care setting: a phase II pilot of problem-solving treatment.  
466 *Clin Ther* 2014; **36**: 1531-1537.
- 467 25. Haaga DA. Introduction to the special section on stepped care models in  
468 psychotherapy. *J Consult Clin Psychol* 2000; **68**: 547-548.

- 469 26. Patel V, Weiss HA, Chowdhary N, Naik S, Pednekar S, Chatterjee S *et al.*  
470 Effectiveness of an intervention led by lay health counsellors for depressive and  
471 anxiety disorders in primary care in Goa, India (MANAS): a cluster randomised  
472 controlled trial. *Lancet* 2010; **376**: 2086-2095.
- 473 27. van't Veer-Tazelaar PJ, van Marwijk HW, van Oppen P, van Hout HP, van der Horst  
474 HE, Cuijpers P *et al.* Stepped-care prevention of anxiety and depression in late life: a  
475 randomized controlled trial. *Arch Gen Psychiatry* 2009; **66**: 297-304.
- 476 28. National Institute for Health and Care Excellent (NICE). 2001. Common mental  
477 health problems: identification and pathways to care. Clinical guideline [CG123].  
478 <https://www.nice.org.uk/guidance/CG123/chapter/1-Guidance#stepped-care>

479 **Titles and legends to figures**

480 Figure 1 Barriers and facilitators to mental wellbeing program participation mapped on the  
481 behaviour change wheel[15]. Barriers to participation appear in bold, while facilitators  
482 appear underlined, with their relative intervention function presented in the outer wheel.