


Central Lancashire Online Knowledge (CLOK)

Title	Stroke in Sierra Leone: Case fatality rate and functional outcome after stroke in Freetown
Type	Article
URL	https://clock.uclan.ac.uk/46013/
DOI	https://doi.org/10.1177/17474930231164892
Date	2023
Citation	Youkee, Daniel, Deen, Gibrilla, Baldeh, Mamadu, Conteh, Zainab Fatmata, Fox-Rushby, Julia, Gbessay, Musa, Johnson, Jotham, Langhorne, Peter, Leather, Andrew et al (2023) Stroke in Sierra Leone: Case fatality rate and functional outcome after stroke in Freetown. International Journal of Stroke. p. 174749302311648. ISSN 1747-4930
Creators	Youkee, Daniel, Deen, Gibrilla, Baldeh, Mamadu, Conteh, Zainab Fatmata, Fox-Rushby, Julia, Gbessay, Musa, Johnson, Jotham, Langhorne, Peter, Leather, Andrew, Lisk, Durodamil Radcliffe, Marshall, Iain, O'Hara, Jessica, Pessima, Sahr, Rudd, Anthony, Soley-Bori, Marina, Thompson, Melvina, Wafa, Hatem, Wang, Yanzhong, Watkins, Caroline Leigh, Williams, Christine, Wolfe, Charles and Sackley, Catherine

It is advisable to refer to the publisher's version if you intend to cite from the work.
<https://doi.org/10.1177/17474930231164892>

For information about Research at UCLan please go to <http://www.uclan.ac.uk/research/>

All outputs in CLOK are protected by Intellectual Property Rights law, including Copyright law. Copyright, IPR and Moral Rights for the works on this site are retained by the individual authors and/or other copyright owners. Terms and conditions for use of this material are defined in the <http://clock.uclan.ac.uk/policies/>



Stroke in Sierra Leone: Case fatality rate and functional outcome after stroke in Freetown

Journal:	International Journal of Stroke
Manuscript ID	IJS-12-22-10562.R1
Manuscript Type:	Research
Date Submitted by the Author:	30-Jan-2023
Complete List of Authors:	<p>Youkee, Daniel; King's College London School of Population Health and Environmental Sciences</p> <p>Deen, Gibrilla; University of Sierra Leone College of Medicine and Allied Health Sciences; Connaught Teaching hospital</p> <p>Paldeh, Mamadu; University of Sierra Leone College of Medicine and Allied Health Sciences</p> <p>Connaught, Zainab; Connaught Teaching Hospital</p> <p>Fox-Rushby, Julia; King's College London School of Population Health and Environmental Sciences; NIHR Applied Research Centre, South London</p> <p>Gbessy, Musa; 34 Military Hospital</p> <p>Johnson, Jotham; Stroke Association Sierra Leone</p> <p>Langhorne, Peter; University of Glasgow, Institute of Cardiovascular and Medical Sciences</p> <p>Leather, Andrew; King's Centre for Global Health and Health Partnerships, School of Population Health and Environmental Sciences, King's College London</p> <p>Lisk, Durodamil; University of Sierra Leone College of Medicine and Allied Health Sciences</p> <p>Marshall, Iain ; King's College London School of Population Health and Environmental Sciences</p> <p>O'Hara, Jessica; King's College London School of Population Health and Environmental Sciences</p> <p>Pessima, Sahr; University of Sierra Leone College of Medicine and Allied Health Sciences</p> <p>Rudd, Anthony; King's College London School of Population Health and Environmental Sciences, Health & Social Care Research</p> <p>Soley-Bori, Marina; King's College London School of Population Health and Environmental Sciences, School of public health</p> <p>Thompson, Melvina; Connaught Teaching Hospital</p> <p>Wafa, Hatem; King's College London School of Population Health and Environmental Sciences</p> <p>Wang, Yanzhong; King's College London, Population Health & Environmental Sciences; NIHR Applied Research Centre, South London</p> <p>Watkins, Caroline; University of Central Lancashire College of Health and Wellbeing</p> <p>Williams, Christine; University of Sierra Leone College of Medicine and Allied Health Sciences</p> <p>Wolfe, Charles; King's College London School of Population Health and Environmental Sciences; NIHR applied research collaboration, South London</p> <p>Sackley, Catherine; King's College London School of Population Health and Environmental Sciences; University of Nottingham Faculty of Medicine and Health Sciences</p>

Keywords:	Stroke, case fatality rate, mortality, sierra leone, africa, barthel index, function

SCHOLARONE™
Manuscripts

Author Accepted Manuscript

Stroke in Sierra Leone: Case fatality rate and functional outcome after stroke in Freetown

Authors:

Daniel Youkee^{1*}, Gibrilla F Deen^{2,3}, Mamadu Baldeh², Zainab F Conteh³, Julia Fox-Rushby^{1,4}, Musa Gbessay⁵, Jotham Johnson⁶, Peter Langhorne⁷, Andrew JM Leather⁸, Durodami R Lisk², Iain J Marshall^{1,4}, Jessica O'Hara¹, Sahr Pessima², Anthony Rudd¹, Marina Soley-Bori¹, Melvina Thompson^{2,3}, Hatem Wafa¹, Yanzhong Wang^{1,4}, Caroline L Watkins⁹, Christine E Williams², Charles DA Wolfe^{1,4}, Catherine M Sackley^{1,10}

*Corresponding author

Twitter: @mirrorsandmaps

Affiliations:

1. School of Life Course and Population Sciences, King's College London, London, UK
2. College of Medicine and Allied health Sciences, University of Sierra Leone, Freetown, Sierra Leone
3. Connaught Teaching Hospital, University of Sierra Leone teaching Hospital Complex, Freetown, Sierra Leone
4. National Institute for Health Research Collaboration for Leadership in Applied Health Research and Care South London, London, UK.
5. 34 Military Hospital, Freetown, Sierra Leone
6. Stroke Association Sierra Leone
7. Academic Section of Geriatric Medicine, School of Cardiovascular and Metabolic Health, University of Glasgow.
8. King's Centre for Global Health and Health Partnerships, School of Life Course and Population Sciences, King's College London, London, UK
9. Faculty of Health and Care, University of Central Lancashire, Preston, United Kingdom
10. Faculty of Medicine and Health Sciences, University of Nottingham, UK

Key words: Stroke, Sierra Leone, Africa, case fatality rate, function, outcomes, Barthel index

Word Count: 4891/5000

Tables and figures

1. *Table one: Case fatality rate by stroke type, at 7 days, 30 days, 90 days, one year and two years post stroke*
2. *Table two: Cox Proportional Hazards model for fatality for all strokes*
3. *Table three: Logistic regression for functional independence with assistance (BI \geq 85) at one year*
4. *Figure 1: Kaplan-Meier estimates for stroke survival from date of stroke onset.*
5. *Figure 2: Sankey plot of flow of functional outcome pre-stroke, seven days, 90 days and one year post stroke*

Abstract

Background: There is limited information on long term outcomes after stroke in Sub-Saharan Africa (SSA). Current estimates of case fatality rate (CFR) in SSA are based on small sample sizes with varying study design and report heterogenous results.

Aims: We report CFR and functional outcomes from a large, prospective, longitudinal cohort of stroke patients in Sierra Leone and describe factors associated with mortality and functional outcome.

Methods: A prospective longitudinal stroke register was established at both adult tertiary government hospitals in Freetown, Sierra Leone. It recruited all patients ≥ 18 years with stroke, using the World Health Organization definition, from May 2019 until October 2021. To reduce selection bias onto the register all investigations were paid by the funder and outreach conducted to raise awareness of the study. Sociodemographic data, National Institute of Health Stroke Scale (NIHSS) and Barthel Index (BI) was collected on all patients on admission, at seven days, 90 days, one year and two years post stroke. Cox proportional-hazards models were constructed to identify factors associated with all-cause mortality. A binomial logistic regression model reports odds ratio (OR) for functional independence at one year.

Results: 986 patients with stroke were included, of which 857 (87%) received neuroimaging. Follow up rate was 82% at one year, missing item data was $<1\%$ for most variables. Stroke cases were equally split by sex and mean age was 58.9 (SD: 14.0) years. 625 (63%) were ischaemic, 206 (21%) primary intracerebral haemorrhage, 25 (3%) subarachnoid haemorrhage and 130 (13%) were of undetermined stroke type. Median NIHSS was 16 (9-24). CFR at 30 days, 90 days, 1 year and 2 years was 37%, 41%, 49% and 53% respectively. Factors associated with increased fatality at any timepoint were male sex HR:1.28 (1.05-1.56), previous stroke HR:1.34 (1.04-1.71), atrial fibrillation HR:1.58 (1.06-2.34), subarachnoid haemorrhage HR:2.31 (1.40-3.81), undetermined stroke type HR: 3.18(2.44-4.14) and in-hospital complications HR: 1.65 (1.36-1.98). 93% of patients were completely independent prior to their stroke, declining to 19% at one year after stroke. Functional improvement was most likely to occur between 7 and 90-days post stroke with 55% patients improving, and 13% improving between 90 days to one year. Increasing age OR: 0.97 (0.95-0.99), previous stroke OR: 0.50 (0.26-0.98), NIHSS OR 0.89 (0.86-0.91), undetermined stroke type OR:0.18 (0.05-0.62) and ≥ 1 in hospital complication OR:0.52 (0.34-0.80) were associated with lower OR of functional independence at one year. Hypertension OR:1.95 (1.14-3.44) and being the primary breadwinner of the household OR:1.59 (1.01-2.49) were associated with functional independence at one year.

Conclusion: Stroke affected younger people, and resulted in high rates of fatality and functional impairment relative to global averages. Key clinical priorities for reducing fatality include preventing stroke-related complications through evidence-based stroke care; improved detection and management of atrial fibrillation, and increasing coverage of secondary prevention. Further research into care pathways and interventions to encourage

care seeking for less severe strokes should be prioritized, including reducing the cost barrier for stroke investigations and care.

Data availability: Requests for access to anonymized data should be made to the SISLE team <https://www.kcl.ac.uk/research/stroke>.

500-500 words

Introduction

It is estimated that stroke is the second leading cause of adult death in Sub Saharan Africa (SSA), and a cause of significant morbidity¹. However, prospective stroke studies of case fatality rate and functional outcome in SSA are limited in number and quality², and long-term outcomes after stroke have not been previously studied in Sierra Leone.

Understanding survival and functional outcome after stroke is important to provide prognostic information for the patient, characterise the natural history of stroke and for health system planning to meet the acute and long-term care needs of patients after stroke³.

The evidence base for CFR after stroke in SSA is heterogenous and of varying quality⁴.

Available data suggest that mortality is higher in some African countries than global estimates⁵. A systematic review of CFR in SSA, published in 2021, included 91 studies⁶ with 9 (9.9%) studies rated as high-quality evidence. CFR at 90 days was 22.3% [95%CI: 16.6–29.2] and at one year was 33.2% (95% CI: 23.6–44.5). For one year CFR, the review included a pooled sample size of 4809 patients, with CFR displaying high heterogeneity ranging from 9.5% to 75.4% across studies. In contrast, a recent prospective hospital study of 564 stroke patients in Ghana, found CFR at 3 months and one year of 43.2% and 49.7% respectively⁷. This compares to global 30-day CFR of 17–30% reported in high income countries and 18–35% in low and middle income countries⁸. Worldwide, CFR differs by stroke type, generally being higher for intracerebral haemorrhage compared to ischaemic stroke at one year^{3,9}. Evidence suggests that prevalence of stroke type and stroke subtypes prevalence differ in SSA, with higher proportions of intracerebral haemorrhage, higher proportions of small vessel disease and lower proportions of cardioembolic ischaemic stroke subtypes, compared to other regions^{10–12}. The different prevalence of stroke types and subtypes may impact case fatality, indeed pooled estimates of CFR at one year in SSA are higher for ischaemic stroke compared to intracerebral haemorrhage⁶. Older age is a nonmodifiable risk factor for stroke strongly associated with both increased case fatality and worse functional outcome after stroke in other regions¹³. However, stroke occurs at a younger age in SSA¹⁴ compared to other regions and life expectancy at birth also differs⁵. Previous work in Sierra Leone suggests stroke occurs at median age of 59 years¹⁵, in the context of an adult life expectancy at birth of 60 years, 20 years younger than other regions¹⁶.

Due to differences in demographics, prevalence of stroke types and access to high quality stroke care, we hypothesise that CFR and variables associated with CFR may differ in Sierra Leone compared to global rates and associations. A previous retrospective hospital based study in Sierra Leone found in-hospital mortality to be associated with prior stroke, hypertension, haemorrhagic stroke and aspiration pneumonia¹⁷. Whilst a prospective study at the same hospital found male sex, pneumonia, subarachnoid haemorrhage and undetermined stroke types were associated with in-hospital death¹⁵. In this paper we

describe long-term case fatality and function after stroke in Sierra Leone and identify factors associated with survival and functional outcome.

Methods

A prospective stroke register was established at the two adult tertiary government hospitals in Freetown, Sierra Leone at Connaught Teaching Hospital from 1st May 2019 until 30th September 2021 and at 34th Military Hospital from 1st February 2021 until 2nd September 2021. All patients 18 years and over meeting the WHO ICD10 definition of stroke were included. The study methods and the health care setting have been previously described¹⁵. All stroke subtypes were included: ischaemic (ICD63); intracerebral haemorrhage (ICD61); sub-arachnoid haemorrhages (ICD60) and undetermined stroke types (ICD62)¹⁸. Classification of pathological stroke subtype, using the Oxford Community Stroke Project (OCSF) classification¹⁹ was conducted by an experienced stroke physician, with reference to the case history, investigation results, and imaging. During the study period, stroke investigations, including CT-scanning, were funded by the grant, National Institute for Health Research (NIHR) (GHR:17:53:66), and provided free-of-charge to patients, to reduce bias in access to investigations. During the study period there was no functional CT scanner at either of the hospitals, so CT scanning was provided at two off-site private radiology centres. The study supported ambulance transfer accompanied by a clinician to enable safe access to imaging. Participants who did not receive neuroimaging, (often in practice due to being too critically ill to transfer for scanning), or those in whom neuroimaging was not conclusive were classified as undetermined stroke type.

Participants were followed up at 90 days, one year and two years post stroke. Patients were primarily contacted by telephone, and those uncontactable were visited at home. We report on follow up from 1st May 2019 to 22nd July 2022. All-cause mortality was recorded from hospital records and as reported by caregiver or relative at follow up. Functional outcome was measured using the Barthel Index (BI) and was retrospectively reported by patients and family seven days prior to stroke, then measured at seven days post stroke, 90 days post stroke, at one year, and each year after stroke. BI was categorised as, completely dependent (BI<60), dependent (BI= 60-84), independent with assistance BI ≥85²⁰ and independent without assistance as BI=100.

Survival curves were constructed for the whole population, and for subgroups by stroke subtype, age, and sex; all using the Kaplan-Meier method. Multivariable Cox proportional-hazards models were conducted to assess the independent effect of variables on all-cause mortality. A logistic regression model, reporting Odds ratios (OR) was created for functional independence at one year. A full description of stroke type classification, risk factors, missing data and regression model development is provided in the appendix.

All data was collected on standardised paper Case Report Forms. Double data entry was conducted, and all data uploaded onto REDCap^{TM21}. Statistical analyses were performed in STATA v17, StataCorp^{TM22}. The study received ethical approval from King's College London (HR-18/19-8467) and approval from the Sierra Leone Ethical and Scientific Review

Committee on 18th December 2018. Written consent was sought from all patients. For those judged not to have capacity, informed consent was sought from the next of kin.

Results

The register recruited 1145 people with suspected strokes. After clinical review and neuroimaging, 986 were confirmed as strokes and were maintained as the core population for analysis, 915 at Connaught Teaching Hospital and 71 at 34th Military Hospital. Stroke cases were equally split by sex and mean age was 58.9 (SD: 14.0) years. During the study period there was no stroke unit at either hospital and no patients received thrombolytic therapy or mechanical thrombectomy. The median time from stroke onset to admission was 24 hours and median length of stay was 7 days (IQR 3-12). 857 (87%) patients underwent neuroimaging, 847 received CT scans and 10 received MRI, comparative statistics of patients who received neuroimaging vs those who did not are presented in the supplementary material. Median NIHSS was 16 (9-24). 625 (63%) patient had ischaemic stroke, 206 (21%) primary intracerebral haemorrhage, 25 (3%) subarachnoid haemorrhage and 130 (13%) were of undetermined stroke type. Ischaemic stroke subtypes by OSCP classification are described in *Supplementary Figure one*. 355 (36.0%) patients died in hospital during the initial admission and 175 (17.7%) post hospital discharge. At one year 182 (18.5%) participants were lost to follow up, follow up counts are reported in *Supplementary table one*. Missing item data is reported in *Supplementary table two*, and was low at under 1% for most variables.

Case Fatality Rate

Case fatality rate was 37.1% at 30 days, 44.4% at 90 days, 49.9% at one year and 53.2% at two years (*table one*). Case fatality rates by stroke type are shown in *Table one* and univariable analysis of CFR at one year is reported in *Supplementary table three*. Ischaemic stroke CFR increased from 25.3% at 30 days to 45.6% at two years, whilst intracerebral haemorrhage CFR increased from 40.3% at 30 days to 51.0% at two years.

Case Fatality	All strokes	Ischaemic	Intracerebral haemorrhage	Subarachnoid haemorrhage	Undetermined
	N=986	N=625 (63%)	N=206 (21%)	N=25 (3%)	N=130 (13%)
30 day case fatality	366 (37.1%)	158 (25.3%)	83 (40.3%)	15 (60%)	110 (84.6%)
90 day case fatality	438 (44.4%)	210 (33.6%)	91 (44.2%)	18 (72%)	19 (31.5%)
One year case fatality	492 (49.9%)	258 (41.3%)	95 (46.1%)	18 (72%)	121 (93.1%)
Two year case fatality	529 (53.2%)	285 (45.6%)	105 (51.0%)	18 (72%)	121 (93.1%)

Table one: Case fatality rate by stroke type, at 7 days, 30 days, 90 days, one year and two years post stroke

Kaplan-Meier survival estimates are shown in *figure 1*. The survival estimates demonstrate that most deaths occur within the first 90 days post stroke, however survival continues to

decrease up to two years post stroke. *Figure 1c* demonstrates significant differences in survival by stroke type ($p=0.0001$), with intracerebral haemorrhage patients initially having a steep decrease in survival compared to ischaemic stroke patients, followed by survival estimates for the two stroke types becoming closer over time. *Figure 1d* demonstrates significantly reduced survival in patients ≥ 55 years ($p=0.0001$). Kaplan-Meier survival curves with censoring hashmarks are reported in *Supplementary figure two*. Univariable analysis of death at one year by stroke type is reported in *Supplementary table four*.

Figure 1: Kaplan-Meier estimates for stroke survival from date of stroke onset. 1a: Kaplan-Meier survival estimate for all strokes (n=986). 1b Kaplan-Meier survival estimate by sex (n=986) logrank test $p=0.75$. 1c Kaplan-Meier survival estimate by stroke type (n=986), logrank test $p=0.0001$. 1d Kaplan-Meier survival estimate by age <55 years vs ≥ 55 years (n=981), logrank test $p=0.0001$.

Cox proportional hazards model for fatality is shown in *table two below*. Male sex, previous stroke, atrial fibrillation, subarachnoid haemorrhage, undetermined stroke type, and hospital complication were all associated with death. The presence of hypertension was associated with survival. A sensitivity analysis excluding in hospital complications is presented in *Supplementary table six*.

Independent Variable	Count	Hazard Ratio	95% CI
Age (each additional year)*	58.9 (SD: 14.0)	1.00	1.00-1.01
Male sex	495 (50.2%)	1.28	1.05-1.56
Previous stroke	128 (13.0%)	1.34	1.04-1.71
Hypertension	831 (84.3%)	0.71	0.57-0.90
Diabetes	212 (21.5%)	1.04	0.84-1.29
Atrial fibrillation	38 (3.9%)	1.58	1.06-2.34
Dyslipidaemia	401 (40.7%)	-	
Current smoker	153 (15.8%)	-	
Alcohol use (any)	255 (26.7%)	-	
Resident of Freetown	822 (84.0%)	-	
Higher education level	367 (37.2%)	0.91	0.75-1.10
Primary breadwinner	424 (43%)	0.88	0.72-1.07
Pre-stroke Barthel Index Mean (SD)	96.7 (12.4)	-	
NIHSS (each additional point)	16 (IQR: 9-24)	1.07	1.06-1.08
¥Intracerebral haemorrhage	206 (20.9%)	1.18	0.93-1.50
¥Subarachnoid haemorrhage	15 (2.5%)	2.31	1.40-3.81
¥Undetermined stroke type	130 (13.2%)	3.18	2.44-4.14
≥1 in hospital complication	396 (40.2%)	1.65	1.36-1.98

Table two: Cox Proportional Hazards model for fatality for all strokes. *Control variables.

¥Stroke type compared to ischaemic stroke (n=186). Variables with dashes (-) were not included in regression model.

Functional Outcome

Progression of functional status displayed by Barthel Index (BI) at seven days prior to stroke, seven days post stroke, 90 days, and one year post stroke is shown in figure 2. Seven days prior to stroke 93% of patients were independent with assistance 5% at seven days post stroke, 28% at 90 days, and 19% at one year. From seven days to 90 days, categorical functional status improved for 272 (34.8%) patients, worsened (including death) for 251 (32.1%), 145 (18.5%) stayed the same and 114 (14.6%) were missing. From 90 days to one year, categorical functional status improved for 56 (13.0%), worsened (including death) for 92 (21.4%), 169 (39.3%) stayed the same and 113 (26.3%) were missing.




Figure 2: Sankey plot of flow of functional outcome pre-stroke, seven days, 90 days and one year post stroke. Percentage demonstrates proportion of patients with functional outcome measured by BI: Independent with assistance BI ≥ 85 , Dependent BI 60-84, Completely dependent BI < 60 .

Logistic regression with Odds ratios (OR) for functional independence with assistance (BI ≥ 85) at one year is shown in *table three*. Increasing age, previous stroke, NIHSS, undetermined stroke type and ≥ 1 in hospital complication were associated with lower OR of functional independence at one year. Whilst hypertension and being the primary breadwinner of the household were associated with higher OR of functional independence at one year.

Independent Variable	OR	95% CI
Age (each additional year)	0.97	0.95-0.99
Male sex	0.89	0.57-1.39
Previous stroke	0.50	0.26-0.98
Hypertension	1.98	1.14-3.44
Diabetes	0.64	0.39-1.05
Atrial fibrillation	0.73	0.20-2.66
Higher education level	1.32	0.86-2.01
Primary breadwinner	1.59	1.01-2.49
NIHSS (each additional point)	0.89	0.86-0.91
¥Intracerebral haemorrhage	1.26	0.77-2.07

¥Subarachnoid haemorrhage	0.36	0.11-1.23
¥Undetermined stroke type	0.18	0.05-0.62
≥1 in hospital complication	0.52	0.34-0.80

Table three: Logistic regression for functional independence with assistance (BI≥85) at one year, n=. ¥Stroke type compared to ischaemic stroke.

Discussion

This is the first publication reporting long term outcomes after stroke in Sierra Leone. We report 30-day CFR of 37.1%, at the higher end of the global estimates of CFR²³. 90 day CFR of 44.4% and one year CFR of 49.7% are higher than pooled CFR estimates in SSA at 90 days of 22.3% (95%CI: 16.6-29.2) and one year 33.2%(95%CI: 23.6-44.5) respectively⁶. Our reported CFR is higher than similar studies in Benin²⁴, Kenya²⁵, and closely matches reported CFR in Ghana⁷. The high CFR reported reflects the severe case mix, our median NIHSS was 16 (9-24), higher than other West African hospital-based stroke registers and significantly higher than seen in UK or North American hospital studies²⁶.

Case fatality in our cohort was higher in patients with intracerebral haemorrhage strokes than ischaemic strokes at all time points, the majority of deaths for patients with intracerebral haemorrhage occurred early in hospital, whilst ischaemic stroke mortality was more evenly distributed across the study period. Relative to global pooled estimates by stroke type, ischaemic stroke CFR was higher than expected and intracerebral haemorrhage CFR was within expected ranges. Our ischaemic stroke 30-day CFR of 25% was higher than worldwide pooled estimates of 13.5% (95% CI 12.3% to 14.7%) at 30 days²⁷. ICH mortality was similar to global estimates, 40% at 30-days compared to 31.3% (95%CI 31.5-41.2) and 46% at one year compared to 50.7% (95%CI = 45.2-56.2)²⁸. This may partially reflect the under-detection of less severe ischaemic strokes in our cohort, which may happen if people with less severe strokes choose not to seek care, due to common barriers in Sierra Leone such as cost of care²⁹, distance³⁰, trust in the formal health system³¹ and health literacy³². Patients with intracerebral haemorrhage in our cohort were younger (*Supplementary table four*), had lower prevalence of diabetes, dyslipidaemia, atrial fibrillation, and higher pre-morbid status (pre-stroke BI) compared to ischaemic strokes. Concurrently, it may reflect a lack hyperacute stroke care for ischaemic strokes and lack of access to quality care for the comorbidities more prevalent in ischaemic stroke patients. Undetermined stroke types were associated with increased CFR as these patients were too sick to transfer for neuroimaging or died before neuroimaging could confirm stroke type (*Supplementary table five*). Subarachnoid haemorrhage CFR of 60% corresponds poorly to global estimates of CFR ranging from 27%-44%³³, reported CFR of 40% in Sudan³⁴, 44.4% in Nigeria³⁵, 45.6% in Kenya³⁶ and reflects both the severity of strokes included in our cohort and lack of timely access to care, including neurosurgical intervention available in our setting^{37, 38}.

We report strokes occurring in younger people, with a mean age of 59 years. As in other settings, age was significantly associated with increased mortality and patients alive at one

year were on average five years younger than those dead at one year. Patients with a previous stroke had increased mortality HR: 1.34 (1.04-1.71), similar to other findings from SSA⁶ suggesting a need to improve the coverage and effectiveness of secondary prevention interventions in Sierra Leone. Atrial fibrillation was associated with increased mortality HR: 1.58 (1.06-2.34). However, only a single 12-lead ECG was conducted in our cohort, therefore atrial fibrillation was likely under-detected. Atrial fibrillation detection should be increased through use of cardiac holters or wearable smart devices with proven diagnostic accuracy³⁹. Atrial fibrillation diagnosis and management remains challenging in our setting, with the prohibitive costs of direct oral anticoagulants⁴⁰, and low levels of access to affordable and reliable international Normalized Ratio (INR) monitoring to allow safe and effective warfarin prescription^{41, 42}. In hospital complications were associated with fatality and worse functional outcome, demonstrating the need to implement evidence based stroke care, such as stroke unit based care, inclusive of swallow screening for the prevention of aspiration pneumonia¹⁵.

Our previous study found male sex to be associated with in hospital fatality¹⁵, we demonstrate this finding again for long term fatality HR 1.28 (1.05-1.56). Further research is needed to understand whether this is due to intrinsic sex survival differences or potentially related to quality of care differences in the gender separated hospital wards. The presence of hypertension was associated with increased survival HR:0.71 (0.57-0.90) in our cohort, pooled findings from SSA found no association between hypertension and CFR⁶, whilst other West African prospective registers found a similar but non-significant direction of effect. This may demonstrate the higher relative fatality of stroke caused by other aetiologies, including renal disease, and malignancy, which were too small a sample size to include in our regression model. Alternatively, it may be due to the influence of hypertension on prevalence of subtypes of haemorrhagic stroke which may influence survival^{43, 44}, which were not included in our regression model. Hypertension remains the primary dominant modifiable risk factor for stroke in our region¹¹, and hypertension detection, management and control should be an urgent priority⁴⁵.

Functional impairment was considerable, 93% of patients were completely independent seven days prior to their stroke, at one year post stroke only 19% were independent with assistance. Most functional recovery was seen between seven and 90 days with 34.8% of patients reporting functional improvement, and a smaller proportion 13.0% improving from 90 days to one year. The functional improvement in some, was matched by a greater amount of decrease in functional outcome and death in others, 32.1% worsening from seven days to 90 days, and 21.4% from 90 days to one year. Rates of functional improvement appear similar to other studies in West Africa^{46, 47}. Functional independence at one year was more likely in younger patients⁴⁸, first-in-a-lifetime strokes and being the primary breadwinner for the household. Socioeconomic status proxies, being the primary breadwinner (significantly) and higher educational attainment (non-significant), were associated with both improved survival and functional outcome at one year. This corresponds to regional⁴⁹ and international findings⁵⁰, and should inform the development of equitable stroke services in Sierra Leone.

Strengths

This is the first study to report long term CFR and functional status after stroke in Sierra Leone. The study benefits from a prospective, multi-centre design and is one of the largest longitudinal studies of stroke patients in SSA to be published. Key elements of the stroke register were prospective methodology, community awareness raising and removal of the cost barrier for stroke investigations to reduce selection bias and increase access to care. During the study period we recorded 381 strokes per year at Connaught Hospital, compared to 179 strokes per year in 2018¹⁷. This is likely to include not only increased presentation of patients to the hospital but also increased awareness and recognition of stroke amongst healthcare workers and the removal of the cost barrier for investigations. Our study benefited from neuroimaging rates of 87% and a follow up rate of 82% at one year.

Limitations

The study is not population based, therefore results cannot be extrapolated to the population level and are influenced by access to care. Selection bias onto the register, with care seeking only for severe strokes and under-detection of less severe strokes likely contributes to the high CFR reported. Due to the lack of an onsite CT scanner, we were unable to provide imaging to the most critically ill patients with the most severe strokes (*supplementary table five*), we therefore do not know the stroke subtype of the sickest patients in our cohort. The study is limited by not using an intracerebral haemorrhage classification system⁵¹, such as the structural vascular lesions (S), medication (M), amyloid angiopathy (A), systemic disease (S), hypertension (H), or undetermined (U) (SMASH-U) criteria⁴³. Atrial fibrillation was only assessed by a single 12-lead ECG, therefore was likely under-detected.

Conclusions

We demonstrate high CFR of 49.7% at one year relative to estimates in SSA and Europe. We report significant functional impairment, 93% of patients were completely independent seven days prior to their stroke, by one year post stroke only 19% were independent with assistance. Key priorities include prevention of stroke related complications through evidence based interventions, such as stroke unit based care, increased detection and improved management of atrial fibrillation, and enhanced coverage of secondary prevention, are key to reducing CFR after stroke in Sierra Leone. Further research into care pathways and interventions to encourage care seeking for less severe strokes should be prioritized.

Funding

This research was funded by the National Institute for Health Research (NIHR) (GHR:17:63:66) using UK aid from the UK Government to support global health research; and is supported by the NIHR Programme Grants for Applied Research scheme (NIHR202339). The views expressed are those of the author(s) and not necessarily those of the NIHR or the Department of Health and Social Care. DY is funded by a Medical Research Council Clinical Research Training Fellowship: MR/W000903/1. CW, JFR and YW are

supported by the National Institute for Health Research (NIHR) Biomedical Research Centre (BRC), Guy's and St Thomas' NHS Foundation Trust and King's College London, London, United Kingdom and the NIHR Applied Research Collaboration (ARC) South London, London, United Kingdom.

Data Availability

The raw data for this study contain both personally identifiable and confidential clinical data. Requests for access to anonymised subsets of data for academic use should be made to the SISLE team where data will be made available subject to academic review and a certificate of a data-sharing agreement: <https://www.kcl.ac.uk/research/stroke>.

Acknowledgements

We acknowledge the Stroke Association Sierra Leone for their work improving lives of stroke survivors in Sierra Leone. We thank all SISLE team members and advisors, Binta Bah, Edward Barrett, Cecily Borgstein, Laura Hucks, Israel Johnson, Ishmael Kebbie, Elizabeth Koroma, Lynn Legg, Jane Monden, Ben Norrving, Albert Sama, Christella Scott, Jurate Wall and Marion Walker.

Authorship Contributions

CS, CDW, AJML, DY, GFD, JFR, AR, PL, CLW acquired the funding. CS, CDW, AJML, DY, GFD, JFR, AR, PL, CLW, IJM conceptualised the research. CS was the Chief Investigator on the grant and GFD and DRL were the Sierra Leone investigators. Investigation was conducted by CS, DY, GFD, ZFC, MB, MG, SP, MT and CW. Data curated by DY, ZFC, SP. DY conducted the analysis and wrote the first draft under the supervision and detailed commenting over time by IJM, JFR and MSB. IJM, JFR, MSB are supervisors of DY's PhD fellowship. All authors approved the final manuscript.

Conflicts of Interest

The authors declare that there is no conflicting or competing interest.

References

1. Vos T, Lim SS, Abbafati C, et al. Global burden of 369 diseases and injuries in 204 countries and territories, 1990–2019: a systematic analysis for the Global Burden of Disease Study 2019. *The Lancet* 2020; 396: 1204-1222. DOI: 10.1016/s0140-6736(20)30925-9.
2. Owolabi MO, Akarolo-Anthony S, Akinyemi R, et al. The burden of stroke in Africa: a glance at the present and a glimpse into the future. *Cardiovasc J Afr* 2015; 26: S27-38. 2015/05/13. DOI: 10.5830/CVJA-2015-038.
3. Bhalla A, Wang Y, Rudd A, et al. Differences in outcome and predictors between ischemic and intracerebral hemorrhage: the South London Stroke Register. *Stroke* 2013; 44: 2174-2181. 2013/07/03. DOI: 10.1161/strokeaha.113.001263.
4. Owolabi M, Olowoyo P, Popoola F, et al. The epidemiology of stroke in Africa: A systematic review of existing methods and new approaches. *J Clin Hypertens (Greenwich)* 2018; 20: 47-55. 2017/12/12. DOI: 10.1111/jch.13152.
5. Akinyemi RO, Ovbiagele B, Adeniji OA, et al. Stroke in Africa: profile, progress, prospects and priorities. *Nature Reviews Neurology* 2021; 17: 634-656. DOI: 10.1038/s41582-021-00542-4.

6. Thierry Adoukonou OnK, Pervenche Fotso Mefo, Mendinatou Agbé tou, Julien Magne, Glwadys Gbaguidi, Dismand Houinato, Pierre-Marie Preux, Philippe Lacroix. Stroke case fatality in sub-Saharan Africa: Systematic review and meta-analysis. *International Journal of Stroke* 2021.
7. Sarfo FS, Akassi J, Ofori E, et al. Long-term determinants of death after stroke in Ghana: Analysis by stroke types & subtypes. *J Stroke Cerebrovasc Dis* 2022; 31: 106639. 20220801. DOI: 10.1016/j.jstrokecerebrovasdis.2022.106639.
8. Feigin VL, Lawes CMM, Bennett DA, et al. Worldwide stroke incidence and early case fatality reported in 56 population-based studies: a systematic review. *The Lancet Neurology* 2009; 8: 355-369. DOI: 10.1016/s1474-4422(09)70025-0.
9. Sennfalt S, Norrving B, Petersson J, et al. Long-Term Survival and Function After Stroke. *Stroke* 2018; STROKEAHA118022913. 2018/12/26. DOI: 10.1161/STROKEAHA.118.022913.
10. Akinyemi RO, Ovbiagele B, Adeniji OA, et al. Stroke in Africa: profile, progress, prospects and priorities. *Nat Rev Neurol* 2021; 17: 634-656. 2021/09/17. DOI: 10.1038/s41582-021-00542-4.
11. Owolabi MO, Sarfo F, Akinyemi R, et al. Dominant modifiable risk factors for stroke in Ghana and Nigeria (GRIEN): a case-control study. *The Lancet Global Health* 2018; 6: e436-e446. DOI: 10.1016/s2214-109x(18)30002-0.
12. Akinyemi RO, Owolabi MO, Ihara M, et al. Stroke, cerebrovascular diseases and vascular cognitive impairment in Africa. *Brain Res Bull* 2019; 145: 97-108. 2018/05/29. DOI: 10.1016/j.brainresbull.2018.05.018.
13. Arnold M, Halpern M, Meier N, et al. Age-dependent differences in demographics, risk factors, co-morbidity, etiology, management, and clinical outcome of acute ischemic stroke. *J Neurol* 2008; 255: 1503-1507. 20080728. DOI: 10.1007/s00415-008-0949-9.
14. Owolabi M, Sarfo F, Howard VJ, et al. Stroke in Indigenous Africans, African Americans, and European Americans: Interplay of Racial and Geographic Factors. *Stroke* 2017; 48: 1169-1175. 2017/04/09. DOI: 10.1161/STROKEAHA.116.015937.
15. Youkee D, Deen G, Barrett E, et al. Prospective Stroke Register in Sierra Leone: Demographics, Stroke Type, Stroke Care and Hospital Outcomes. *Front Neurol* 2021; 12: 712060. 2021/09/25. DOI: 10.3389/fneur.2021.712060.
16. Population Division of the Department of Economic and Social Affairs of the United Nations Secretariat. *World Population Prospects 2022*. <https://population.un.org/wpp/> accessed 7th January 2023
17. Russell JBW, Charles E, Conteh V, et al. Risk factors, clinical outcomes and predictors of stroke mortality in Sierra Leoneans: A retrospective hospital cohort study. *Ann Med Surg (Lond)* 2020; 60: 293-300. 2020/11/19. DOI: 10.1016/j.amsu.2020.10.067.
18. World Health Organization, The ICD-10 Classification of Mental and Behavioural Disorders: Diagnostic Criteria for Research (Geneva: World Health Organization, 1992).
19. Bamford J, Sandercock P, et al. A prospective study of acute cerebrovascular disease in the community: the Oxfordshire Community Stroke Project--1981-86. 2. Incidence, case fatality rates and overall outcome at one year of cerebral infarction, primary intracerebral and subarachnoid haemorrhage. *J Neurol Neurosurg Psychiatry*. 1990 Jan;53(1):16-22. doi: 10.1136/jnnp.53.1.16.
20. Quinn TJ, Langhorne P and Stott DJ. Barthel index for stroke trials: development, properties, and application. *Stroke* 2011; 42: 1146-1151. 2011/03/05. DOI: 10.1161/strokeaha.110.598540.
21. Harris PA, Taylor R, Minor BL, et al. The REDCap consortium: Building an international community of software platform partners. *J Biomed Inform* 2019; 95: 103208. 2019/05/13. DOI: 10.1016/j.jbi.2019.103208.
22. Statacorp. Stata Multivariate Statistics Reference Manual Release 11. 2009.
23. Thayabaranathan T, Kim J, Cadilhac D, et al. Global Stroke Statistics 2022. *Int J Stroke* 2022; 17474930221123175. 20220817. DOI: 10.1177/17474930221123175.

24. Adoukonou T, Agbétou M, Bangbotché R, et al. Long-Term Mortality of Stroke Survivors in Parakou: 5-Year Follow-Up. *J Stroke Cerebrovasc Dis* 2020; 29: 104785. 20200319. DOI: 10.1016/j.jstrokecerebrovasdis.2020.104785.
25. Kaduka L, Muniu E, Oduor C, et al. Stroke Mortality in Kenya's Public Tertiary Hospitals: A Prospective Facility-Based Study. *Cerebrovasc Dis Extra* 2018; 8: 70-79. 20180612. DOI: 10.1159/000488205.
26. Rost NS, Bottle A, Lee JM, et al. Stroke Severity Is a Crucial Predictor of Outcome: An International Prospective Validation Study. *Journal of the American Heart Association* 2016; 5: e002433. DOI: doi:10.1161/JAHA.115.002433.
27. Zhang R, Wang Y, Fang J, et al. Worldwide 1-month case fatality of ischaemic stroke and the temporal trend. *Stroke Vasc Neurol* 2020; 5: 353-360. 20200713. DOI: 10.1136/svn-2020-000371.
28. Pinho J, Costa AS, Araújo JM, et al. Intracerebral hemorrhage outcome: A comprehensive update. *J Neurol Sci* 2019; 398: 54-66. 20190114. DOI: 10.1016/j.jns.2019.01.013.
29. Phull M, Grimes CE, Kamara TB, et al. What is the financial burden to patients of accessing surgical care in Sierra Leone? A cross-sectional survey of catastrophic and impoverishing expenditure. *BMJ Open* 2021; 11. DOI: 10.1136/bmjopen-2020-039049.
30. Aenishanslin J, Kamara A and Magnusson L. Experiences accessing and using rehabilitation services for people with physical disabilities in Sierra Leone. *Disabil Rehabil* 2020; 1-10. 2020/05/01. DOI: 10.1080/09638288.2020.1755375.
31. Pieterse P and Longe T. When free healthcare is not free. Corruption and mistrust in Sierra Leone's primary healthcare system immediately prior to the Ebola outbreak. *Int Health* 2015; 7: 400-404. 2015/04/25. DOI: 10.1093/inthealth/ihv024.
32. Idriss A, Diaconu K, Zol C, et al. Rural-urban health-seeking behaviours for non-communicable diseases in Sierra Leone. *BMJ Global Health* 2020; 5: e002024. DOI: 10.1136/bmjgh-2019-002024.
33. Etminan N, Chang HS, Hackenberg K, et al. Worldwide Incidence of Aneurysmal Subarachnoid Hemorrhage According to Region, Time Period, Blood Pressure, and Smoking Prevalence in the Population: A Systematic Review and Meta-analysis. *JAMA Neurol* 2019; 76: 588-597. 2019/01/20. DOI: 10.1001/jamaneurol.2019.0006.
34. Al-Mistarehi AH, Elsayed MA, Ibrahim RM, et al. Clinical Outcomes of Primary Subarachnoid Hemorrhage: An Exploratory Cohort Study from Sudan. *Neurohospitalist* 2022; 12: 249-263. 20220218. DOI: 10.1177/1941874421116825.
35. Alkali NH, Bwala SA, Akano AO, et al. Stroke risk factors, symptoms, and 30-day case fatality in Abuja, Nigeria. *Niger Med J* 2013; 54: 129-135. DOI: 10.4103/03001652.110051.
36. Waweru P and Gatimu SM. Mortality and functional outcomes after a spontaneous subarachnoid haemorrhage: A retrospective multicentre cross-sectional study in Kenya. *PLoS One* 2019; 14: e0217832. 20190612. DOI: 10.1371/journal.pone.0217832.
37. Roushdy T, Aref H, Kesraoui S, et al. Stroke services in Africa: What is there and what is needed. *Int J Stroke* 2022; 17: 972-982. 20220104. DOI: 10.1177/1747493021106415.
38. Dokponou YCH, Kotecha J, Bandyopadhyay S, et al. Continental Survey of Access to Diagnostic Tools and Endovascular Management of Aneurysmal Subarachnoid Hemorrhage in Africa. *Front Surg* 2021; 8: 690714. 20210720. DOI: 10.3389/fsurg.2021.690714.
39. Prasitlumkum N, Cheungpasitporn W, Chokesuwattanaskul A, et al. Diagnostic accuracy of smart gadgets/wearable devices in detecting atrial fibrillation: A systematic review and meta-analysis. *Arch Cardiovasc Dis* 2021; 114: 4-16. 20200910. DOI: 10.1016/j.acvd.2020.05.015.
40. Pyykönen M, Linna M, Tykkyläinen M, et al. Patient-specific and healthcare real-world costs of atrial fibrillation in individuals treated with direct oral anticoagulant agents or warfarin. *BMC Health Serv Res* 2021; 21: 1299. 20211203. DOI: 10.1186/s12913-021-07125-5.
41. Ramakumar V, Benz AP and Karthikeyan G. Long-term oral anticoagulation for atrial fibrillation in low and middle income countries. *Indian Heart J* 2021; 73: 244-248. 20210210. DOI: 10.1016/j.ihj.2021.02.003.

42. Bonny A, Ngantcha M, Yuyun MF, et al. Cardiac arrhythmia services in Africa from 2011 to 2018: the second report from the Pan African Society of Cardiology working group on cardiac arrhythmias and pacing. *Europace* 2020; 22: 420-433. DOI: 10.1093/europace/euz354.
44. Meretoja A, Strbian D, Putaala J, et al. SMASH-U: a proposal for etiologic classification of intracerebral hemorrhage. *Stroke* 2012; 43: 2592-2597. 2012/08/04. DOI: 10.1161/strokeaha.112.661603.
45. Sarfo FS, Ovbiagele B, Gebregziabher M, et al. Unraveling the risk factors for spontaneous intracerebral hemorrhage among West Africans. *Neurology* 2020; 94: e998-e1012. 2020/02/23. DOI: 10.1212/wnl.00000000000009056.
46. Parati G, Lackland DT, Campbell NRC, et al. How to Improve Awareness, Treatment, and Control of Hypertension in Africa, and How to Reduce Its Consequences: A Call to Action From the World Hypertension League. *Hypertension* 2022; 79: 1949-1961. 20220531. DOI: 10.1161/hypertensionaha.121.18884.
47. Ojagbemi A and Owolabi M. Predictors of functional dependency after stroke in Nigeria. *J Stroke Cerebrovasc Dis* 2013; 22: e381-387. 20130513. DOI: 10.1016/j.jstrokecerebrovasdis.2013.04.015.
48. Imarhiagbe FA and Abidakun A. Functional motor recovery in stroke survivors-determinants in a sub-Saharan african stroke unit. *East Afr Med J* 2014; 91: 119-124.
49. Harvey RL. Predictors of Functional Outcome Following Stroke. *Phys Med Rehabil Clin N Am* 2015; 26: 583-598. 20150926. DOI: 10.1016/j.pmr.2015.07.002.
50. Sarfo FS and Ovbiagele B. Key determinants of long-term post-stroke mortality in Ghana. *J Neurol Sci* 2022; 434: 120123. 2022/01/03. DOI: 10.1016/j.jns.2021.120123.
51. Marshall LJ, Wang Y, Crichton S, et al. The effects of socioeconomic status on stroke risk and outcomes. *Lancet Neurol* 2015; 14: 1200-1218. 2015/11/20. DOI: 10.1016/s1474-4422(15)00200-8.
52. Rannikmae K, Woodfield R, Anderson CS, et al. Reliability of intracerebral hemorrhage classification systems: A systematic review. *Int J Stroke* 2016; 11: 626-636. 2016/04/20. DOI: 10.1177/1747193016641962.