

Central Lancashire Online Knowledge (CLoK)

Title	The use of statistical parametric mapping to determine altered movement	
	patterns in people with chronic low back pain	
Туре	Article	
URL	https://clok.uclan.ac.uk/46396/	
DOI	https://doi.org/10.1016/j.jbiomech.2023.111601	
Date	2023	
Citation	Wattananon, Peemongkon, Kongoun, Sasithorn, Chohan, Ambreen and Richards, James (2023) The use of statistical parametric mapping to determine altered movement patterns in people with chronic low back pain. Journal of Biomechanics, 153. ISSN 0021-9290	
Creators	Wattananon, Peemongkon, Kongoun, Sasithorn, Chohan, Ambreen and Richards, James	

It is advisable to refer to the publisher's version if you intend to cite from the work. https://doi.org/10.1016/j.jbiomech.2023.111601

For information about Research at UCLan please go to http://www.uclan.ac.uk/research/

All outputs in CLoK are protected by Intellectual Property Rights law, including Copyright law. Copyright, IPR and Moral Rights for the works on this site are retained by the individual authors and/or other copyright owners. Terms and conditions for use of this material are defined in the <u>http://clok.uclan.ac.uk/policies/</u>

1 2 PATTERNS IN PEOPLE WITH CHRONIC LOW BACK PAIN 3 4 5 6 7 Peemongkon Wattananon¹, Sasithorn Kongoun¹, Ambreen Chohan², Jim Richards² ¹ Spine Biomechanics Laboratory, Faculty of Physical Therapy, Mahidol University. 999 Phuttamonthon 4 Road, Salaya, Nakhon Pathom 73170, Thailand. 8 9 Email: peemongkon.wat@mahidol.ac.th; sasithorn.kon@mahidol.edu 10 ² Allied Health Research Unit, University of Central Lancashire. 11 Preston, Lancashire PR1 2HE, United Kingdom. 12 Email: AChohan@uclan.ac.uk; JRichards@uclan.ac.uk 13 14 15 This study was funded by Mahidol University Mini-Research Cluster Fund (MiniRC) and Mahidol University (MU 16 Talent program). 17 18 This study was approved by the Institutional Review Board of Mahidol University (COA No. MU-CIRB 19 2020/084.1806). 20 21 The authors have no disclosures or conflict of interest to report. 22 23 Address correspondence to: 24 Peemongkon Wattananon, PT, PhD.

- 25 Mail stop: Faculty of Physical Therapy, Mahidol University.
- 26 999 Phuttamonthon 4 Road, Salaya, Nakhon Pathom 73170 Thailand.
- 27 Email: peemongkon.wat@mahidol.edu

THE USE OF STATISTICAL PARAMETRIC MAPPING TO DETERMINE ALTERED MOVEMENT

1 Highlights

- 2 3 4 5 • Statistical Parametric Mapping differentiated segment movement contributions and timing
 - Lumbar segments contributed more during flexion, extension, and lateral bending •
 - Thoracic segments contributed more in rotation during rotation in sitting
 - Clinicians should observe relative contributions between segments

- **1** ABSTRACT
- 2

3 Kinematics studies have generally focused on the quantity of movement using discrete parameters such as 4 maximum and minimum angles to compare between people with chronic low back pain (CLBP) and healthy 5 individuals. However, discrete parameters cannot be used to fully describe movement patterns and segmental 6 contributions. This study aimed to explore the use of Statistical Parametric Mapping (SPM) to characterize quality of 7 movement by examining if differences in movement patterns exist between groups, and within-group segmental 8 contributions, during active movement tests. Twenty-one individuals with CLBP and nine healthy individuals were 9 recruited. Inertial Measurement Unit (IMUs) were attached at thoracic (T3) and lumbar (L1) spine, and pelvis (S1) to 10 collect active trunk flexion, extension, rotation, and lateral bend. SPM was used to analyze between-group movement 11 patterns and within-group segmental contributions. SPM revealed no significant differences (P>0.05) between groups. 12 However, a greater lumbar contribution (P < 0.001) was observed during 10-40% of flexion followed by a greater 13 pelvic contribution (P<0.001) during 60-90% of flexion, while a greater lumbar than thoracic contribution (P<0.001) 14 was observed during flexion and the return to upright position in individuals with CLBP. Individuals with CLBP used 15 a greater thoracic contribution compared to lumbar contribution (P < 0.001) during rotation, while a greater lumbar 16 contribution compared to pelvic contribution was observed (P<0.001) during lateral bending. Our findings suggest 17 that SPM approach was able to detect differences in thoracic, lumbar, and pelvic velocity contributions and timings 18 between segments in individuals with CLBP. These findings may help improving inter-rater reliability of clinical 19 observations. 20 21 Keywords: Statistical Parametric Mapping; Movement Patterns; Quality of Movement; Segmental contributions; Low 22 back pain 23 24 Word count 25 Abstract = 24826 Main text = 308027

1 1. INTRODUCTION

2 A recent clinical practice guideline suggests clinicians should use specific muscle activation and movement 3 control exercises for patients with chronic low back pain (CLBP) who have underlying movement control impairments 4 (George et al., 2021). This would suggest the need to assess active trunk movement in patients with CLBP. Clinical 5 observations of aberrant movement patterns during active trunk movements in different planes (flexion/extension, 6 rotation, and lateral bend) are a critical component in physical examination (Biely et al., 2014; Gombatto et al., 2007; 7 Hicks et al., 2005; Marich et al., 2020; O'Sullivan, 2000). Although studies have demonstrated the ability of clinical 8 observations to identify patients with CLBP, the reliability of such clinical observation studies have shown the inter-9 rater reliability to be less than the intra-rater reliability (Hicks et al., 2003; Rabin et al., 2013). This could be due to 10 different clinicians considering different segments or references to determine the presence, or not, of aberrant 11 movements.

12 Several studies have attempted to characterize movement patterns in patients with CLBP and healthy 13 individuals (Laird et al., 2014; Zheng et al., 2022). These studies focused on the quantity of movement, rather than 14 quality of movement (Laird et al., 2014; Zheng et al., 2022). One systematic review showed that patients with LBP 15 had less lumbar motion and moved slower than healthy individuals (Laird et al., 2014). However, findings were 16 derived from discrete parameters (e.g., maximum range of motion and velocity). This over-simplification of discrete 17 parameter analysis may limit the clinical utility of such results. In addition, clinical observations of aberrant movement 18 patterns primarily focus on the quality of movement (e.g., segmental contributions and timing) of the thoracic and 19 lumbar spine and pelvis (Biely et al., 2014; Hicks et al., 2005; Marich et al., 2020; O'Sullivan, 2000; Rabin et al., 20 2013). Those aberrant movement patterns are commonly described in terms of movement control and/or segmental 21 coordination in the same patient, rather than comparing against healthy individuals. For instance, altered lumbopelvic 22 rhythm is defined as a greater lumbar contribution in the early phase of active forward bending. This suggests that the 23 within-group characterization of movement patterns, particularly in people with CLBP, would be valuable to explore. 24 Although kinematic data revealed that patients with LBP had reduced lumbar range of motion and velocity compared 25 with healthy individuals (Laird et al., 2014), it is difficult for clinicians to determine whether patients had altered range 26 of motion and velocity. Therefore, the inclusion of the objective assessment of movement contributions and timing 27 into any analysis into clinical practice may help our understanding of the efficacy of, and response to, interventions in 28 individuals with LBP.

29 The measurement of joint and segment kinematics using inertial measurement units (IMUs) have 30 demonstrated concurrent validity against optoelectronic system, and between-session test-retest reliability to track 31 lumbopelvic motion in three-dimensional space (Bauer et al., 2016; Bauer et al., 2015; Seel et al., 2014), opening up 32 the opportunity for assessments of quality of movement and stability within clinical settings (Budini et al., 2018). Due 33 to the portable nature and relatively low cost of IMUs, these can be used to evaluate a variety of physical activities 34 and even have the potential to assess movement in the community (Fennema et al., 2019). Therefore, kinematic data 35 from IMUs may provide information to help clinicians better identify aberrant movement patterns, thereby improving 36 interrater reliability.

1 Statistical parametric mapping (SPM) has been recently utilized to investigate time-series data including 2 kinematics, electromyography, and ground reaction force data (Papi et al., 2020; Pataky et al., 2013; Robinson et al., 3 2015). Papi et al. (2020) utilized SPM to analyze movement patterns during functional movements including sit to 4 stand and walking in patients with LBP. The findings support the advantage of SPM over discrete parameter analysis 5 as it takes the amplitude and timing over the whole time series into consideration (Papi et al., 2020). Therefore, this 6 approach can be used to investigate movement patterns, as well as contribution and timing during active movement 7 tests which is a critical part of the physical examination for patients with CLBP. This may help refine clinical 8 observations, particularly when considering the timing and relative segmental contributions during specific movement 9 tasks.

10 Therefore, this study aimed to compare movement patterns recorded using IMUs between people with CLBP 11 and healthy individuals during active trunk flexion, rotation, and lateral bending, as well as within-group movement 12 contributions and timing of the thoracic and lumbar spine and pelvis. We hypothesized that people with CLBP would 13 demonstrate different movement patterns, segmental contributions and timing when compared with healthy 14 individuals.

15

16 2. METHODS

17 2.1 Participants

18 Twenty-one individuals with CLBP and nine healthy individuals aged between 18 and 40 years were recruited 19 as part of another ongoing study. The inclusion criteria for CLBP were having low back pain for over 3 months, or a 20 recurrent pattern of LBP with at least two episodes that interfered with activities of daily living and/or required 21 treatment. The inclusion criteria for healthy individuals were no previous history of LBP for 6 months. Participants 22 were excluded if they had clinical signs of systematic disease, definitive neurologic signs (weakness or numbness in 23 the lower extremity), previous spinal surgery, or any lower extremity condition that could alter trunk movement. All 24 participants provided written informed consent before data collection, and the study was approved by the institutional 25 review board of Mahidol University (COA No. MU-CIRB 2020/084.1806). We performed a sample size calculation 26 to ensure that the data available was sufficient to perform the planned data analysis. The sample size calculation was 27 based on a meta-analysis which demonstrated that patients with LBP had significantly slower movement than those 28 healthy individuals with a standardized mean difference of 1.24 (Laird et al., 2014). The sample size calculation was 29 performed using G*Power (version 3.1.9.6) using an independent one tailed t-test with the overall effect size (Cohen's 30 d) of 1.24, an alpha value of 0.05 and an 80% power. This yielded a sample size of at least 9 participants in each 31 group, indicating our sample would be sufficient for the planned statistical analysis.

32 2.2 Instrumentation

Three Trigno Avanti Inertial Measurement Units (Delsys Inc., MA, USA) were attached to the thoracic (T3),
 lumbar (L1), and sacral (S1) spinous processes (Figure 1). These data were used to record flexion, rotation, and lateral
 bending movement patterns during active movement tests. Data collection was performed using EMGworks 4.7.3
 (Delsys Inc., MA, USA) at 370 Hz. This system has been used to investigate human movement in several studies

37 (Costello et al., 2020; Khobkhun et al., 2020; Kolářová et al., 2022), and preliminary work demonstrated the average

coefficient of multiple determination was 0.85, indicating excellent test-retest reliability. The average cross correlation coefficients between IMU and electromagnetic tracking system were 0.95, 0.88, and 0.95 in sagittal,

3 frontal, and transverse planes, respectively. These findings indicate excellent concurrent validity.

4 2.3 Procedure

5 Demographic data including age, sex, BMI, and low back pain characteristics were recorded, and the IMUs 6 attached. The participants were then asked to perform 3 consecutive repetitions of active movement tests including 7 trunk flexion, extension, rotation to the right and left in sitting, and lateral bending to the right and left at a comfortable 8 pace with 1-minute rest between tasks. One practice trial was allowed to check the data and to familiarize each 9 participant with the tasks.

10 2.4 Data analysis

11 Data was exported in c3d format and analysis was performed using Visual3D (C-Motion Inc., MD, USA). 12 All data were filtered using a lowpass Butterworth filter at 15 Hz, and the start and end points were defined which 13 allowed the identification of the "Go" and "Return" phases for each repetition of the movements using the IMU 14 positioned on T3. The data were then time-normalized to 101 data points to represent 100% of the movement. Data 15 from T3 were used to represent trunk velocity. L1 relative to S1 represented lumbar velocity, while T3 relative to L1 16 represented thoracic velocity, and S1 relative to the global reference represented pelvic velocity. Therefore, total trunk, 17 thoracic, lumbar, and pelvic velocities in the three planes of movement were derived. However, participants can have 18 deviation in frontal and/or transverse planes either to the right or left during flexion and extension. Therefore, we used 19 the absolute deviation during the movement for data analysis.

20 2.5 Statistical analysis

21 Statistical analyses were performed using SPSS program (SPSS version 21, IBM Corp., NY, USA) to 22 compare demographic data between groups. SPM was used to determine differences in movement patterns between 23 people with CLBP and healthy individuals and individual segment contributions and timing within each group. Python 24 (version 3.9.4, Python Software Foundation, https://www.python.org) with open-source code were used to perform 25 SPM analysis. The SPM two-sample t test is based on Random Field Theory (Adler & Taylor, 2007), which is the 26 vector-field equivalent to the t-test which was calculated at each percent of movement to establish a smooth random 27 curve and 95% confidence interval critical threshold (Pataky et al., 2013; Robinson et al., 2015). Significant difference 28 occurs when the SPM curves (SPM(t)) cross the critical threshold (t*) at any percent of movement, and associated p-29 values were calculated using Random Field Theory (Adler & Taylor, 2007; Pataky et al., 2013; Robinson et al., 2015). 30 When the t-values are greater than the upper border or less than the lower border this would be considered as statistical 31 difference. In other words, the area outside the upper and lower critical t-value would demonstrate the presence of a 32 statistical difference. SPM two-sample t tests were used to compare between group angular velocities for the 33 movements in each plane of movement for each segment (total trunk, thoracic, lumbar, and pelvic angular velocity). 34 In addition, within-group SPM paired t tests were performed between segments (thoracic vs lumbar spine and lumbar 35 spine vs pelvis) for the primary planes of movement for each task for the two groups separately.

36

37 3. RESULTS

Demographic data and clinical characteristics are presented in Table 1. No significant differences (P>0.05)
were seen for age, sex, and BMI between groups. People with CLBP had an average pain in the past 7 days of 4 out
of 10. More than 11 (57.2%) had a history of pain for greater than one year, and 14 (66.7%) had pain for at least half
the days or nearly every day. Results also demonstrated that LBP interfered with their daily living, work, social life,
and chores. SPM revealed no significant differences (P>0.05) in movement patterns between people with CLBP and
healthy individuals for each segment and each plane of movement.

7 Within-group segment contributions and timing comparisons between lumbar spine and pelvis (Figure 2; 8 lumbar spine vs pelvis during forward bend) demonstrated that people with CLBP used a greater lumbar contribution 9 during the early phase of movement (10-40%) followed by a greater pelvic contribution during the later phase (60-10 90%) during flexion, while healthy individuals used a shared lumbar and pelvic contribution in the early phase with a 11 greater pelvic contribution during the middle phase (30-70%). During the return to an upright position (Figure 2; 12 lumbar spine vs pelvis during return to upright), individuals with CLBP used a greater pelvic contribution during the 13 early phase (0-30%) followed by a greater pelvic contribution (40-100%), indicating an over reliance on a particular 14 segment within the different phases of the movement. Conversely, healthy individuals seemed to use a more shared 15 pattern of movement between the segments. When considering the thoracic and lumbar contributions (Figure 2; 16 thoracic vs lumbar spine), individuals with CLBP tended to use a greater lumbar contribution than healthy individuals 17 during flexion and the return to upright position.

18 Differences (P<0.001) were also seen in thoracic and lumbar contributions and timing during rotation and 19 return to the central position (Figure 3). During rotation and return, individuals with CLBP tended to use a greater 20 thoracic contribution and less lumbar contribution, while healthy individuals used a more shared pattern between the 21 thoracic and lumbar spine. Similarly, differences (P < 0.001) were observed between the lumbar and pelvic 22 contributions during lateral bending and return to an upright posture (Figure 4) in which individuals with CLBP tended 23 to use a greater lumbar contribution than pelvic contribution, whereas healthy individuals again used a more shared 24 pattern of movement between the lumbar spine and pelvis. However, there were no significant differences (P>0.05) 25 in the segment contributions and timings within-group during the active trunk extension task.

26

27 4. DISCUSSION

28 Using SPM analysis, we did not find any significant differences in movement patterns between people with 29 CLBP and healthy individuals. Non-significant results indicate that both people with CLBP and healthy individuals 30 have similar segment velocity profiles when performing trunk flexion, extension, rotation to the right and left in sitting, 31 and lateral bending to the right and left. Our findings were not consistent with a previous systematic review which 32 could be due to methodological differences in which the studies included used maximum velocity using discrete point 33 analysis, where our approach considered the whole time-series data (Laird et al., 2014; Zheng et al., 2022). In addition, 34 their results did not specify acute or chronic LBP, whereas our study included CLBP. Evidence suggests that 35 individuals with acute and subacute low back pain use stiffening strategies to minimize pain exacerbation (Jones et 36 al., 2012; Wattananon et al., 2022), which would result in lower lumbar velocities. However, in cases of CLBP, 1 individuals might have adopted a different movement strategy, resulting in adapting a higher lumbar segment velocity

2 (van Dieën et al., 2019).

People with CLBP demonstrated greater lumbar contribution during the early phase of forward bend
comparing with pelvis and thoracic spine. Greater lumbar contributions have been associated with increases in the risk
of injury (Sahrmann et al., 2017; Van Dillen et al., 2003), and it has been suggested that excessive lumbar movement
during daily activities can cause an accumulation of soft tissue stress leading to microtrauma and low back pain
(Sahrmann et al., 2017; Van Dillen et al., 2003).

8 To the best of our knowledge, no previous study has investigated angular velocity during rotation in sitting. 9 Only one study using the SPM approach has investigated range of motion in different spinal segments (thoracic, 10 lumbar, and pelvic segments) and planes (sagittal, frontal, and transverse planes) (Papi et al., 2020). They found that 11 patients with LBP had greater thoracic rotation than healthy individuals (Papi et al., 2020). Therefore, a greater 12 thoracic contribution might be a compensatory movement to achieve the rotation task.

Similarly, individuals with CLBP used a greater lumbar contribution than pelvic contribution during lateral bending and the return to an upright posture. This was again in contrast to healthy individuals who tended to use a shared pattern of movement between the lumbar and pelvic segments. Such greater lumbar contribution could increase stress to the soft tissues surrounding the lumbar spine, thereby causing pain, or contributing to persistence or recurrence of low back pain (Sahrmann et al., 2017; Van Dillen et al., 2003).

Although our kinematic data did not show significant differences between individuals with CLBP and healthy individuals, within-group comparisons demonstrated significant differences in the contributions and timing between segments. These within-group findings are consistent with the assessment of quality of movement when clinicians observe aberrant movements during active trunk movement during physical examinations (Biely et al., 2014; Gombatto et al., 2007; Hicks et al., 2005; Marich et al., 2020; O'Sullivan, 2000). Our results suggest lumbar spine and pelvis as reference segments during trunk flexion which could be used to refine definition of aberrant movement; thereby improving inter-rater reliability of clinical observation.

25 The use of SPM allows the exploration of differences over the whole time series which has advantages over 26 the more common approach of using discrete parameter analysis (e.g., peak, maximum, minimum, etc.) to represent 27 the movement characteristics (Laird et al., 2014; Zheng et al., 2022). Discrete parameters do not represent movement 28 patterns because they can occur at different time points during the movement (Papi et al., 2020). For instance, the 29 contribution between the lumbar spine and pelvis during active trunk flexion and return in Figure 2 clearly showed 30 that peak lumbar velocity occurred before peak pelvic velocity. However, using discrete peak velocity showed no 31 differences in segmental contributions, which could lead to misinterpretation of findings. Unlike discrete parameter 32 analysis, SPM approach takes both spatial and temporal data into consideration (Adler & Taylor, 2007; Pataky et al., 33 2013; Robinson et al., 2015). SPM can identify the time when significant differences between lumbar and pelvic 34 velocity have occurred. In addition, SPM offers more control for both type I and II errors (Pataky et al., 2013; Robinson 35 et al., 2015), and appears to provide a better approach when analyzing movement patterns.

The findings of this study should be considered in light of the following limitations. We used differencesbetween group to calculate sample size. Therefore, our healthy individual sample size might be relatively small for

2 needs to be recalculated for each purpose (e.g., between-group and within-group comparisons). Although differences 3 between segments were found in the CLBP group and not in the control group, indicating a different contribution of 4 the different segments, a limitation is the differences in the two group sizes for the within group analysis, with the 5 differences seen in the larger CLBP group (n=21). Therefore, it is possible that the lack of differences seen in the 6 control group (n=9) could be due to the smaller group size. We investigated individuals with CLBP which could limit 7 the generalizability to other low back pain populations. We attached IMUs on the skin which could introduce soft 8 tissue noise affecting kinematic data, however this protocol was able to determine different relative contributions of 9 segments within individuals with CLBP when compared with healthy controls. Lastly, our study aimed to explore the 10 utility of the SPM approach to characterize quality of movement across the whole time series; therefore, we considered 11 the contributions and timings of the movement of neighboring pairs of segments rather than all the spinal segments 12 relative to the pelvis. We did not control for multiple comparisons for within-group segment contributions and timing

within-group comparisons. Replication of our study with larger sample size is necessary, and sample size calculation

- 13 comparisons or allow for confounder variables such as age, BMI, and sex.
- 14

1

15 5. CONCLUSION

16 The SPM approach was able to determine differences in thoracic, lumbar, and pelvic velocity contributions 17 and timings between segment in individuals with CLBP. Our results suggest clinical observation should focus on the 18 contribution and timing between lumbar spine and pelvis, as well as thoracic and lumbar spine during active trunk 19 flexion, the thoracic and lumbar spine during rotation in sitting, and lumbar spine and pelvis during lateral bending. 20 These findings may help improving inter-rater reliability of clinical observations.

21

22 ACKNOWLEDGEMENT

We would like to thank Mahidol University for providing travel grant (Academic and staff mobility program).
 This study was funded by Mahidol University Mini-Research Cluster Fund (MiniRC) and Mahidol University (MU
 Talent). We also would like to thank all participants for providing data in this study.

26

27 REFERENCES

- 28 Adler, R. J., & Taylor, J. E. (2007). Random fields and geometry (Vol. 80). Springer.
- Biely, S. A., Silfies, S. P., Smith, S. S., & Hicks, G. E. (2014). Clinical observation of standing trunk movements:
 what do the aberrant movement patterns tell us? *J Orthop Sports Phys Ther*, 44(4), 262-272.
- Budini, K., Richards, J., Cole, T., Levine, D., Trede, R., George, L. S., & Selfe, J. (2018). An exploration of the use
 of Inertial Measurement Units in the assessment of dynamic postural control of the knee and the effect of bracing
 and taping. *Physiother Pract Res*, *39*, 91-98.
- Costello, K. E., Eigenbrot, S., Geronimo, A., Guermazi, A., Felson, D. T., Richards, J., & Kumar, D. (2020).
 Quantifying varus thrust in knee osteoarthritis using wearable inertial sensors: A proof of concept. *Clin Biomech*
- **36** (*Bristol, Avon*), 80, 105232.

- Fennema, M. C., Bloomfield, R. A., Lanting, B. A., Birmingham, T. B., & Teeter, M. G. (2019). Repeatability of
 measuring knee flexion angles with wearable inertial sensors. *Knee*, 26(1), 97-105.
- George, S. Z., Fritz, J. M., Silfies, S. P., Schneider, M. J., Beneciuk, J. M., Lentz, T. A., Gilliam, J. R., Hendren, S.,
 & Norman, K. S. (2021). Interventions for the management of acute and chronic low back pain: revision 2021.
- 5 J Orthop Sports Phys Ther, 51(11), 1-60.
- Gombatto, S. P., Collins, D. R., Sahrmann, S. A., Engsberg, J. R., & Van Dillen, L. R. (2007). Patterns of lumbar
 region movement during trunk lateral bending in 2 subgroups of people with low back pain. *Phys Ther*, 87(4),
 441-454.
- 9 Hicks, G. E., Fritz, J. M., Delitto, A., & McGill, S. M. (2005). Preliminary development of a clinical prediction rule
 10 for determining which patients with low back pain will respond to a stabilization exercise program. *Arch Phys* 11 *Med Rehabil*, 86(9), 1753-1762.
- Hicks, G. E., Fritz, J. M., Delitto, A., & Mishock, J. (2003). Interrater reliability of clinical examination measures for
 identification of lumbar segmental instability. *Arch Phys Med Rehabil*, 84(12), 1858-1864.
- Jones, S. L., Henry, S. M., Raasch, C. C., Hitt, J. R., & Bunn, J. Y. (2012). Individuals with non-specific low back
 pain use a trunk stiffening strategy to maintain upright posture. *J Electromyogr Kinesiol*, 22(1), 13-20.
- Khobkhun, F., Hollands, M. A., Richards, J., & Ajjimaporn, A. (2020). Can we accurately measure axial segment
 coordination during turning using inertial measurement units (IMUs)? *Sensors (Basel)*, 20(9).
- 18 Kolářová, B., Richards, J., Haltmar, H., Lippertová, K., Connell, L., & Chohan, A. (2022). The effect of motor imagery
 19 on quality of movement when performing reaching tasks in healthy subjects: A proof of concept. *J Bodyw Mov* 20 *Ther*, 29, 161-166.
- Laird, R. A., Gilbert, J., Kent, P., & Keating, J. L. (2014). Comparing lumbo-pelvic kinematics in people with and
 without back pain: a systematic review and meta-analysis. *BMC Musculoskelet Disord*, 15, 229.
- Marich, A. V., Hwang, C. T., Sorensen, C. J., & van Dillen, L. R. (2020). Examination of the lumbar movement
 pattern during a clinical test and a functional activity test in people with and without low back pain. *Pm r*, *12*(2),
 140-146.
- O'Sullivan, P. B. (2000). Lumbar segmental 'instability': clinical presentation and specific stabilizing exercise
 management. *Man Ther*, 5(1), 2-12.
- Papi, E., Bull, A. M. J., & McGregor, A. H. (2020). Alteration of movement patterns in low back pain assessed by
 Statistical Parametric Mapping. *J Biomech*, *100*, 109597.
- Pataky, T. C., Robinson, M. A., & Vanrenterghem, J. (2013). Vector field statistical analysis of kinematic and force
 trajectories. *J Biomech*, 46(14), 2394-2401.
- Rabin, A., Shashua, A., Pizem, K., & Dar, G. (2013). The interrater reliability of physical examination tests that may
 predict the outcome or suggest the need for lumbar stabilization exercises. *J Orthop Sports Phys Ther*, 43(2), 83 90.
- Robinson, M. A., Vanrenterghem, J., & Pataky, T. C. (2015). Statistical Parametric Mapping (SPM) for alpha-based
 statistical analyses of multi-muscle EMG time-series. *J Electromyogr Kinesiol*, 25(1), 14-19.

- Sahrmann, S., Azevedo, D. C., & Dillen, L. V. (2017). Diagnosis and treatment of movement system impairment
 syndromes. *Braz J Phys Ther*, 21(6), 391-399.
- Seel, T., Raisch, J., & Schauer, T. (2014). IMU-based joint angle measurement for gait analysis. Sensors (Basel),
 14(4), 6891-6909.
- van Dieën, J. H., Reeves, N. P., Kawchuk, G., van Dillen, L. R., & Hodges, P. W. (2019). Motor control changes in
 low back pain: divergence in presentations and mechanisms. *J Orthop Sports Phys Ther*, 49(6), 370-379.
- Van Dillen, L. R., Sahrmann, S. A., Norton, B. J., Caldwell, C. A., McDonnell, M. K., & Bloom, N. J. (2003).
 Movement system impairment-based categories for low back pain: stage 1 validation. *J Orthop Sports Phys Ther*, 33(3), 126-142.
- Wattananon, P., Silfies, S. P., & Wang, H. K. (2022). Patients with low back pain use stiffening strategy to compensate
 for movement control during active prone hip rotation: A cross-sectional study. *J Back Musculoskelet Rehabil*,
 35(2), 373-382.
- 13 Zheng, Z., Wang, Y., Wang, T., Wu, Y., & Li, Y. (2022). A systematic review and meta-analysis on comparative
- 14 kinematics in the lumbopelvic region in the patients suffering from spinal pain. *J Healthc Eng*, 2022, 7369242.

Table 1. Demographic data and clinical characteristics

Parameter	CON(n=9)	CLBP (n = 21)	
Age (year)	25.0 (6.6)	30.5 (6.9)	
Sex (%female)	6 (75.0%)	11 (52.4%)	
$BMI (kg/m^2)$	21.9 (3.7)	25.2 (4.3)	
Average pain in the past 7 days		4.0 (1.8)	
Duration			
- Less than 1 months		3 (14.3%)	
- 1-3 months		3 (14.3%)	
- 3-6 months		2 (9.5%)	
- 6 months-1 year		1 (4.8%)	
- 1-5 years		9 (42.9%)	
- more than 5 years		3 (14.3%)	
Frequency in the past 6 months			
- Every day or nearly every day		3 (14.3%)	
- At least half the days		11 (52.4%)	
- Less than half the days		7 (33.3%)	
Daily interference			
- Not at all		3 (14.3%)	
- A little bit		8 (38.1%)	
- Somewhat		3 (14.3%)	
- Ouite a bit		5 (23.8%)	
- Verv much		2 (9.5%)	
Work interference			
- Not at all		4 (19.0%)	
- A little bit		8 (38.1%)	
- Somewhat		3 (14.3%)	
- Quite a bit		2 (9.5%)	
- Very much		4 (19.0%)	
Social interference			
- Not at all		7 (33.3%)	
- A little bit		8 (38.1%)	
- Somewhat		2 (9.5%)	
- Quite a bit		0 (0%)	
- Very much		4 (19.0%)	
Chore interference			
- Not at all		4 (19.0%)	
- A little bit		8 (38.1%)	
- Somewhat		4 (19.0%)	
- Quite a bit		2 (9.5%)	
- Very much		3 (14.3%)	

Figure legends



Figure 1. Positioning of IMUs on the thoracic (T3), lumbar (L1), and sacral (S1) spinous processes.



Figure 2. Differences in lumbar and pelvic contribution, as well as thoracic and lumbar contribution during flexion and return to upright



Figure 3. Differences in thoracic and lumbar contribution during rotation to right/left and return



Figure 4. Differences in lumbar and pelvic contribution during lateral bend to right/left and return