

Central Lancashire Online Knowledge (CLoK)

Title	Working effectively with people who have received a diagnosis of personality disorder
Type	Article
URL	https://clock.uclan.ac.uk/46740/
DOI	https://doi.org/10.7748/mhp.2023.e1634
Date	2023
Citation	Elliott, Alison, Lamph, Gary and Kendall, Zoe (2023) Working effectively with people who have received a diagnosis of personality disorder. Mental Health Practice. ISSN 1465-8720
Creators	Elliott, Alison, Lamph, Gary and Kendall, Zoe

It is advisable to refer to the publisher's version if you intend to cite from the work.
<https://doi.org/10.7748/mhp.2023.e1634>

For information about Research at UCLan please go to <http://www.uclan.ac.uk/research/>

All outputs in CLoK are protected by Intellectual Property Rights law, including Copyright law. Copyright, IPR and Moral Rights for the works on this site are retained by the individual authors and/or other copyright owners. Terms and conditions for use of this material are defined in the <http://clock.uclan.ac.uk/policies/>

Working effectively with people who receive a diagnosis of personality disorder

Alison Elliott, Gary Lamph and Zoe Kendall

Citation

Elliott A, Lamph G, Kendall Z (2023) Working effectively with people who receive a diagnosis of borderline personality disorder. Mental Health Practice. doi: 10.7748/mhp.2023.e1634

Peer review

This article has been subject to external double-blind peer review and checked for plagiarism using automated software

Correspondence

alisonelliott0@gmail.com I would prefer my UCLan email to be used – aelliott1@uclan.ac.uk

Conflict of interest

[Q do you have any conflicts of interest to declare?] **No Conflict of interest**

Accepted

14 December 2022

Published online

xxxx

Abstract

Personality disorder is a contentious diagnostic label that is associated with high levels of stigma, leading many practitioners and people with lived experience to call for a change in its use. This article explores this area further by addressing some of the knowledge deficits and highlighting opportunities for self-reflection and analysis. It also focuses on the importance of formulation for making sense of past experience, and as a tool to develop practitioner insight into suitable psychologically informed interpersonal approaches. Nurses frequently encounter people who have been given a diagnosis of personality disorder across various settings. This article provides a critique of this diagnosis and discusses the issues that are associated with this label.

Author details

Alison Elliott, senior lecturer, School of Nursing, University of Central Lancashire, Preston, England; Gary Lamph, senior research fellow in mental health, School of Nursing, University of Central Lancashire, Preston, England; Zoe Kendall, lived experience Knowledge and Understanding Framework (KUF)] development lead, Leeds and York Partnership NHS Foundation Trust, Leeds, England

Twitter handles - @allyel1408, @gazlamph

Keywords

personality disorder, mental health, service user experience, service users, stigma

Aims and intended learning outcomes

This article aims to enhance nurses' understanding of the sensitivities, misunderstandings, stigma and challenges that surround personality disorder. It discusses the important role nurses have in the care and support of people who have this diagnosis, differential diagnosis or associated issues. The article was co-produced, incorporating professional and lived experience perspectives. However, the authors acknowledge that the stance taken is not necessarily representative of all views held in this field of practice, particularly in relation to the use of language and diagnostic labels. After reading this article and completing the time out activities you should be able to:

- » Appreciate the misunderstandings and stigma that people with complex emotional needs commonly experience.
- » Recognise the crucial role of formulation and reflection in your nursing practice.
- » Consider how you could adapt your approach to meet the needs of individual **service users**.
- » Understand your contribution and responsibility in advocating for **service users'** individual needs.

Introduction

Personality disorder is defined as: *'an enduring pattern of inner experience and behaviour that deviates markedly from the expectations of the individual's culture, is pervasive and inflexible, has an onset in adolescence or early adulthood, is stable over time, and leads to distress or impairment'* (American Psychiatric Association (APA) 2013).

Globally, personality disorder affects up to 7.8% of community populations (Winsper et al, 2020) and comorbidity with other mental health conditions is not uncommon (Hayward and Moran 2007). Personality disorders are biopsychosocial disorders that can adversely affect people's ability to function in everyday life. They are often described as developmental conditions in which adverse childhood experiences have psychological effects due to the social, environmental and biological factors that determine a person's ways of coping (Sampson et al 2006).

The DSM-5 TR (APA 2022) describes 10 types of personality disorder, with Borderline Personality Disorder (BPD), also known as emotionally unstable personality disorder (World Health Organization 1990), being the type most commonly diagnosed. These diagnostic labels refer to same condition but are often used interchangeably, despite a misconception they that are different. BPD is characterised by interpersonal difficulties, emotional dysregulation, impulse control and self-image deficits (Lieb et al 2004), and it is often assumed that the term 'borderline' means that someone may be on the 'cusp' of diagnosis. This misconception is concerning because people with this diagnosis are often those with the most unmet needs and highest levels of risk, particularly to themselves (Bohus et al, 2021). Despite the BPD diagnosis being highly contentious, it continues to be used in the Diagnostic Statistical Manual of Mental Disorders, Fifth Edition (DSM-5) (APA 2022), however, in the International Classification of Diseases 11th Revision (ICD-11) (World Health Organization, the way in which personality disorder is diagnosed has moved away from this categorical model whereby a person either has it or does not and must fit within one of the 10 types. Instead, there is a severity continuum with mild, moderate and severe personality difficulties. If severe difficulties are identified, then more descriptive terms relating to specific difficulties can be considered, for example 'with borderline pattern qualifier' (Mulder 2021). The differences in the DSM-5 and ICD-11, alongside the lack of agreement in the influential experts, leading their revisions, may add to the confusion and resentment many service users describe surrounding this diagnosis, while also causing confusion for practitioners.

Considering the systemic failures, high levels of stigma, the aetiology, and high levels of trauma encountered by people who receive this diagnosis, the sensitivities that arise are to be expected (Lamph et al 2021).

If one's personality is fundamentally considered to be a representation of one's being, it may be insensitive to describe it as 'disordered' or flawed in an often-traumatised client group, who may have had challenging childhood experiences. Doing so can activate emotions in the service user that are vulnerable to escalation, and may lead to further dissonance from service and practitioner responses. Therefore, it is important that education, knowledge and awareness are attended to, and that service users are directed to evidence-based interventions and support of benefit to them (Lamph et al 2021).

A study by Lamph et al (2021) explored the sensitivities surrounding the personality disorder label with practitioners and people with lived experience. It found that the importance of receiving sensitive and accurate information when receiving this label was paramount, but in practice this is not always people's experience. Practitioners are relied on to provide accurate information about this diagnosis and its implications; however, this can be challenging for them because of the changes to nurse education standards and the shift towards a generic nursing syllabus (Connell et al 2022). Therefore, enhancing nurses' understanding of personality disorder, the critical debate that surrounds the diagnosis and providing space for reflection and discussion are essential to enhancing working relationships with this misunderstood and highly stigmatised client group.

Reflect on the issues associated with the label of personality disorder. What may be the advantages and disadvantages of receiving this diagnosis? How would you feel if someone you know received this diagnosis? How do you understand the effect of early relationships on personality development and interpersonal effectiveness? Reflect on the information you provide to inform service users about the diagnosis in practice. Does this include written information? Is it clear, hopeful and optimistic?

Alternative conceptualisations of personality disorder

One of the issues associated with the personality disorder diagnosis is that it is deficit focused, locating problems within the individual, and ignoring such issues as past experiences, and current social context (Read and Harper 2022). Therefore, the authors suggest that there may be more helpful ways of conceptualising ‘personality disorder’.

In 2018, The British Psychological Society issued the Power Threat Meaning Framework (Johnstone et al 2018), which presented an opportunity to approach the presentation of emotional distress in a different way. This framework suggests that an alternative approach to diagnosis and the focus on problematic behaviours is to consider what may have happened to someone, how power may have been misused in their life, and how the individual has made sense of these experiences (Johnstone et al 2018). Given that many people who receive a diagnosis of ‘personality disorder’ have experienced adversity in childhood (Porter et al 2020), this framework provides a more compassionate way of framing their difficulties. The emphasis moves from ‘what’s wrong with you?’ to ‘what has happened to you?’, providing an opportunity for insight and kindness. This is a more helpful way of engaging with the origins and development of people’s coping strategies and in understanding the effects of past experiences.

Groenewald (2018) described trauma as the ‘suffering that remains’. Trauma can be physical, psychological, emotional or sexual, can be experienced at any life stage, and is the subjective experience of an individual. Exposure to traumatic events is not unusual, with studies estimating that around 80-90% of the general population in the UK may have experienced one such event (Sweeney et al 2018, Jowett et al 2020), and the experience of childhood adversity and trauma has significant and far-reaching consequences. In the late 1990s, Felitti et al (1998) investigated the association between childhood trauma and adult health in more than 13,400 people (predominantly white, middle-class Americans), and found that childhood trauma was common in that at least half reported one or more traumatic experiences. Their findings were supported in a later survey by Public Health Wales (2015) which reported that of 2,028 adults, 47% had experienced at least one adverse childhood experience, such as neglect, physical, emotional or sexual abuse.

Studies have consistently shown that people in contact with mental health services have experienced higher rates of interpersonal violence and trauma than the general population (Khalifeh et al 2015). Mauritz et al (2013) estimated that around half of those in the mental health system had experienced physical abuse and more than one third had experienced sexual abuse in childhood or adulthood, which is significantly higher than in the general population. Childhood trauma is associated with various physical health conditions in adults, including diabetes mellitus, cancer, heart disease and obesity (Afifi et al 2016).

Despite the evidence in relation to the experience and effects of trauma, few practitioners routinely ask about it, perhaps due to concerns about retraumatising people. However, by failing to do so, they risk compromising the understanding of someone’s difficulties (Nicki 2016) and their subsequent access to appropriate interventions and support (Lietz et al 2014). A study by Purkey et al (2018), with 26 women who were frequent users of healthcare services, found that most of them had not been asked about trauma or adverse childhood experiences by healthcare professionals, and that most participants believed that their history of adverse childhood experiences was important to their health and that providers should ask about these. This was supported by Rye et al (2021), who identified that practitioners having an understanding of trauma had positive effects on their relationships with service users.

It has been suggested that BPD would be more appropriately described as complex post-traumatic stress disorder (C-PTSD) (Herman 1992, Ford and Courtois 2021). Framing their experiences in this way can be helpful for individuals because it normalises their responses to trauma, and provides a non-stigmatising explanation for both service users and practitioners in relation to the origins and development of their difficulties. According to Substance Abuse and Mental Health Administration (2014), it is helpful to consider the ‘3 Es’ of trauma – events(s), experience of event(s) and effect. Wilson et al (2013) suggested that professionals should take into account the role that trauma and prolonged traumatic stress has in the lives of service users. By doing so they can influence entire organisations and systems that then may behave differently, and provide a strengths-based ‘safe’ base, using a collaborative, trustworthy, transparent and empowering approach (Boles 2017, Rye et al 2021).

TIME OUT 2

Think about someone you know with a diagnosis of personality disorder and consider how their childhood experiences may have affected them. What sense do you think that person made of them as a child? How might these experiences relate to their presentation and difficulties?

Is your organisation understanding of an individual's experiences of trauma and is this considered in the care and treatment offered?

Use of formulation

Given the issues related to diagnosis, and the need for practitioners to understand experiences of trauma, the authors suggest that a formulation-driven approach to working with people is the most effective way of ensuring a collaborative, empathic, empowering and strengths-based approach – as advocated by trauma-focused approaches (Shaw et al 2017).

While there is no universally agreed definition of formulation, The British Psychological Society (2011) suggested that its essential features are:

- » It provides an overview of a person's presenting issues by making sense of information gathered during assessment.
- » It suggests how the person's issues may link to and maintain one another by drawing on psychological principles and theories.
- » It suggests a plan of intervention based on those psychological principles and theories.
- » It is open to revision and reformulation.
- » It is a collaborative process (Cox 2020) that can provide individuals with a means of understanding the origins of their difficulties and what maintains these (van Bilsen and Thomson 2011).

There are various models of formulation, including: generic models such as the Five-Aspects model (Greenberger and Padesky 1995); problem-specific formulations such as Clark and Wells' (1995) cognitive model of social phobia; and idiosyncratic formulations, which can be helpful if individuals present with multiple issues or with issues for which there is no validated model.

One widely used approach is the '5Ps formulation' (Box 1) (Macneil et al 2012), which provides a framework comprising five areas, each of which can be explored collaboratively with an individual.

Box 1. 5Ps formulation

Presenting problem

- » What is the problem or issues that has arisen?

Predisposing factors

- » What has happened to you?
- » Explore earlier experiences that might be affecting the person's current presentation
- » Are there any vulnerability factors you can illicit?

Precipitating factors

- » What has led to the problem or difficulty the person is experiencing on this occasion? Can you identify the trigger?

Perpetuating factors

- » What is keeping the problems going? Is there anything fuelling these problems or making them more challenging to overcome?

Protective factors

- » Are there any strengths or protective factors you can identify?
 - » Is there anything that may have helped this time or in the past?
 - » Have they felt this way before, and if so, can they recall what helped?
 - » Develop a plan to overcome or assist with the presenting problem
- (Macneil et al, 2012)

The main concept of the Power Threat Meaning Framework is that those who present with complex needs have been subjected to the misuse of power (Boyle 2022). Therefore, the authors suggest that engaging in an individualised, strengths-based approach based on a collaborative formulation is wholly appropriate. According to Katsakou et al (2012), this approach is likely to support service users with their recovery, provided it includes treatment goals that are important to them, long-term plans as to how these goals will be met, and which services may be involved.

TIME OUT 3

Think about a person you work with and start to develop a 5Ps formulation, as a way of better understanding them and their difficulties. How would you explain the term 'formulation' to them and engage them in developing this formulation further?

Evidence-based interventions

There is a growing body of evidence on effective treatments for personality disorder, particularly BPD. This treatment is usually provided in specialist mental health secondary services. National Institute for Health and Care Excellence (NICE) (2009) guidelines do not promote the use of pharmacological intervention for BPD due to a lack of evidence of its effectiveness, unless it is directed at treating comorbid mental health issues such as anxiety and depression. Despite this guidance, medicines such as anti-psychotics are often prescribed to people with personality disorder. NICE (2009) guidelines recommend psychological therapy approaches such as dialectic behavioural therapy (DBT), which is underpinned by a combination of cognitive behavioural interventions and mindfulness exercises (Linehan 1993). Stoffers et al (2012) conducted a systematic review of psychological treatments for BPD and found DBT to have the strongest evidence.

Mentalisation-based therapy is a structured therapy which has a growing evidence base. It includes both group and individual sessions underpinned by psychoanalytic approaches (Bateman and Fonagy 2012), designed to improve service users capacity to understand the intentional mental states of themselves and others– to think about thinking, which can improve emotional stability (Gardner et al, 2019). While it is becoming more frequently offered as a treatment approach across the UK, it does not yet have the depth of replication studies that are present in the DBT literature (Stoffers et al 2012).

Schema-focused therapy and Transference-focused therapy are effective interventions for BPD, however require further research and replication (Stoffers et al 2012). Schema-focused therapy is underpinned by cognitive therapy but focuses on early childhood and developmental experiences (Young et al 2003), whereas in Transference-focused therapy, which is a type of psychoanalytic psychotherapy, there is an emphasis on the relationship between therapist and service user and therapy aims to explore unhelpful interpersonal and thinking patterns. Cognitive behavioural therapy for personality disorders is also indicated as an effective treatment for borderline personality disorder and avoidant personality disorder (Davidson 2008).

DBT and mentalisation-based therapy are long term psychological interventions provided using multimodal approaches with a combination of individual and group-based interventions (Richards et al 2012). Cognitive behavioural therapy, schema-focused therapy and transference-focused therapy are usually delivered as an individual one-to-one therapy. Other newer approaches to working effectively with personality disorder include structured clinical management, in which there is an emphasis on the provision of consistency, continuity and coherence of interventions, and on problem-solving, effective crisis planning, medication review and assertive follow-up if service users disengage (Bateman and Krawitz 2013; Bateman and Fonagy, 2009). Despite this approach becoming increasingly popular within mental health services, further research of its effectiveness is required. A new approach known as structured psychological support is currently being trialled and focuses on the provision of brief low-intensity interventions in secondary mental health services (Crawford et al 2020).

At present, evidence-based psychological interventions for people with personality disorder are only available to those with the most severe presentations in specialist secondary mental health services. There is a need to develop whole systems approaches to working effectively with people with personality disorder, but research to support this and the development of such interventions are in their infancy (Lamph et al 2019, 2020) There is often a reported disparity or 'postcode lottery' of service provision for this client group, which will hopefully be addressed via changes in the healthcare system (Royal College of Psychiatrists 2020).

TIME OUT 4

Reflect on what evidence-based approaches are available in your area – are service users able to access a wide range of them? Explore what is available locally, what are the referral criteria, and what are waiting lists like? Are service users invited to identify and select therapeutic interventions?

Interpersonal effectiveness

It has been suggested that 'personality disorder' is essentially an interpersonal disorder (Wilson et al 2017, Gardner et al 2020), which reinforces the importance of developing effective therapeutic relationships. This suggests that to work with individuals

effectively, practitioners need to have high levels of self-awareness and an awareness of their own schemas and triggers, as well as those of their colleagues and team. Schemas are memories, thoughts and feelings that develop through childhood and adolescence that significantly affect an individual's reactions to internal or external triggers that relate to their early experiences (Puri et al 2021). According to several studies, people who present with BPD tend to have developed maladaptive or unhelpful schemas as a result of their early experiences (Bach and Farrell 2018, Puri et al 2021). Therefore, an awareness of one's own schemas and responses is important to prevent these from becoming a barrier to sensitive relational work. This awareness is important since there is evidence to suggest that those who present with personality difficulties may have an issue related to theory of mind, which is the ability to attribute mental states (beliefs, emotions, intent and desires) to oneself and others (Premack and Woodruff 1978).

One way of conceptualising how nurses and other healthcare professionals work with individuals is the 'boundary seesaw' (Hamilton 2010), where the ideal position would be in the centre of the seesaw, but there are times when we move up and down the seesaw, for example 'pacifying' service users, or becoming more 'controlling' rather than 'negotiating' depending on factors such as our relationships with service users, colleagues, the prevailing culture within the team, resources, and our role within the team and with the service user. The authors suggest that insight into this is essential, because without it practitioners could potentially reinforce some of the harmful messages that exist in healthcare in relation to this client group.

Such harmful messages serve to maintain unhelpful narratives, therapeutic nihilism (which is the belief that people cannot be treated effectively) and stigma in relation to people who present with personality difficulties for example that they are 'untreatable', manipulative (Nicki 2016, Rye et al 2021) and undeserving of treatment when compared with people with mental illness (Lanfredi et al 2019), meaning that people might be offered inappropriate or inadequate interventions (Paris and Black 2015), and may be excluded from services (Ferguson 2016) due to organisational anxieties about how best to manage their difficulties and support them.

This demonstrates the need for practitioners to work in a relational way with individuals, and to have an optimistic and hopeful approach that supports the notion that 'recovery' – an idea that has different meanings for different people – is possible. It has also been suggested that using the term 'recovery' in this context is problematic, because some people believe it implies there was something fundamentally 'wrong' with them. However, others view it as an open-ended journey, meaning that they learn to cope with their difficulties and live a meaningful life despite these (Katsakou et al 2012). Practices such as supervision and reflection have been shown to support practitioners who work with people who present with personality difficulties and complex needs, and can enable them to develop therapeutic, collaborative, trauma-informed and empowering relationships (Rigby and Longford 2004, Farrell et al 2010).

TIME OUT 5

Think about your team and how you initiate, maintain and develop therapeutic relationships. Where are you and your colleagues usually positioned on the boundary seesaw? Think about when and why that changes, and how you and your team could mitigate against the effect of factors that move you towards either end of the seesaw?

Conclusion

The diagnosis of personality disorder is a highly sensitive and contentious label. Those who receive this diagnosis have often endured significant trauma and adversity in early life, then are further stigmatised when given this diagnosis, with many experiencing of rejection, lack of treatment options and exclusion from services. Despite some significant improvements and training initiatives to enhance staff knowledge and challenge stigma, many people continue to report feeling misunderstood.

Mental health nurses have an important role in the care and support of people who have this diagnosis, differential diagnosis or associated issues. The authors encourage mental health nurses to engage in ongoing reflective practice and undertake any local personality disorder training initiatives to develop their knowledge, since this area of practice is rapidly evolving.

TIME OUT 6

Identify how working effectively with people who receive a diagnosis of BPD applies to your practice and the requirements of your regulatory body

TIME OUT 7

References

- Afifi TO, MacMillan HL, Boyle M et al (2016) Child abuse and physical health in adulthood. *Health Reports*. 27, 3, 10-18.
- American Psychiatric Association (2013) *Diagnostic and Statistical Manual of Mental Disorders*, Fifth Edition. APA, Arlington VA.
- American Psychiatric Association. (2022). *Diagnostic and statistical manual of mental disorders* (5th ed., text revision). American Psychiatric Association Publishing.
- Bach B, Farrell JM (2018) Schemas and modes in borderline personality disorder: the mistrustful, shameful, angry, impulsive, and unhappy child. *Psychiatry Research*. 259, 323-329. doi: 10.1016/j.psychres.2017.10.039
- Bateman, A., Fonagy, P. (2009). Randomized controlled trial of outpatient mentalization-based treatment versus structured clinical management for borderline personality disorder. *American Journal of Psychiatry*, 200 Dec; 166(12): 1355-64. doi: 10.1176/appi.ajp.2009.09040539.
- Bateman AW, Fonagy P (2012) *Handbook of Mentalizing in Mental Health Practice*. American Psychiatric Publishing, New York NY.
- Bateman AW, Krawitz R (2013) *Borderline Personality Disorder: An Evidence-Based Guide for Generalist Mental Health Professionals*. Oxford University Press, Oxford.
- Boles J (2017) Trauma-informed care: an intentional approach. *Paediatric Nursing*. 43, 5, 250-255.
- Bohus, M., Stoffers-Winterling, J., Sharp, C., Krause-Utz, A., Schmahl, C. (2021). Borderline Personality Disorder. *The Lancet*; London Vol. 398, Iss. 10310 Oct 23, 2021; 1528-1540. DOI:10.1016/S0140-6736(21)00476-1
- Boyle M (2022) Power in the power threat meaning framework. *Journal of Constructivist Psychology*. 35, 1, 27-40. doi: 10.1080/10720537.2020.1773357
- Clark DM, Wells A (1995) A cognitive model of social phobia. In Heimberg R, Leibowitz M, Hope DA et al (Eds) *Social Phobia: Diagnosis, Assessment and Treatment*. Guilford Press, New York NY, 69-93.
- Connell C, Jones E, Haslam M et al (2022) Mental health nursing identity: a critical analysis of the UK's Nursing and Midwifery Council's pre-registration syllabus change and subsequent move towards genericism. *Mental Health Review Journal*. 27, 4, 472-483. doi: 10.1108/MHRJ-02-2022-0012
- Cox LA (2020) Use of individual formulation in mental health practice. *Mental Health Practice*. doi: 10.7748/mhp.2020.e1515
- Crawford MJ, Thana L, Parker J et al (2020) Structured Psychological Support for people with personality disorder: feasibility randomised controlled trial of a low-intensity intervention. *BJPsych Open*. 6, 2, e25. doi: 10.1192/bjo.2020.7
- Davidson KM (2008) Cognitive-behavioural therapy for personality disorders. *Psychiatry*. 7, 3, 117-120. doi: 10.1016/j.mppsy.2008.01.005
- Farrell GA, Shafiei T, Salmon P (2010) Facing up to 'challenging behaviour': a model for training in staff-client interaction. *JAN*. 66, 7, 1644-1655. doi: 10.1111/j.1365-2648.2010.05340.x
- Felitti VJ, Anda RF, Nordenberg MD et al (1998) Relationship of childhood abuse and household dysfunction to many of the leading causes of death in adults. The Adverse Childhood Experiences (ACE) Study. *American Journal of Preventative Medicine*. 14, 4, 245-258. doi: 10.1016/S0749-3797(98)00017-8
- Ferguson A (2016) Borderline personality disorder and access to services: a crucial social justice issue. *Australian Social Work*. 69, 2, 206-214. doi: 10.1080/0312407X.2015.1054296
- Ford JD, Courtois CA (2021) Complex PTSD and borderline personality disorder. *Borderline Personality Disorder and Emotion Dysregulation*. 8, 1, 16. doi: 10.1186/s40479-021-00155-9
- Gardner KJ, Wright KM, Elliott A et al (2020) The weirdness of having a bunch of other minds like yours in the room: the lived experiences of mentalization-based therapy for borderline personality disorder. *Psychology and Psychotherapy*. 93, 3, 572-586. doi: 10.1111/papt.12243
- Greenberger D, Padesky CA (1995) *Mind Over Mood: A Cognitive Therapy Treatment Manual for Clients*. Guilford Press, New York NY.
- Groenewald A (2018) 'Trauma is suffering that remains'. The contribution of trauma studies to prophetic studies. *Acta Theologica*. 38, Suppl 26, 88-102. doi: 10.18820/23099089/actat.Sup26.5
- Hamilton L (2010) The boundary seesaw model: good fences make for good neighbours. In Tennant A, Howells K (Eds) *Using Time, Not Doing Time: Practitioner Perspectives on Personality Disorder and Risk*. Wiley-Blackwell, Oxford, 181-194.
- Hayward M, Moran P (2007) Comorbidity of personality disorder and mental illness. *Psychiatry*. 7, 3, 102-104. doi: 10.1016/j.mppsy.2008.01.010

- Herman J (1992) *Trauma and Recovery: The Aftermath of Violence – From Domestic Abuse to Political Terror*. Basic Books, New York NY.
- Johnstone L, Boyle M, Cromby J et al (2018) The Power Threat Meaning Framework: Towards the Identification of Patterns in Emotional Distress, Unusual Experiences and Troubled or Troubling Behavior, As an Alternative to Functional Psychiatric Diagnosis. The British Psychological Society, Leicester.
- Jowett S, Karatzias T, Albert I (2020) Multiple and interpersonal trauma are risk factors for both post-traumatic stress disorder and borderline personality disorder: a systematic review on the traumatic backgrounds and clinical characteristics of comorbid post-traumatic stress disorder/borderline personality disorder groups versus single-disorder groups. *Psychology and Psychotherapy*. 93, 3, 621-638. doi: 10.1111/papt.12248
- Katsakou C, Marougka S, Barnicot K et al (2012) Recovery in borderline personality disorder (BPD): a qualitative study of service users' perspectives. *PLoS One*. 7, 5, e36517. doi: 10.1371/journal.pone.0036517
- Khalifeh H, Moran P, Borschmann R et al (2015) Domestic and sexual violence against patients with severe mental illness. *Psychological Medicine*. 45, 4, 875-886. doi: 10.1017/S0033291714001962
- Lamph G, Baker J, Dickinson T et al (2019) Personality disorder co-morbidity in primary care 'Improving Access to Psychological Therapy' (IAPT) services: a qualitative study exploring professionals' perspectives of working with this patient group. *Personality and Mental Health Journal*. 13, 3, 168-179. doi: 10.1002/pmh.1454
- Lamph G, Baker J, Dickinson T et al (2020) Personality disorder co-morbidity in primary care 'Improving Access to Psychological Therapy' (IAPT) services: a qualitative study exploring patient perspectives on treatment experience. *Behavioural and Cognitive Psychotherapy*. 49, 2, 144-158. doi: 10.1017/S1352465820000594
- Lamph G, Dorothy J, Jaynes T et al (2021) A qualitative study of the label of personality disorder from the perspectives of people with lived experience and occupational experience. *Mental Health Review Journal*. 27, 1, 31-47. doi: 10.1108/MHRJ-05-2020-0035
- Lanfredi M, Ridolfi ME, Occhialini G et al (2019) Attitudes of mental health staff toward patients with borderline personality disorder: an Italian cross-sectional multisite study. *Journal of Personality Disorders*. 1-16. doi: 10.1521/pedi_2019_33_421
- Lieb K, Zanarini MC, Schmahl C et al (2004) Borderline personality disorder. *The Lancet*. 364, 9432, 453-461. doi: 10.1016/S0140-6736(04)16770-6
- Lietz CA, Lacasse JR, Cheung JR (2014) A case study approach to mental health recovery: understanding the importance of trauma-informed care. *Ethical Human Psychology and Psychiatry*. 16, 3, 167-182. doi: 10.1891/1559-4343.16.3.167
- Linehan MM (1993) *Skills Training Manual for Treating Borderline Personality Disorder*. Guilford Press, London.
- Macneil CA, Hasty MK, Conus P et al (2012) Is diagnosis enough to guide interventions in mental health? Using case formulation in clinical practice. *BMC Medicine*. 10, 111. doi: 10.1186/1741-7015-10-111
- Mauritz MW, Goossens PJ, Draijer N et al (2013) Prevalence of interpersonal trauma exposure and trauma-related disorders in severe mental illness. *European Journal of Psychotraumatology*. 4. doi: 10.3402/ejpt.v4i0.19985
- Mulder RT (2021) ICD-11 personality disorders: utility and implications of the new model. *Frontiers in Psychiatry*. 12, 655548. doi: 10.3389/fpsyt.2021.655548
- National Institute for Health and Care Excellence (2009) *Borderline Personality Disorder: Recognition and Management*. Clinical guideline No. 78. NICE, London.
- Nicki A (2016) Borderline personality disorder, discrimination, and survivors of chronic childhood trauma. *International Journal of Feminist Approaches to Bioethics*. 9, 1, 218-245. doi: 10.3138/ijfab.9.1.218
- Paris J, Black DW (2015) Borderline personality disorder and bipolar disorder. what is the difference and why does it matter? *Journal of Nervous and Mental Disease*. 23, 1, 3-7. doi: 10.1097/NMD.0000000000000225
- Porter C, Palmier-Claus J, Branitsky A et al (2020) Childhood adversity and borderline personality disorder: a meta-analysis. *Acta Psychiatrica Scandinavica*. 141, 1, 6-20. doi: 10.1111/acps.13118
- Premack D, Woodruff G (1978) Does the chimpanzee have a theory of mind? *Behavioral and Brain Sciences*. 1, 4, 515-526. doi: 10.1017/S0140525X00076512
- Public Health Wales (2015). *Adverse Childhood Experiences and their association with health-harming behaviours in the Welsh adult population*. The Policy, Research and International Development Directorate, Public Health Wales NHS Trust.
- Purkey E, Patel R, Beckett T et al (2018) Primary care experiences of women with a history of childhood trauma and chronic disease: trauma-informed care approach. *Canadian Family Physician*. 64, 3, 204-211.
- Puri P, Kumar D, Muralidharan K et al (2021) Evaluating schema modes and cognitive distortions in borderline personality disorder: a mixed-method approach. *Journal of Clinical Psychology*. 77, 9, 1973-1984. doi: 10.1002/jclp.23126
- Read J, Harper DJ (2022) The power threat meaning framework: addressing adversity, challenging prejudice and stigma, and transforming services. *Journal of Constructivist Psychology*. 35, 1, 54-67. doi: 10.1080/10720537.2020.1773356
- Richards DA, Bower P, Pagel C et al (2012) Delivering stepped care: an analysis of implementation in routine practice. *Implementation Science*. 7, 3. doi: 10.1186/1748-5908-7-3

- Rigby M, Longford J (2004) Development of a multi-agency experiential training course on personality disorder. *Psychiatric Bulletin*. 28, 9, 337-341. doi: 10.1192/pb.28.9.337
- Royal College of Psychiatrists (2020) PS01/20: Services for People Diagnosable with Personality Disorder. Position Statement. www.rcpsych.ac.uk/docs/default-source/improving-care/better-mh-policy/position-statements/ps01_20.pdf?sfvrsn=85af7fbc_2 (Last accessed: 23 January 2023.)
- Rye E, Anderson J, Pickard M (2021) Developing trauma-informed care: using psychodynamic concepts to help staff respond to the attachment needs of survivors of trauma. *Advances in Mental Health and Intellectual Disabilities*. 15, 5, 201-208. doi: 10.1108/AMHID-12-2020-0033
- Sampson MJ, McCubbin RA, Tyrer P (2006) *Personality Disorder and Community Mental Health Teams: A Practitioners Guide*. John Wiley & Sons, Chichester.
- Shaw J, Higgins C, Quartey C (2017) The impact of collaborative case formulation with high risk offenders with personality disorder. *Journal of Forensic Psychiatry & Psychology*. 28, 6, 777-789. doi: 10.1080/14789949.2017.1324579
- Stoffers JM, Vollm BA, Rucker G et al (2012) Psychological therapies for people with borderline personality disorder. *Cochrane Database of Systematic Reviews*. Issue 5. CD012955. doi: 10.1002/14651858.CD012955.pub2
- Substance Abuse and Mental Health Administration (2014) *SAMHSA's Concept of Trauma and Guidance for a Trauma-Informed Approach*. SAMHSA, Rockville MD.
- Sweeney A, Filson B, Kennedy A et al (2018) A paradigm shift: relationships in trauma-informed mental health services. *BJPsych Advances*. 24, 5, 319-333. doi: 10.1192/bja.2018.29
- The British Psychological Society (2011) *Good Practice Guidelines on the Use of Psychological Formulation*. BPS, Leicester.
- van Bilsen H, Thomson B (2011) *CBT for Personality Disorders*. SAGE Publications, London.
- Wilson C, Pence DM, Conradi L (2013) Trauma-informed care. *Encyclopaedia of Social Work*. 1-15. doi: 10.1093/acrefore/9780199975839.013.1063
- Wilson S, Stroud CB, Durbin CE (2017) Interpersonal dysfunction in personality disorders: a meta-analytic review. *Psychological Bulletin*. 143, 7, 677-734. doi: 10.1037/bul0000101
- Winsper, C., Bilgin, A., Thompson, A., Marwaha, S., Chanen, A. M. (2020). The prevalence of personality disorders in the community: a global systematic review and meta-analysis. *British Journal of Psychiatry*; London Vol. 216, Iss. 2. (Feb 2020): 69-78. DOI:10.1192/bjp.2019.166
- World Health Organisation (2019). *International Classification of Diseases, Eleventh Revision (ICD-11)*, World Health Organization (WHO) 2019/2021 <https://icd.who.int/browse11>.
- Young JE, Klosko J, Weishaar ME (2003) *Schema Therapy: A Practitioner's Guide*. Guilford, New York NY.