

## Central Lancashire Online Knowledge (CLOK)

Title	The impact of injury on match running performance following the return to competitive match-play over two consecutive seasons in elite European soccer players
Type	Article
URL	<a href="https://clock.uclan.ac.uk/47762/">https://clock.uclan.ac.uk/47762/</a>
DOI	<a href="https://doi.org/10.7752/jpes.2023.05142">https://doi.org/10.7752/jpes.2023.05142</a>
Date	2023
Citation	Morgans, Ryland, Rhodes, David, Bezuglov, Eduard, Etemad, O, Di Michele, Rocco, Teixeira, José, Modric, Toni, Versic, S. and Oliveira, Rafael (2023) The impact of injury on match running performance following the return to competitive match-play over two consecutive seasons in elite European soccer players. <i>Journal of Physical Education and Sport</i> , 23 (5). pp. 1142-1149. ISSN 2247-8051
Creators	Morgans, Ryland, Rhodes, David, Bezuglov, Eduard, Etemad, O, Di Michele, Rocco, Teixeira, José, Modric, Toni, Versic, S. and Oliveira, Rafael

It is advisable to refer to the publisher's version if you intend to cite from the work.  
<https://doi.org/10.7752/jpes.2023.05142>

For information about Research at UCLan please go to <http://www.uclan.ac.uk/research/>

All outputs in CLOK are protected by Intellectual Property Rights law, including Copyright law. Copyright, IPR and Moral Rights for the works on this site are retained by the individual authors and/or other copyright owners. Terms and conditions for use of this material are defined in the <http://clock.uclan.ac.uk/policies/>

## The impact of injury on match running performance following the return to competitive match-play over two consecutive seasons in elite European soccer players

MORGANS R<sup>1</sup>, RHODES D<sup>2</sup>, BEZUGLOV E<sup>3</sup>, ETEMAD O<sup>4</sup>, DI MICHELE R<sup>5</sup>, TEIXEIRA J<sup>6</sup>, MODRIC T<sup>7</sup>,  
VERVIC S<sup>8</sup>, OLIVEIRA R<sup>9</sup>

<sup>1,2</sup>Football Performance Hub, University of Central Lancashire, Preston PR1 2HE, UK

<sup>1,3</sup>Department of Sports Medicine and Medical Rehabilitation, Sechenov First State Medical University, Moscow, RUSSIA FEDERATION

<sup>4</sup>Medical Department, CSKA PFC, Moscow, RUSSIA FEDERATION

<sup>5</sup>Department of Biomedical and Neuromotor Sciences, University of Bologna, ITALY

<sup>6,9</sup>Research Centre in Sports Sciences, Health and Human Development, 5001-801 Vila Real, PORTUGAL

<sup>6</sup>Department of Sport and Physical Education, Polytechnic Institute of Bragança, 5300-253 Bragança, PORTUGAL

<sup>6</sup>Polytechnic Institute of Guarda, 6300-559 Guarda, PORTUGAL

<sup>7,8</sup>Faculty of Kinesiology, University of Split, Split, CROATIA

<sup>7,8</sup>High Performance Sport Center, Croatian Olympic Committee, Zagreb, CROATIA

<sup>9</sup>Sports Science School of Rio Maior-Polytechnic Institute of Santarém, 2040-413 Rio Maior, PORTUGAL

<sup>9</sup>Life Quality Research Centre, 2040-413 Rio Maior, PORTUGAL

Published online: May 31, 2023

(Accepted for publication May 15, 2023)

DOI:10.7752/jpes.2023.05142

### Abstract

Based on the assessment and diagnosis, the rest period following a moderate/severe injury may lead to de-conditioning for the injured player and therefore an association with a prolonged rehabilitation, re-conditioning and return to sport is observed post-injury. The aim of the present study was to assess the impact of all injuries on match running performance following the return to competitive match-play over two consecutive seasons in elite European soccer players. A retrospective analysis was conducted utilizing data related to a player's injury and match running performance. A club physiotherapist consistently recorded availability and injury data in a standardized format. Linear mixed modelling analysis revealed no difference between PRE and POST1, POST2, and POST3 for total distance, running distance, high-intensity distance, and sprint distance (all  $p > 0.05$ ). Although, maximum speed was significantly ( $p < 0.05$ ) lower in POST1 and POST2 when compared to PRE, in both cases with a large ( $ES = 1.88$ ) effect. No significant difference was observed for maximum speed between PRE and POST3 ( $p = 0.07$ ). There were very low correlations between the number of days absent and changes in maximum speed between POST1 and PRE ( $r = 0.09$ , 95% CI -0.42 to 0.56), and POST2 and PRE ( $r = 0.10$ , 95% CI -0.42 to 0.57), respectively. In conclusion, no variation in distance variables were found regardless of one, two or three matches post-injury compared to pre-injury status. Moreover, maximum speed was lower during the first three matches post-injury, although the mean value was slightly lower. Finally, a low correlation between absent days and maximum speed loss between pre-injury and following one and two matches were found.

**Key Words:** soccer; injury analysis; running performance; match data; sprint; maximal speed

### Introduction

The literature examining injuries has highlighted a high injury incidence in elite teams (Ekstrand et al., 2011a, 2011b). Injuries are an inevitable drawback in any sport and are debilitating and costly (Häggglund et al., 2013a, 2013b; Knowles et al., 2007). Specifically, muscle injuries in professional soccer are a major concern to teams as a negative impact on performance through lower player availability and club economy as well as the potential for increased risk of subsequent injury can occur (Fanchini et al., 2020). A professional team with a squad of 25 players can expect approximately 16 time-loss muscle injuries per season with most (92%) affecting the hamstring, quadriceps, adductor, and calf muscle groups (Ekstrand et al., 2016). Furthermore, the average number of muscle injuries in men's professional soccer can be categorized as six hamstring, three quadriceps, three adductor, one/two calf, and one/two 'other' muscle groups (McCall et al., 2015). Thus, the prevention of injuries is an essential aspect in sport management to optimize team performance and resources (Seow and Massey, 2022). Furthermore, based on the assessment and diagnosis, the rest period following a moderate/severe injury may lead to de-conditioning for the injured player and therefore an association with a prolonged

rehabilitation, re-conditioning and return to sport (RTS) is observed post-injury (Brukner et al., 2018; Pollock et al., 2016; Taberner et al., 2022).

A further practical problem may be that players' RTS may occur prior to fully completing the re-conditioning stage of the rehabilitation process due to internal pressure from the manager/coaching staff and external pressure, namely match outcome and the media. Thus, minor to mild injuries normally involve only a few absent days, and minimal de-conditioning, and may be underestimated. As a result, the player may be required to compete under sub-optimal physical (or even psychological) condition, which may lead to reduced match performance and further injury risk. Therefore, it is necessary to understand if and to what extent match running performance is altered in the first matches following RTS compared to pre-injury performance.

A number of recent studies have assessed the post- vs. pre-injury match performance in soccer players of different age groups and categories, in particular high-intensity running and sprint performance, although the vast majority of this research has only focused on hamstring strains (Hoppen et al., 2022; Jimenez et al., 2020; Whiteley et al., 2021; Whiteley et al., 2022). Only one study (Raya-Gonzalez et al., 2022) examined the impact of all injury types in elite soccer (Spain La Liga, across two seasons), where injuries were identified according to the UEFA guidelines for epidemiological studies as events that caused absence from the next training session or match (Hagglund et al., 2005). Although, Raya-Gonzalez et al. (2022) did not collect and analyse data from official club medical staff, instead publicly available data from the clubs' website was examined. Thus, the validity and reliability of the data may be questionable, as the authors analyzed match performance changes following players sustaining an injury. The main findings of this study showed reductions in playing time following moderate and major injuries. Moreover, reductions post-minor injury were observed during jogging (>6 km/h) and running (6 to 12 km/h). Additionally, greater distances were covered when running at 18-21 km/h and 21-24 km/h by 161 professional Spanish soccer players from La Liga who had suffered major injuries. Lastly, maximum speed also decreased following moderate and major injuries. Overall, these studies highlighted reductions in high-intensity and sprint distance and maximum speed in the matches played immediately following RTS (Hoppen et al., 2022; Jimenez et al., 2020; Raya-Gonzalez et al., 2022; Whiteley et al., 2021; Whiteley et al., 2022). However, the results of these studies are not consistent due to the specific nature of the study and the effects of re-conditioning training. Despite current RTS guidelines in the existing literature regarding timelines following injury in elite soccer (Taberner et al., 2022), limited information is available regarding match running performance post-injury.

Alongside physical performance, injury incidence and subsequent player availability has also been commonly cited as a key factor in the success of teams over the course of a soccer season. Carling et al. (2015) investigated the relationship between injury, player availability and team performance across five consecutive French Ligue 1 seasons. The authors found that during the season with the highest player availability, the team was able to attain the highest points total and conceded the least number of goals across the five examined seasons. While there may be many other contextual factors that contribute towards these achievements, this may suggest that higher player availability may play an important role in the overall success of a soccer team over the course of a season.

Considering the previous literature and the fact that only one study was found to analyze the injury impact on match performance variables (Raya-Gonzalez et al., 2022), the aim of the present study was to assess the impact of any injury, identified according to the UEFA definition (an event involving the absence from subsequent training/match), on match running performance following RTS over two consecutive seasons in elite European soccer players. The study hypothesis was that the first few matches post-injury will result in lower running demands when compared with pre-injury status as suggested by Raya-González et al. (2022).

## Material and methods

A retrospective analysis was conducted utilizing player injury and match running performance data collected in an elite European Premier League soccer team. A club physiotherapist consistently recorded availability and injury data on a daily basis in a standardized format for each player. The consensus on definitions and data collection procedures in studies of soccer injuries (Fuller et al., 2006) only administered and recorded time loss injuries. Players that were previously injured at study commencement were included, although were not incorporated into the analysis, while new players recruited during the study period were included. The study was conducted according to the requirements of the Declaration of Helsinki and was approved by the local Ethics Committee of the University of Central Lancashire (N 0104 dated 7/12/20) and the Premiership club from which the subjects volunteered (Winter & Maughan, 2009). To ensure confidentiality, all data were anonymized prior to analysis. All players provided consent for their injury-related and match running performance data to be collected and utilized for research purposes.

### Participants

Eleven professional outfield soccer players from a European Premiership club were involved in the study. Data from the complete 2019-20 and 2020-21 seasons included 11 senior players (age  $23.6 \pm 6.3$  years, weight  $76.1 \pm 7.8$  kg, height  $1.78 \pm 0.08$  m). The inclusion criteria for the study were: (i) listed on the first-team roster of the Premiership club at the start of the 2019-20 and/or 2020-21 seasons, (ii) regularly trained with the

first-team, (iii) participated in at least 80% of training sessions and matches, (iv) did not use dietary supplements during the study, (v) not injured during the study, and (vi) did not participate in any other training program during the study period. Additionally, the exclusion criteria for the study were: (i) long-term injury, (ii) joining the team late in either of the study seasons, (iii) lack of full, complete data for training and/or match-play, (iv) an insufficient number of satellite connection signals, and (v) goalkeepers, due to significant variations in physical demands compared with outfield players. Players were assigned to one of five positions as match demands for these differ considerably. The methodology of differentiating specialized positions was adapted from previous research (Aalbers et al., 2019). Goalkeepers were excluded from the investigation due to the specific nature of their match activity and low running demands (Ingebrigtsen et al., 2015; Bradley et al., 2009).

All injuries (n = 85, including minimal to severe injuries) recorded in rostered outfield players (n = 31) from the study team across the 2019-20 and 2020-21 seasons were initially considered. Subsequently, only players (n = 22) having completed at least 10 matches and for a minimum of 30-minutes were included, thus recording a total of 59 injuries. Finally, only injuries where at least 3 matches were completed and for a minimum of 30-minutes in the month preceding the injury, and 3 matches completed for a minimum of 30-minutes in the month following the last day-off due to injury were examined. Therefore, a total of 16 injuries from 11 players were considered for analysis. Examined injuries included muscle (n = 7), ligament/tendon (n = 4), contusions (n = 3) and concussions/effusions (n = 2). The mean and standard deviation (SD) of number of absent days was  $12 \pm 15$ , with a range of 1 to 49.

#### *Data collection*

Each season was split into two phases to mirror the official fixture schedule. The first phase was sub-divided into pre-season (mid-June to mid-July) that lasted 5-weeks and the competitive season (mid-July to mid-December) that lasted 25-weeks. The second phase was also sub-divided into pre-season (mid-January to end-February) that lasted 6-weeks and the competitive season (end-February to mid-May) that lasted 11-weeks.

All injuries were assessed, diagnosed and categorized by the club's physiotherapist and head doctor and defined as injury incidence resulting in a modified training program, or missed training sessions or matches (Rogalski et al., 2012). Injuries were further categorized by body-site (injury location and region), injury type (trauma, overuse), sub-type (fracture, other bone injury, dislocation/subluxation, sprain/ligament, meniscus/cartilage, concussion, muscle strain, tendon, hematoma/contusion, nerve injury and synovitis), nature (training, match or undefined), description and days missed. The definition of injury severity differed slightly to previously reported methods (Rogalski et al., 2012), although are consistent with Fuller et al. (2006) where injury severity was defined as the number of days from the date of injury to the date of return to full participation in team training and availability for match selection (Fuller et al., 2006).

A two-camera optical tracking system (InStat, Moscow, Russia) was utilized to record and examine the match running performance across two consecutive seasons. The matches were filmed using two full HD, static cameras positioned on the centre line of the field, not less than 3-metres from the field and 7-metres in height. A consistent 25Hz format was provided. Data were linearly interpolated to 50Hz, smoothed using a 5-point moving average and then down-sampled to 10Hz, which allowed analysis of all player actions with and without the ball (Morgans et al., 2022). The installation process, reliability and validity of InStat have been previously reported (FIFA, 2019; Morgans et al., 2022). Physical performance was analyzed using the InStat Analysis Software System. Match running performance metrics included: time on pitch (mins); total distance covered (m); high-intensity distance (m; total distance covered 5.5-7m/s); sprint distance (m; total distance covered >7m/s); explosive distance (m; distance covered with an acceleration  $>1.12\text{m.s}^{-2}$ ); acceleration distance (m; distance covered during accelerations  $+3\text{m.s}^{-2}$ ); deceleration distance (m; distance covered during decelerations  $+3\text{m.s}^{-2}$ ) and lasted more than 0.5-sec (Algroy et al., 2021). All distance variables were divided by individual playing time (expressed in minutes) to calculate a distance per minute value.

#### *Statistical analysis*

All data are presented as mean and 95% confidence interval (CI), or as median and interquartile range (IQR). Physical performance data were averaged across the last three matches completed by a player prior to injury occurrence, to obtain a pre-injury match performance level (PRE). Linear mixed models with random intercepts on individual players's IDs were used to examine, for every match performance variable, the differences between PRE and respectively the first (POST-1), second (POST-2), and third (POST-3) match performed following RTS. Differences were standardized by the estimated between-subject SD to calculate an effect size (ES), that was evaluated as trivial ( $<0.2$ ), small (0.2-0.6), moderate (0.6-1.2), large (1.2-2.0), very large (2.0-4.0) and extremely large ( $>4.0$ ) (Hopkins et al., 2009). When a significant difference between PRE and any post-injury match was observed, Pearson's r correlation coefficients calculated between individual changes in that performance variable (POST minus PRE) and the number of absent days due to injury, to determine the relationship between such variables. Furthermore, for each variable, the number of matches played from RTS to achieving or overcoming the pre-injury performance values were calculated (that is 1, 2, 3 or >3),

and were expressed as the median and IQR. The analysis were conducted with R software, version 4.2.0. For all analyzes, statistical significance was set at  $p < 0.05$ .

## Results

Table 1 shows the estimated means and 95% CIs for the examined match running performance variables before and following injury, while Table 2 displays the estimated differences with 95% CIs and effect sizes. The linear mixed modelling analysis revealed no difference between PRE and POST1, POST2, and POST3 for total distance, running distance, high-intensity distance, and sprint distance (all  $p > 0.05$ ). Although, maximum speed was significantly ( $p < 0.05$ ) lower in POST1 when compared to PRE, and in POST2 when compared to PRE, in both cases with a large ( $ES = 1.88$ ) effect. No significant difference was observed for maximum speed between PRE and POST3 ( $p = 0.07$ ).

Table 1. Estimated means and 95% CIs for the examined running performance variables in matches preceding an injury (PRE) and in the first (POST1), second (POST2), and third (POST3) match following RTS.

	PRE	POST1	POST2	POST3
Total distance (m/min)	118 (113 - 122)	118 (114 - 122)	119 (115 - 123)	121 (116 - 125)
Running distance (m/min)	20.5 (18.2 - 22.7)	21.0 (18.8 - 23.3)	21.5 (19.2 - 23.7)	21.7 (19.4 - 23.9)
High-intensity distance (m/min)	8.4 (7.0 - 10.0)	8.6 (7.0 - 10.0)	8.3 (6.8 - 9.8)	9.0 (7.6 - 10.5)
Sprint distance (m/min)	1.6 (1.2 - 2.1)	1.4 (0.9 - 1.8)	1.3 (0.8 - 1.7)	1.3 (0.8 - 1.7)
Maximum speed (m/s)	8.3 (8.1 - 8.5)	8.0 (7.8 - 8.2)	8.1 (7.9 - 8.3)	8.1 (7.9 - 8.3)

Table 2. Estimated differences and effect sizes for POST1, POST2 and POST3 vs. PRE.

	POST1	POST2	POST3
Total distance (m/min)	0.5; ES = 0.11	1.5; ES = 0.33	3.0; ES = 0.67
Running distance (m/min)	0.6; ES = 0.21	1.0; ES = 0.36	1.2; ES = 0.43
High-intensity distance (m/min)	0.1; ES = 0.06	-0.1; ES = -0.06	0.6; ES = 0.33
Sprint distance (m/min)	-0.3; ES = -0.75	-0.3; ES = -0.75	-0.3; ES = -0.75
Maximum speed (m/s)	-0.3*; ES = -1.88	-0.3*; ES = -1.88	-0.2; ES = -1.25

\* $p < 0.05$

A very low correlation between number of absent days and changes in maximum speed between POST1 and PRE ( $r = 0.09$ , 95% CI -0.42 to 0.56), and POST2 and PRE ( $r = 0.10$ , 95% CI -0.42 to 0.57) were observed respectively.




Figure 1. The relationship between the number of days absent and change in maximum speed between PRE and POST1




Figure 2. The relationship between the number of days absent and change in maximum speed between PRE and POST2

The median (IQR) for the number of matches to attain pre-injury running performance was 1 (1 to 2) for total distance; 2 (1 to 2.25) for running distance; 1.5 (1 to 3) for high-intensity distance; 2 (1 to 4) for sprint distance; and 3 (1.75 to 4.00) for maximum speed, respectively.

## Discussion

The aim of the present study was to assess the impact of all injury types on match running performance following RTS competitive match-play across two consecutive seasons in elite European soccer players. The main findings showed no substantial variations for distance variables regardless of one to three matches post-injury compared to pre-injury status. Only maximum speed was found to be lower in POST1 and POST2, although the mean value was slightly lower. Moreover, no correlations between absent days and maximum speed reductions in POST1 and POST2 were found, which contrasts the Raya-Gonzalez et al. (2022) study, who only highlighted a difference for moderate and severe injuries. Such results did not confirm the hypothesis of the present study.

Regarding the non-variations for running distance variables, specifically for total distance and running distance, the results supported the Raya-Gonzalez et al. (2022) findings. The present study reinforced Raya-Gonzalez et al. (2022) results regardless of injury severity (minor - from 4 to 7 missed days, moderate - from 8 to 28 missed days, and major - more than 28 missed days). However, injury severity was not analyzed in the present study due to the limited number of each injury type. Even so, the present data contrasts Raya-Gonzalez et al. (2022) findings who reported a reduction in playing time experienced by all players who suffered injury, while the present study analyzed mainly starting players.

Previous research has suggested that players suffering injury severity of four to seven days may have participated in official match-play prior to full recovery in terms of health readiness (Wing, 2018). The present study, also reported more injuries of similar severity, noted no differences in high-intensity and sprint distance. Such results may be relevant to clarify each study and thus the need to analyse players specific context. There are other possible justifications for the non-variation results in running distance variables. For instance, a previous study reported that following a major injury, players performed high-intensity running in the fourth match post-injury (Jiménez-Rubio et al., 2020). This may be related to specific high-speed running (Taberner et al., 2019), along with greater control of the training load and fatigue during the re-conditioning program (Jiménez-Rubio et al., 2020). In contrast, it was previously suggested that players that suffered minor injury, may be included as a starting player even prior to full recovery, which may decrease match performance (Raya-Gonzalez et al., 2022). However, minor injuries were most prevalent in the present study, although such results were not confirmed.

Furthermore, maximum speed remained similar pre- and post-injury which opposes the Raya-Gonzalez et al. (2022) study who found increases post-minor injury, and no changes following moderate and major injuries. Again, this does not seem to be confirmed in the present study considering that the higher number of injuries remained in the minor category. On the one hand, Raya-Gonzalez et al. (2022) attributed such findings

to maximum speed that decreased following moderate and major injuries. Such a scenario was similar in the present study, since maximum speed showed to be lower in POST1 and POST2, although the mean value was slightly lower. On the other hand, Jiménez-Rubio et al. (2020) study found significant improvements in maximum speed. Nonetheless, there are methodological differences between the studies since in the Jiménez-Rubio et al. (2020) study, all players completed the same re-conditioning program following a grade IIb hamstring strain, while the present study included distinct types of injuries and different types of re-conditioning programs. Therefore, the current findings suggest that more than three official matches are required to fully recover to pre-injury running performance levels. This may also contribute towards an impaired maximum speed performance during match-play, that requires careful consideration for coaches given the importance of this variable (Faude et al., 2012).

Additionally, there was a low correlation between absent days and maximum speed loss. This suggests that players tended to reduce maximum speed following any injury, irrespective of the number of days absent. Even following only one or two absent days, players maximum speed attained were slower in the first two matches completed. Some speculations for this finding may be associated with psychological mechanisms. For instance, fear of worsening the injury if not 100% confident that the injury has fully healed despite being allowed to play, factors that were not controlled in the present study.

Despite the interesting results of this study, there are some limitations to address. Firstly, the limited number of players and injuries was due to only including injuries with at least four matches played pre-injury and at least four matches played post-RTS. Thus, it is suggested that future research should assess longer periods of time with potentially a greater number of participants and injury and running performance data. Secondly, different injury types were examined collectively (i.e., sprains, strains, traumatic, overuse, etc.). This may also be regarded as a study strength, considering that maximum speed was reduced regardless of injury severity classification. Nonetheless, a wider injury sample may be differentiated according to type and varying rehabilitation training for future research. Thirdly, only external load data was analyzed while no different biomechanical or physiological tests were employed to compare pre- and post-injury status as suggested previously (Jiménez-Rubio et al., 2020; Raya-Gonzalez et al., 2022). Fourthly, injured players were not compared with a control group of non-injured players as well as contextual factors such as match result and quality of opposition (Gonçalves et al., 2020; Gonçalves et al., 2019; Rago et al., 2021). These factors may have an impact on running distances and/or maximum speed that may provide greater insights for the present research. Finally, the differences in positional demands were not considered, as players with various tactical roles may perform 7 to 61 sprints (Taylor et al., 2017), as this was not possible due to the small sample size.

In comparison to Raya-Gonzalez et al. (2022) that was the only previous study found relating to this topic in elite soccer players, this was the first analysis conducted on an elite soccer team from a European league utilizing injury data collected from the club medical staff. From a methodological point of view, the number of days absent for each injury was considered rather than classifying injuries into discrete categories (based on absent days). Furthermore, it is also relevant to highlight that four injury types were included in the present study which were not analyzed separately due to the limited number of incidents. This may also be a limitation, although this is the real-world scenario of any professional team and should be considered.

## Conclusions

In conclusion, no variation in the examined distance metrics were found regardless of one to three matches post-injury when compared to pre-injury status. Moreover, it was reported that maximum speed was lower during the first three matches post-injury, although the mean value was slightly lower. Finally, a low correlation between absent days and maximum speed loss between pre-injury and following one and two matches, respectively were found. The practical implications of this study for coaches and performance staff are that the number of absent days for each injury is significant in order to individualize the players profile in relation to the positional demands within the unit and team. Such information is relevant to understand when players are ready to RTS competitive match-play based on key performance markers and the comparison to pre-injury levels.

## Acknowledgements

The authors would like to express their gratitude to the club that participated in the study.

## Conflict of interest

The author(s) declare no potential conflict of interest with respect to the research, authorship, and/or publication of this article.

## Funding

This research was funded by the Portuguese Foundation for Science and Technology, I.P., Grant/Award Number UIDP/04748/2020. The funders had no role in the design of the study; in the collection, analyses, or interpretation of data; in the writing of the manuscript, or in the decision to publish the results.



## References

- Aalbers, B., Van Haaren, J., editors. (2019). Distinguishing between roles of football players in play-by-play match event data. International Workshop on Machine Learning and Data Mining for Sports Analytics; 2019: Springer.
- Algroy, E., Grendstad, H., Riiser, A., Nybakken, T., Saeterbakken, A. H., Vidar Andersen, V., & Gundersen, H. S. (2021). Motion Analysis of MatchPlay in U14 Male Soccer Players and the Influence of Position, Competitive Level and Contextual Variables. *International Journal of Environmental Research and Public Health*, 18, 7287.
- Bradley, P. S., Sheldon, W., Wooster, B., Olsen, P., Boanas, P., & Krstrup, P. (2009). High-intensity running in English FA Premier League soccer matches. *Journal of Sports Science*, 27, 159–168.
- Brukner, P., Cook, J. L., & Purdam, C. R. (2018). Does the intramuscular tendon act like a free tendon? *British Journal of Sports Medicine*, 52(19), 1227e1228.
- Carling, C., Le Gall, F., McCall, A., Nedelec, M., & Dupont, G. (2015). Squad management, injury and match performance in a professional soccer team over a championship-winning season. *European Journal of Sport Science*, 15 (7), 573-582.
- Ekstrand, J., Hagglund, M., & Walden, M. (2011a). Injury incidence and injury patterns in professional soccer: The UEFA injury study. *British Journal of Sports Medicine*, 45, 553–558.
- Ekstrand, J., Hagglund, M., & Walden, M. (2011b). Epidemiology of muscle injuries in professional football (soccer). *American Journal of Sports Medicine*, 39(6), 1226–32. <https://doi.org/10.1177/0363546510395879>
- Ekstrand, J., Walden, M., & Hagglund, M. (2016). Hamstring injuries have increased by 4% annually in men's professional football, since 2001: A 13-year longitudinal analysis of the UEFA elite club injury study. *British Journal of Sports Medicine*, 50, 731e737
- Fanchini, M., Steendahl, I. B., Impellizzeri, F. M., Pruna, R., Dupont, G., Coutts, A. J., Meyer, T., & McCall, A. (2020). Exercise-Based Strategies to Prevent Muscle Injury in Elite Footballers: A Systematic Review and Best Evidence Synthesis. *Sports Medicine*, <https://doi.org/10.1007/s40279-020-01282-z>
- Faude, O., Koch, T., & Meyer, T. (2012). Straight sprinting is the most frequent action in goal situations in professional football. *Journal of Sports Science*, 30(7), 625–31.
- FIFA (2019). Handbook of test methods for Electronic Performance Tracking Devices.
- Fuller, C. W., Ekstrand, J., Junge, A., Andersen, T. E., Bahr, R., Dvorak, J., Hagglund, M., McCrory P., & Meeuwisse, W. H. (2006). Consensus statement on injury definitions and data collection procedures in studies of football (soccer) injuries. *Clinical Journal of Sport Medicine*, 16, 97–106.
- Gonçalves, B., Coutinho, D., & Exel, J. (2019). Extracting spatial-temporal features that describe a team match demands when considering the effects of the quality of opposition in elite football. *PLoS One*, 14, 1–20.
- Gonçalves, L. G. C., Kalva-Filho, C. A., & Nakamura, F.Y. (2020). Effects of match-related contextual factors on weekly load responses in professional Brazilian soccer players. *International Journal of Environmental Research and Public Health*, 17, 1–13.
- Hagglund, M., Walden, M., & Ekstrand, J. (2013a). Risk factors for lower extremity muscle injury in professional soccer: The UEFA injury study. *American Journal of Sports Medicine*, 41, 327–335.
- Häggglund, M., Walden, M., & Magnusson, H. (2013b). Injuries affect team performance negatively in professional football: an 11-year follow-up of the UEFA champions League injury study. *British Journal of Sports Medicine*, 47, 738–42.
- Häggglund, M., Walden, M., Bahr, R., & Ekstrand, J. (2005). Methods for epidemiological study of injuries to professional football players: developing the UEFA model. *British Journal of Sports Medicine*, 39(6), 340-346.
- Hopkins, W. G., Marshall, F. W., Batterham, A. M., & Hanin, J. (2009). Progressive statistics for studies in sport medicine and exercise science. *Medicine Science in Sports Exercise*, 41, 3-13.
- Hoppen, M. I., Reurink, G., de Boode, V. A., van der Kaaden, L., Jagtman, L., Glazenburg, T., & Tol, J. L. (2022). Return to match running performance after a hamstring injury in elite football: a single-centre retrospective cohort study. *BMJ Open Sport & Exercise Medicine*, 8(1), e001240.
- Ingebrigtsen, J., Dalen, T., Hjelde, G. H., Drust, B., & Wisløff, U. (2015). Acceleration and sprint profiles of a professional elite football team in match play. *European Journal of Sport Science*, 15, 101-110.
- Jiménez-Rubio, S., Navandar, A., Rivilla-García, J., Paredes-Hernández, V., & Gómez-Ruano, M. Á. (2020). Improvements in match-related physical performance of professional soccer players after the application of an on-field training program for hamstring injury rehabilitation. *Journal of Sport Rehabilitation*, 29(8), 1145–50. <https://doi.org/10.1123/jsr.2019-0033>
- Knowles, S. B., Marshall, S. W., & Miller, T. (2007). Cost of injuries from a prospective cohort study of North Carolina high school athletes. *Injury Prevention*, 13, 416–21.
- McCall, A., Carling, C., Davison, M., Nedelec, M., Le Gall, F., & Berthoin, S. (2015). Injury risk factors, screening tests and preventative strategies: a systematic review of the evidence that underpins the



- 
- perceptions and practices of 44 football (soccer) teams from various premier leagues. *British Journal of Sports Medicine*, 49(9), 583–9. <https://doi.org/10.1136/bjsports-2014-094104>
- Morgans, R., Bezuglov E., Orme, P., & Di Michele, R. (2022). Technical and physical performance across five consecutive seasons in Russian Premier League soccer. *International Journal of Sports Science & Coaching*, 1–9. DOI: 10.1177/17479541221089247
- Pollock, N., Patel, A., & Chakraverty, J., (2016). Time to return to full training is delayed and recurrence rate is higher in intratendinous ('c') acute hamstring injury in elite track and field athletes: Clinical application of the British athletics muscle injury classification. *British Journal of Sports Medicine*, 50, 305e310
- Rago, V., Rebelo, A., & Krstrup, P., (2021). Contextual Variables and Training Load Throughout a Competitive Period in a Top-Level Male Soccer Team. *Journal of Strength and Conditioning Research*, 35, 3177–3183.
- Raya-González, J., Pulido, J. J., Beato, M., Ponce-Bordón, J. C., López del Campo, R., Resta, R., & García-Calvo, T. (2022). Analysis of the effect of intuire on match performance variables in professional soccer players: a retrospective, experimental longitudinal design. *Sports Medicine-Open*, 8(1), 31.
- Rogalski, B., Dawson, B., Heasman, J., & Gabbett, T. J. (2013). Training and game loads and injury risk in elite Australian footballers. *Journal of Science and Medicine in Sport*, 16, 499–503.
- Seow, D., & Massey A. (2022). Correlation between preseason body composition and sports injury in an English Premier League professional football team. *BMJ Open Sport & Exercise Medicine*, 8, e001193. doi:10.1136/bmjsem-2021-001193
- Taberner, M., Allen, T., & Cohen, D. D. (2019). Progressing rehabilitation after injury: consider the 'control-chaos continuum.' *British Journal of Sports Medicine*, 53(18), 1132–6.
- Taylor, J. B., Wright, A. A., Dischiavi, S. L., Townsend, M. A., & Marmon, A. R. (2017). Activity demands during multi-directional team sports: a systematic review. *Sports Medicine*, 47(12), 2533–2551. doi:10.1007/s40279-017-0772-5
- Whiteley, R., Gregson, W., Bahr, R., Tabben, M., Chamari, K., Lolli, L., & Salvo, V. D. (2022). High-speed running during match-play before and after return from hamstring injury in professional footballers. *Scandinavian Journal of Medicine & Science in Sports*, 32(10), 1502-1509. DOI: 10.1111/sms.14219
- Whiteley, R., Massey, A., Gabbett, T., Blanch, P., Cameron, M., Conlan, G., & Williams, M. (2021). Match high-speed running distances are often suppressed after return from hamstring strain injury in professional footballers. *Sports Health*, 13(3), 290-295. DOI: 10.1177/1941738120964456
- Wing, C. (2018). In-season strength and power training considerations for professional soccer teams competing within national level competitions. *Strength and Conditioning Journal*, 40(3), 12–22.
- Winter, E. M., & Maughan, R. J. (2009). Requirements of ethics approval. *Journal of Sports Science*, 27, 85.