

Central Lancashire Online Knowledge (CLOK)

Title	Are non-starters accumulating enough load compared with starters? Examining load, wellness, and training/match ratios of a European professional soccer team
Type	Article
URL	https://clock.uclan.ac.uk/49280/
DOI	https://doi.org/10.1186/s13102-023-00743-y
Date	2023
Citation	Oliveira, Rafael, Canário-Lemos, Rui, Morgans, Ryland, Rafael-Moreira, Tiago, Vilaça-Alves, José and Brito, João Paulo (2023) Are non-starters accumulating enough load compared with starters? Examining load, wellness, and training/match ratios of a European professional soccer team. BMC Sports Science, Medicine and Rehabilitation, 15 (1).
Creators	Oliveira, Rafael, Canário-Lemos, Rui, Morgans, Ryland, Rafael-Moreira, Tiago, Vilaça-Alves, José and Brito, João Paulo

It is advisable to refer to the publisher's version if you intend to cite from the work.
<https://doi.org/10.1186/s13102-023-00743-y>

For information about Research at UCLan please go to <http://www.uclan.ac.uk/research/>

All outputs in CLOK are protected by Intellectual Property Rights law, including Copyright law. Copyright, IPR and Moral Rights for the works on this site are retained by the individual authors and/or other copyright owners. Terms and conditions for use of this material are defined in the <http://clock.uclan.ac.uk/policies/>

RESEARCH

Open Access



Are non-starters accumulating enough load compared with starters? Examining load, wellness, and training/match ratios of a European professional soccer team

Rafael Oliveira^{1,2,3*}, Rui Canário-Lemos^{4,5}, Ryland Morgans⁶, Tiago Rafael-Moreira^{5,7}, José Vilaça-Alves^{3,4,5} and João Paulo Brito^{1,2,3}

Abstract

Background The aims of the study were to: (i) compare accumulated load and wellness between starters and non-starters of a European professional soccer team; (ii) analyze the relationships between wellness and load measures and; (iii) compare training/match ratio (TMr) of external and internal load between starters and non-starters.

Methods Ten players were considered starters while seven were classified as non-starters over a 16-week period in which six training sessions and match day (MD) were considered in each weekly micro-cycle. The following measures were used: wellness (fatigue, quality of sleep, muscle soreness, stress, and mood); load (rated of perceived exertion (RPE), session-RPE (s-RPE), high-speed running (HSR), sprinting, accelerations (ACC) and decelerations (DEC)). Accumulated wellness/load were calculated by summing all training and match sessions, while TMr was calculated by dividing accumulated training load by match data for all load measures and each player. Mann–Whitney U test was used for wellness variables, while independent T-test was used for the remaining variables to compare groups. Moreover, relationships among variables were explored using the Spearman's Rho correlation coefficient.

Results The main results showed that non-starters presented higher significant values for fatigue ($p < 0.019$; $g = 0.24$) and lower significant values for duration ($p < 0.006$; $ES = 1.81$) and s-RPE ($p < 0.001$; $ES = 2.69$) when compared to starters. Moreover, positive and very large correlation was found between quality of sleep and RPE, while negative and very large correlation were found between stress and deceleration, and mood and deceleration (all, $p < 0.05$). Finally, non-starters presented higher values in all TMr than starters, namely, RPE ($p = 0.001$; $g = 1.96$), s-RPE ($p = 0.002$; $g = 1.77$), HSR ($p = 0.001$; $g = 2.02$), sprinting ($p = 0.002$; $g = 4.23$), accelerations ($p = 0.001$; $g = 2.72$), decelerations ($p < 0.001$; $g = 3.44$), and duration ($p = 0.003$; $g = 2.27$).

Conclusions In conclusion, this study showed that non-starters produced higher TMr in all examined variables despite the lower match and training durations when compared with starters, suggesting that physical load was adjusted appropriately. Additionally, higher RPE was associated with improved sleep while higher number of decelerations were associated with decreased wellness, namely, stress and mood for non-starters.

*Correspondence:

Rafael Oliveira

rafaeloliveira@esdrm.ipsantarem.pt

Full list of author information is available at the end of the article



© The Author(s) 2023. **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated in a credit line to the data.

Keywords Fatigue, Football, Mood, Muscle soreness, Load, Sleep, Stress, Wellbeing, Training load, Load quantification, External load, Internal load, Sports training

Introduction

The quantification of training and match load/demands on soccer players is a common practice [1–3]. Specifically, the monitorisation of athletes include the quantification of training/match demands (e.g., locomotor/mechanical and psychophysiological), and the wellness and readiness of players [4]. On the one hand, locomotor/mechanical demands or just physical demands are associated with external load monitoring using global positioning system (GPS) variables (e.g., distances covered at various running speeds or accelerations). While psychophysiological demands are associated with internal load/intensity monitoring employing subjective and/or objective measures (e.g. rating of perceived exertion (RPE) and heart rate, respectively) [2, 5]. While, wellness is usually measured by questionnaires as previously proposed by Hooper and Mackinnon [6] and McLean et al. [7]. These questionnaires assess various items such as fatigue, quality of sleep, muscle soreness, mood and stress [7], that may vary depending on the specific load applied [4]. Thus, internal load can be estimated from questionnaires whose initial validity presented some concerns, although these issues have recently been resolved due to numerous scientific studies demonstrating individual response sensitivity to training-load changes [8–10].

Despite the relevance that has been attributed to monitoring measures, external load monitoring often raises questions surrounding technology reliability and validity, as well as inconsistencies in those reported measures [11, 12]. However, internal load monitoring has made assumptions regarding player adherence, as well as uncertainty surrounding the impact of these measures on player performance [13]. Furthermore, despite questions over the reliability of GPS data [11] and on internal load monitoring instruments, research has supported their implementation in a variety of team sports [13].

Factors such as player age, competitive level, playing style and playing status are some examples of variables that can affect physical demands. In this regard, some studies reported significant differences between playing status, where starters presented higher values than non-starters in professional male and female soccer players [14–17]. However, those studies only examined variables such as training monotony, strain, acute:chronic workload ratios, while further analysis may provide additional key findings for coaches and performance staff. For instance, analysing accumulated load with and without match data may provide different

interpretations and it is expected that starters and/or players with higher match participation present greater values of accumulated load [18]. Additionally, analyzing the relationship between total load, including match participation and wellness is warranted as wellness has previously been modulated by such factors [19]. The relationship between wellness and load is fundamental for coaches in order to understand the player's state of fitness and freshness considering the total workload. Only in this way, is it possible to have an adequate and individualized load management [20]. Through a comprehensive and integrated monitoring strategy to assess changes in players (such as external load, internal load, wellness status, and readiness for training and match), can coaches implement appropriate training programs and recovery strategies to optimize player performance.

Some calculations such as training/match ratio (TMr) may also be extremely helpful to understand the physical demands induced by competitive match-play in starters and to further understand if compensatory training sessions are an adequate stimulus for non-starters [21, 22]. Furthermore, TMr may be fundamental to decreasing injury risk and optimizing performance [23]. The TMr are calculated by the division of the accumulated weekly load by the match load [22]. If the TMr provides a value equal to one, it suggests that the accumulated load of the week is equal to the match [22]. Usually, players in a standard micro-cycle should have a TMr of at least one, as exposure to match demands is the first prevention of the risk of injury, however, if excessively high, it also increases its risk of occurrence [24]. Even so, the oscillation between higher or lower values can vary greatly from context to context, from player to player and from metric to metric. League characteristics and game/training models can have an influence on what is accumulated training load during the week. In this way, there are teams that, due to their game model, end up accumulating more ACC/DEC than HSR and sprint [25], players who, due to their position, may be more exposed to certain types of efforts [26] and even individual characteristics that, like the moderators who influence how load is tolerated by players [27].

Due to scant recent studies, it would seem reasonable to compare starters and non-starters data [21]. For example, Stevens et al. [21] reported lower TMr values for non-starters compared to starters, specifically for high-speed running (HSR). However, Clemente et al.

[22] analyzed the differences between varying micro-cycle schedules (3 versus 4 versus 5, but did not consider playing status). Nonetheless, the authors found that both TMr of HSR (HSRr) and TMr of sprint distance (SPDr) were higher in week in five sessions than weeks with three or four sessions. They concluded that training sessions were not adjusted according to weekly variations in match demands. Understanding individual TMr may be important to adjust training characteristics considering the needs of each player.

Therefore, the aims of this study were to: (i) compare accumulated load and wellness between starters and non-starters of a European professional soccer team; (ii) analyze the relationships between wellness and load measures and; (iii) compare TMr of external and internal load between starters and non-starters. The study hypothesis was that non-starters would present lower accumulated load and wellness values, and lower TMr when compared with starters [21]. Consequently, trivial-to-small correlations would be observed between load and wellness specifically for non-starters, considering that non-starters tend to present lower load during micro-cycles [21].

Materials and methods

Experimental approach to the problem

This was an observational study design. The players were monitored daily for wellness measures (sleep quality, muscle soreness, fatigue, stress and mood), internal and external load during 70 training sessions and 15 official matches over a 16-week period during the 2022/23 in-season period (July to November).

All micro-cycles were organized according to the following match, and thus all micro-cycles included only one match. In addition, when a player (non-starter) did not participate in MD, MD + 1 consisted of exercises to expose players to higher levels of HSR and sprinting distances and accelerations (ACC) and decelerations (DEC), while starters performed a recovery session. Therefore, to accomplish HSR and sprint distances, chasing opponents and goal-scoring exercises were performed, while small-sided games were performed to accomplish ACC and

DEC targets. Some examples of scoring exercise included running to shoot on goal and then returning to defend the next player who will be shooting in the opposite goal, this way HSR and sprinting were accomplished. Ball possession exercises included two squares and three teams. The team that recovers the ball connects with the opposite square and the team that lost ball possession has to press immediately to regain the ball. Table 1 presents the number of training sessions and matches for each day of the micro-cycle.

Participants

Seventeen highly trained/national level professional male soccer players (age: 25 ± 2.8 years; body mass: 71 ± 6.6 kg; body height: 177.9 ± 6.7 cm; fat mass: $8.6 \pm 1.0\%$; professional experience: 7.4 ± 2.7 years) participated in the current study. Players belonged to a European soccer team that played in the national first division. From the 17 players included, 7 were defenders, 5 were midfielders and 5 were attackers.

The eligibility criteria for participant inclusion were: (i) participating in 80% of the total number of training sessions (full session duration) [19] and; (ii) completed all wellness and training reports over the data collection period. In addition, players were considered starters if they started the match and completed at least 60-min in three consecutive matches while non-starters were the remaining players [14–17]. Thus, 10 players were considered starters (age: 25.1 ± 2.2 years; body mass: 73.4 ± 8.7 kg; body height: 180.7 ± 9.0 cm; fat mass: $8.7 \pm 1.1\%$; professional experience: 7.3 ± 2.3 years), while seven were considered non-starters (age: 26.1 ± 4.6 years; body mass: 71.4 ± 7.2 kg; body height: 178.0 ± 6.3 cm; fat mass: $8.9 \pm 1.5\%$; professional experience: 8.4 ± 4.2 years).

Prior to data collection, participants were fully informed of the study design and signed an informed consent. The study followed the ethical guidelines for human study as suggested by the Declaration of Helsinki (2013). Furthermore, the study was approved by the research Ethics Committee of the Polytechnic Institute of Santarém, Santarém, Portugal (Nº24-2022ESDRM).

Table 1 Number of training/match sessions of the 16-week period

Day of the week	MD + 1	MD-5	MD-4	MD-3	MD-2	MD-1	MD ^a
Number of sessions	13	6	9	13	15	16	16
Duration for starters (min)	24.2 ± 6.3	38.4 ± 2.6	44.8 ± 3.6	39.3 ± 1.0	35.2 ± 2.1	39.2 ± 5.6	80.2 ± 17.3
Duration for non-starters (min)	38.2 ± 6.5	39.0 ± 2.1	45.1 ± 3.5	40.8 ± 2.2	36.0 ± 2.0	26.6 ± 2.3	43.8 ± 16.3

MD Match day, MD + 1 One day after the match day, MD-5 Five days before match day, MD-4 Four days before match day, MD-3 Three days before match day, MD-2 Two days before match day, MD-1 One day before match day

^a for non-starters this was considered a training session

Wellness quantification

Following previous recommendations of overtraining monitoring [6], the wellness questionnaire of McLean et al. [7] was applied individually 30-min prior to each training/match session through a google form specifically designed. The questionnaire uses a scale of 1–5 arbitrary units (A.U.) and has five questions assessing fatigue, quality of sleep, muscle soreness, stress and mood (in which 5 is very fresh, very restful, very great, very relaxed and very positive mood, respectively, while 1 suggests always tired, insomnia, very sore, highly stressed and highly annoyed/irritable/down, respectively) [7]. All players were familiarized with the questionnaire during the previous full season.

Internal load quantification

The CR-10 Borg's scale [28] was employed to monitor players internal load (RPE) relative to the physiological and/or psychological effect of the work [1]. Using a valid and reliable practical tool is imperative for monitoring the training load imposed on athletes during every training session. Foster et al. [29] proposed a method based on RPE. This method, known as the session-RPE method, considers both the intensity and duration of a training session. Several studies confirmed the validity, reliability and internal consistency of the session-RPE method in soccer with players of varying levels [30, 31].

Following 20- to 30-min after each session, every player provided a perceived exertion value using a google form specifically designed by answering the following question: "how intense was the training session?". The scale varied from 0 to 10 A.U., where each value rated as: 0 – nothing to all; 0.5 – extremely weak; 1 – very weak; 2 – weak; 3 – moderate; 4 – somewhat strong; 5 – strong; 7 – very strong; and 10 – extremely strong.

The score was used as a measure of internal intensity: RPE. In addition, the duration of the entire training session and/or match in minutes was multiplied by the RPE to generate the session-RPE (s-RPE) (A.U.) [29, 32]. All players were already familiarized with the questionnaire during the previous full season.

External load quantification

Locomotor demands were measured using a 10 Hz GPS Vector S7 (Catapult Innovations, Melbourne, Australia). To avoid inter-unit bias, the same unit was used for each player throughout the analysis period [33]. The unit was placed on the upper back of the players, within a custom-made vest, 30-min prior to each session (training and match) and removed immediately post-session.

All data were transferred to a Microsoft Excel spreadsheet from Catapult Openfield software (Version 1.21.1). The data were grouped by training session, although only

actual training time was considered for analysis, that is, data from the warm-up onwards, with transition time between exercises and rest times removed.

The GPS device has previously been validated for accuracy and reliability regarding sprint acceleration profiles [34] and in measures such as distance, speed and average acceleration where the coefficient of variation ranged from 0.1 to 3.9% (reliability for running distance measures and average acceleration) [35]. The following metrics were used for analysis: (i) high-speed running (HSR, 20–25 km/h) and sprinting (SPD, > 25 km/h) based on Houtmeyers et al. [36], number of accelerations (ACC, > 2 m/s²) and number of decelerations (DEC, < 2 m/s²) based on Gonçalves et al. [37].

Accumulated wellness/load and training/match ratio

Accumulated wellness/load consisted of the sum of each measure during all training sessions of the micro-cycle and was calculated per player, thus providing the weekly load for each measure (MD included).

Moreover, accumulated load was also calculated without MD data in order to calculate training/match ratio (TMr) for all internal and external measures. Ratios were then calculated by dividing accumulated load (without MD data) by MD data (TMr = weekly load/match demands) [22]. This calculation was made for each player and then, average values were used for each group (starters and non-starters). Consequently, the following measures were obtained: RPEr (rating of perceived exertion ratio); s-RPEr (session rating of perceived exertion ratio); HSRr (high-speed running ratio); SPDdr (sprinting distance ratio); ACCr (accelerations ratio) and; DECdr (decelerations ratio). The same ratio was calculated for session duration (Dr).

Statistical analysis

Descriptive statistics are presented as mean ± standard deviation. Normality and homogeneity of the different variables was tested using the Shapiro–Wilk and Levene tests, respectively. Only wellness variables did not present normal distribution ($p < 0.05$). Thus, Mann–Whitney U test was used for wellness variables while independent T-test was used for the remaining variables. Significant results were considered at $p < 0.05$, while the Hedges effect-size was performed to determine the effect magnitude through the difference of two means divided by the standard deviation from the data and the following criteria were used: < 0.2 = trivial, 0.2 to 0.6 = small effect, 0.6 to 1.2 = moderate effect, 1.2 to 2.0 = large effect, and > 2.0 = very large [38].

Finally, the relationship between wellness and load variables were explored using the Spearman's Rho correlation coefficient. The magnitude of correlations were

classified as trivial (0.00 to 0.09), small (0.10 to 0.29), moderate (0.30 to 0.49), large (0.50 to 0.69), very large (0.70 to 0.89), and nearly perfect (>0.90) [39].

All statistical procedures were executed in the IBM SPSS Statistics for Windows version 23.0 (IBM Corp, Armonk, NY:USA).

Results

Table 2 presents the accumulated training demands (MD data included) for all variables. Non-starters presented significantly higher values for fatigue and lower

significant values for duration and s-RPE when compared with starters. There were no other significant differences.

Table 3 presents the correlation coefficients among wellness and the physical demands using the accumulated micro-cycle (including the MD) values for starters. Significant positive and nearly perfect correlations were found between fatigue and muscle soreness, muscle soreness and mood, and stress and mood. In addition, significant positive very large correlations were found between fatigue and stress, muscle soreness and stress, fatigue and mood, and ACC and DEC. Finally, there was a significant

Table 2 Descriptive statistics (mean \pm standard deviation, and confidence interval, CI, 95%) for accumulated wellness and load demands

Variables	Starters	Non-starters	p-value	Effect size
Quality of sleep (A.U.)	26.2 \pm 2.4 (24.5–27.9)	26.9 \pm 3.0 (24.1–29.7)	0.536	-
Fatigue (A.U.)	22.8 \pm 3.0 (20.6–25.0)	26.3 \pm 2.7 (23.8–28.8)	0.019*	0.24
Muscle Soreness (A.U.)	22.4 \pm 12.8 (19.9–25.0)	25.5 \pm 3.0 (22.7–28.3)	0.070	-
Stress (A.U.)	24.4 \pm 3.5 (22.4–27.4)	25.2 \pm 3.8 (21.7–28.7)	0.740	-
Mood (A.U.)	24.9 \pm 3.1 (22.7–27.2)	25.1 \pm 4.0 (21.4–28.0)	>0.999	-
RPE (A.U.)	36.4 \pm 2.0 (34.9–37.8)	35.7 \pm 3.0 (32.9–38.5)	0.594	-
Duration (min)	291.3 \pm 11.9 (282.8–299.9)	269.5 \pm 16.4 (254.4–284.7)	0.006*	1.81
s-RPE (A.U.)	1837.3 \pm 105.6 (1761.8–1912.9)	1552.4 \pm 170.5 (1394.7–1710.2)	0.001*	2.69
High speed running (m)	1107.9 \pm 208.9 (208.9–1257.4)	1132.4 \pm 86.3 (1052.6–1212.2)	0.745	-
Sprinting (m)	278.9 \pm 87.2 (216.6–341.3)	320.0 \pm 54.6 (269.5–370.5)	0.289	-
Accelerations (nr)	291.7 \pm 37.0 (265.2–318.2)	306.4 \pm 21.1 (286.9–325.9)	0.359	-
Decelerations (nr)	268.0 \pm 26.9 (248.0–289.3)	283.3 \pm 28.3 (257.1–309.5)	0.315	-

RPE Rate of perceived exertion using the CR-10 Borg's scale, s-RPE Multiplication of session time by the RPE score, A.U Arbitrary units, m Meters, min Minutes, * denotes significant difference between starters versus non-starters ($p < 0.05$)

Table 3 Correlation coefficient (r) for wellness and load demands in starters

Variables	Fatigue	Muscle Soreness	Stress	Mood	RPE	Time of session	s-RPE	HSR	Sprinting	ACC	DEC
Quality of sleep	0.527	0.527	0.503	0.479	-0.333	-0.030	-0.588	0.079	0.079	0.442	0.115
Fatigue		0.915 $p < 0.001$	0.745 $p = 0.013$	0.782 $p = 0.008$	-0.030	-0.139	-0.188	0.236	0.236	0.467	0.491
Muscle Soreness			0.806 $p = 0.005$	0.915 $p < 0.001$	0.030	-0.370	-0.345	0.358	0.224	0.370	0.382
Stress				0.903 $p < 0.001$	-0.224	0.030	-0.115	0.030	-0.224	0.067	0.127
Mood					-0.055	-0.188	-0.333	0.103	-0.006	0.067	0.127
RPE						-467	0.188	0.261	0.479	0.261	0.406
Time of session							0.527	-0.442	-0.200	-0.333	-0.236
s-RPE								0.115	0.103	-0.103	0.224
HSR									0.612	0.648 $p = 0.043$	0.685 $p = 0.029$
Sprinting										0.430	0.430
ACC											0.891 $p = 0.001$

RPE Rate of perceived exertion using the CR-10 Borg's scale, s-RPE Multiplication of session time by the RPE score, HSR High speed running (20–25 km/h), ACC Acceleration, DEC Deceleration, Bold denotes significant correlations

large correlation between HSR and ACC and HSR and DEC.

Table 4 presents the correlation coefficients among wellness and physical demands using the accumulated micro-cycle (including the MD) values for non-starters. Significant positive and nearly perfect correlations were found between: stress and mood. Moreover, significant positive very large correlations were found between fatigue and muscle soreness, quality of sleep and RPE. Finally, significant negative very large correlations were found between stress and DEC, and mood and DEC.

Figure 1 shows that non-starters presented higher values in all TMr than non-starters, namely, RPEr ($p=0.001$; $g=1.96$), s-RPEr ($p=0.002$; $g=1.77$), HSRr ($p=0.001$; $g=2.02$), SPD_r ($p=0.002$; $g=4.23$), ACC_r ($p=0.001$; $g=2.72$), DEC_r ($p<0.001$; $g=3.44$), Dr ($p=0.003$; $g=2.27$).

Discussion

The aims of this study were to: (i) compare accumulated load and wellness between starters and non-starters of a European professional soccer team; (ii) analyze the relationship between wellness and load measures and; (iii) compare TMr of external and internal load between starters and non-starters. The main results of the study reported that non-starters presented significantly higher values for fatigue and significantly lower values for duration and s-RPE when compared with starters. In addition, starters showed positive correlations between fatigue and muscle soreness, muscle soreness and mood, stress and mood, fatigue and stress, muscle soreness and stress, fatigue and mood, ACC and DEC, HSR and ACC, and HSR and DEC. While non-starters reported correlations between stress and mood, fatigue and muscle soreness, quality of sleep and RPE, and negative correlations between stress and DEC, and mood and DEC.

There is a lack of evidence comparing differences between the varying status of players (starter versus non-starter) of load and wellness measures, thus this may partly be explained by the specification of MD+1 training. In these training sessions, non-starters performed compensatory training that focussed on physical demands to ensure these players were provided with a stimulus similar to that of starters (from the previous match). However, the characteristics of these training tasks for non-starters, sometimes with more analytical exercises, focusing on fitness development and less specific dynamics of the game [40], resulted in higher values for duration and s-RPE on MD+1 (for instance, starters presented 24 min and 101.8 A.U. while non-starters 38 min and 230.6 A.U. of duration and s-RPE, respectively). In this regard, some authors [41] have reported that wellness measures have sometimes revealed an

element of subjectivity when associated with load monitoring variables measured through more conventional objective markers. Furthermore, it has also been argued that despite the growing interest and use of wellness measures, there is still a lack of a general frame referenced in the current literature [42].

Nonetheless, the other notable study result was that TMr comparisons reported higher values for non-starters in all variables despite lower match and training durations when compared with starters. Once again, it may be speculated that the compensatory training session (MD+1) as a strategy to resolve differential weekly training load between starting and non-starting players' may partly explain these results, specifically highlighting the effort and commitment during match minutes. Regarding the first aim, the present study results are not in accordance with previous research, that showed higher fatigue, stress and muscle soreness levels for starters compared with non-starters over a 40-week period [17]. Such findings were also corroborated by Nobari et al. [18] that found higher weekly acute s-RPE, fatigue, and stress occurred in starters when compared with non-starters at the end of the competitive season, albeit in elite young soccer players [43]. Furthermore, it is relevant to note that despite the non-significant differences in external load measures, starters showed lower values in comparison to non-starters thus contradicting previous literature that attributed differences to match playing time [44, 45]. The findings of the existing literature do not fully explain the present results as accumulated duration was higher for starters. Thus, the external load of the study players seems to be adjusted appropriately for starters and non-starters.

Considering the second aim of the study, the most relevant associations occurred between load and wellness measures. In this regard, starters only showed an association within wellness measures (e.g., fatigue and muscle soreness) and within external load measures (e.g., sprint and acceleration), respectively. However, non-starters highlighted associations between the quality of sleep and RPE, stress and DEC, and mood and DEC. A study examining youth soccer players found higher external and internal intensity to be associated with improved sleep (quality and quantity), and feeling rested, although did not consider playing status [46] that contrasts the present results. Contrastingly, a different study in youth soccer players found that high-intensity training had no impact on the following night's sleep quality [47]. While a further study conducted with professional soccer players also reported that sleep quality was not affected by higher intensity sessions (MD included) [48]. Again, these previous studies did not consider the starting status of players, starters and non-starters. Thus, more research

Table 4 Correlation coefficient (r) for wellness and load demands in non-starters

Variables	Fatigue	Muscle Soreness	Stress	Mood	RPE	Time of session	s-RPE	HSR	Sprinting	ACC	DEC
Quality of sleep	0.321	0.321	0.179	0.286	0.893 p = 0.007	0.143		0.214	-0.286	-0.036	-0.357
Fatigue		0.786 p = 0.036	0.321	0.393	0.107	0.464	0.250	-0.393	-0.250	-0.536	-0.429
Muscle Soreness			0.714	0.750	0.179	0.679		0.536	< 0.001	-0.536	-0.643
Stress				0.964 p < 0.001	-0.071	0.179		0.250	0.071	-0.750	-0.786 p = 0.036
Mood					0.071	0.143	0.286	0.214	0.036	-0.786	-0.857 p = 0.014
RPE						0.214	0.750	0.250	-0.071	0.286	-0.107
Time of session							0.714	-0.241	-0.036	0.214	-0.143
s-RPE								0.071	-0.214	0.250	-0.357
HSR									0.714	< 0.001	0.214
Sprinting										0.036	0.464
ACC											0.679

RPE Rate of perceived exertion using the CR-10 Borg's scale, s-RPE Multiplication of time of session by the score of RPE, HSR High speed running (20–25 km/h), ACC Acceleration, DEC Deceleration, Bold denotes significant correlations




Fig. 1 Accumulative weekly match load, training load, and training/match ratios for RPE, s-RPE, HSR, SPD, ACC, DEC and Duration. RPEr: rating of perceived exertion ratio; s-RPEr: session rating of perceived exertion ratio; HSRr: high-speed running ratio; SPD: sprinting distance ratio; ACCr: accelerations ratio; DEC: decelerations ratio; Dr: duration ratio. * denotes significant difference for starters ($p < 0.05$)

is warranted to investigate these specific groups. Nonetheless, the relationship between perceived exertion and a higher number of DEC seems to decrease wellness, namely, stress and mood while sleep is improved for non-starters.

Furthermore, relevant findings of the current study were related to aim (iii), where non-starters revealed higher TMR in all variables regardless of the lower match and training durations when compared with starters. Such findings are in contrast with some studies that attributed higher match-play minutes as a possible explanation for such differences. However, those studies found higher values of HSR, sprint distance, number of ACC and number of repeated sprints for starters [44, 45, 49]. This may suggest that coaches and performance staff adjusted training sessions for non-starters regardless of match participation. Considering the comparison of starters and non-starters, it seems that only Stevens et al. [21] conducted such analysis utilizing TMR. The authors found lower TMR values for non-starters when compared to starters with significant differences for HSR, while the remaining running metrics and accelerometry variables were also higher for starters, thus contradicting the current study findings.

Additionally, the calculation of TMR using both RPE and s-RPE was, to our knowledge, the first study to report such a method. Nonetheless, the external TMR values found in the present study for starters ranged from 1.3 to

2.3 A.U, while non-starters presented a range between 2.9 and 5.1 A.U, that is similar to those previously reported by Clemente et al. [22] who found a range of ~1 to 4 A.U for Portuguese soccer players. Although, these data were higher than those reported by Stevens et al. [21] for HSR (0.22) and sprint distance (0.03) in Dutch soccer players. However, such ratios are dependent on the number of training sessions per week. Therefore, possible explanations for the different values may be associated with different training strategies that included additional conditioning exercises and/or supplementary sessions that may have been conducted as previously suggested [22]. Nevertheless, in the present study it was found that non-starters had higher TMR values for ACC and DEC which seems to suggest that additional training for non-starters (performed on MD + 1) consisted of small-sided games, as this game format increases the number of ACC/DEC and decreases HSR and sprint distances covered [50, 51].

Notwithstanding, the present study had some limitations. The main limitation is the small sample size that derived from only one team, and across a restricted time period of 16-weeks. Furthermore, as reported in Table 1, different training days per week occurred that may have influenced TMR values. Even so, average values were employed and produced relevant information for coaches and performance staff. Moreover, playing position differences were not considered for analysis and for some positions, such as wide players (attacking and defending),

perform higher accumulated values, TMr and the perceived various wellness status, as these positions produce greater running effort than other positions [52]. Consequently, future studies should attempt to analyze larger sample sizes, include different training frequencies [22] and analyze regular weeks with one match versus congested weeks with more than one match as previously suggested [21]. Additionally, the generalization of these results to other teams/countries, competitive standards and ages is not recommended and thus further replication studies are required. For instance, recently no external load difference was observed between an under-18 and a first team [53], although the analysis of TMr would provide more insights for coaches. Finally, a detailed description of training drills in future research may facilitate a more precise practical implementation.

Although these results may be dependent on the analyzed team, this study showed that it is possible to plan higher load for non-starters during the in-season without negatively impacting training for starters. From a load perspective, non-starters need to be physically prepared to replace or substitute a starter. Moreover, TMr analysis facilitates the interpretation and contextualization of data, and consequently allows the training prescription to be planned accordingly to achieve the appropriate load and also allows coaches and performance staff to communicate effectively with each other and players as previously suggested [21]. Finally, the individuality and how athletes recover varies from player to player. Some athletes perceive a high load and recover well, while other athletes recover slower, hence the moderators explain that absolute and/or equal load can be tolerated differently. Thus, this supports the notion to interpretate results with caution.

Conclusions

In summary, it was observed that non-starters presented significantly higher values for fatigue (suggesting less fatigue) and lower significant values for duration and s-RPE than starters. Specifically, non-starters reported higher TMr in all variables despite lower match and training durations when compared with starters suggesting that physical load was adjusted appropriately.

Furthermore, non-starters showed positive relationships between stress and mood, fatigue and muscle soreness, and quality of sleep and RPE. Moreover, non-starters reported negative relationships between stress and DEC and mood and DEC. These findings seemed to suggest that perceived exertion and a higher number of DEC may contribute to decreased wellness, namely, quality of sleep and stress and mood for non-starters. The current study also provides practical information

regarding the micro-cycle periodization, and suggests that the type of exercises used during the compensatory training sessions were appropriate for this specific group. Finally, although this study provides average values, it is not recommended to apply these findings to other teams and coaches from similar contexts.

Abbreviations

TMr	Training/match ratio
MD	Match day
MD + 1	One day after the match day
MD-5	Five days before match day
MD-4	Four days before match day
MD-3	Three days before match day
MD-2	Two days before match day
MD-1	One day before match day
RPE	Rating of perceived exertion
s-RPE	Session rating of perceived exertion
HSR	High-speed running
ACC	Acceleration
DEC	Deceleration
GPS	Global positioning system
A.U.	Arbitrary units
RPEr	Rating of perceived exertion ratio
s-RPEr	Session rating of perceived exertion ratio
HSRr	High-speed running ratio
SPDr	Sprinting distance ratio
ACCr	Accelerations ratio
DECr	Decelerations ratio
Dr	Duration ratio

Acknowledgements

Not applicable.

Authors' contributions

Conceptualization, R.O., R.C-L. and J.P.B.; methodology, R.O., R.C-L. and J.P.B.; data collection, R.C-L.; formal analysis, R.O.; data curation, R.O. and R.C-L.; writing—original draft preparation, R.O., R.C-L. T.R-M., J. V-A., R.M. and J.P.B.; writing—review and editing, R.O., R.C-L. T.R-M., J. V-A., R.M. and J.P.B.; supervision, R.O., R.M. and J.P.B.; funding, R.O. and J.P.B. All authors have read and agreed to the published version of the manuscript.

Funding

This research was funded by the Portuguese Foundation for Science and Technology, I.P., Grant/Award Number UIDP/04748/2020. The funders had no role in the design of the study; in the collection, analyses, or interpretation of data; in the writing of the manuscript, or in the decision to publish the results.

Availability of data and materials

The data presented in this study are available on request from the corresponding author.

Declarations

Ethics approval and consent to participate

The study was conducted according to the guidelines of the Declaration of Helsinki and approved by the Ethics Committee of Polytechnic Institute of Santarém (Nº24-2022ESDRM). Informed consent was obtained from all subjects involved in the study.

Consent for publication

Not applicable.

Competing interests

The authors declare no competing interests.

Author details

¹Sports Science School of Rio Maior–Polytechnic Institute of Santarém, Rio Maior 2040-413, Portugal. ²Life Quality Research Centre, Rio Maior 2040-413, Portugal. ³The Research Centre in Sports Sciences, Health Sciences and Human Development (CIDESD), Vila Real 5001-801, Portugal. ⁴University of Trás-Os-Montes E Alto Douro, Vila Real 5001-801, Portugal. ⁵Research Group in Strength Training and Fitness Activities, GEETFAA, 5001-801 Vila Real, Portugal. ⁶Football Performance Hub, University of Central Lancashire, Preston PR1 2HE, UK. ⁷Research Center of the Polytechnic Institute of Maia (N2i), Maia Polytechnic Institute (IPMAIA), Castêlo da Maia, Maia 4475-690, Portugal.

Received: 21 March 2023 Accepted: 1 October 2023

Published online: 10 October 2023

References

- Halsen SL. Monitoring Training Load to Understand Fatigue in Athletes. *Sport Med*. 2014;44:139–47.
- Miguel M, Oliveira R, Loureiro N, García-Rubio J, Ibáñez SJ. Load measures in training/match monitoring in soccer: A systematic review. *Int J Environ Res Public Health*. 2021;18:1–26.
- Williams S, Trewartha G, Cross MJ, Kemp SPT, Stokes KA. Monitoring What Matters: A Systematic Process for Selecting Training-Load Measures. *Int J Sports Physiol Perform*. 2017;12(Suppl 2):S2–101–S2–106.
- Gabbett TJ, Nassis GP, Oetter E, Pretorius J, Johnston N, Medina D, et al. The athlete monitoring cycle: a practical guide to interpreting and applying training monitoring data. *Br J Sports Med*. 2017;51:1451–2.
- Bourdon PC, Cardinale M, Murray A, Gastin P, Kellmann M, Varley MC, et al. Monitoring athlete training loads: Consensus statement. *Int J Sports Physiol Perform*. 2017;12:161–70.
- Hooper SL, Mackinnon LT. Monitoring Overtraining in Athletes: Recommendations. *Sport Med*. 1995;20:321–7.
- McLean BD, Coutts AJ, Kelly V, McGuigan MR, Cormack SJ. Neuromuscular, endocrine, and perceptual fatigue responses during different length between-match microcycles in professional rugby league players. *Int J Sports Physiol Perform*. 2010;5:367–83.
- Gastin PB, Meyer D, Robinson D. Perceptions of wellness to monitor adaptive responses to training and competition in elite Australian football. *J strength Cond Res*. 2013;27:2518–26.
- Montgomery PG, Hopkins WG. The Effects of Game and Training Loads on Perceptual Responses of Muscle Soreness in Australian Football. *Int J Sports Physiol Perform*. 2013;8:312–8.
- Buchheit M, Racinais S, Bilsborough JC, Bourdon PC, Voss SC, Hocking J, et al. Monitoring fitness, fatigue and running performance during a pre-season training camp in elite football players. *J Sci Med Sport*. 2013;16:550–5.
- Cummins C, Orr R, O'Connor H, West C. Global positioning systems (GPS) and microtechnology sensors in team sports: a systematic review. *Sports Med*. 2013;43:1025–42.
- Ammann L, Altmann S. Training and match load ratios in professional soccer—should we use player- or position-specific match reference values? *Front Sport Act Living*. 2023;5:1151828.
- Burgess DJ. The Research Doesn't Always Apply: Practical Solutions to Evidence-Based Training-Load Monitoring in Elite Team Sports. *Int J Sports Physiol Perform*. 2017;12:S2–136–S2–141.
- Nobari H, Oliveira R, Clemente FM, Adsuar JC, Pérez-Gómez J, Carlos-Vivas J, et al. Comparisons of Accelerometer Variables Training Monotony and Strain of Starters and Non-Starters: A Full-Season Study in Professional Soccer Players. *Int J Environ Res Public Heal*. 2020;17:6547.
- Nobari H, Praça GM, Clemente FM, Pérez-Gómez J, Carlos Vivas J, Ahmadi M. Comparisons of new body load and metabolic power average workload indices between starters and non-starters: A full-season study in professional soccer players. *Proc Inst Mech Eng Part P J Sport Eng Technol*. 2020. <https://doi.org/10.1177/1754337120974873>.
- Fernandes R, Brito JP, Vieira LHP, Martins AD, Clemente FM, Nobari H, et al. In-season internal load and wellness variations in professional women soccer players: Comparisons between playing positions and status. *Int J Environ Res Public Health*. 2021;18(23):12817.
- Oliveira R, Ceylan Hİ, Brito JP, Martins A, Nalha M, Mendes B, et al. Within- and between-mesocycle variations of well-being measures in top elite male soccer players: a longitudinal study. *J Mens health*. 2022;18:1–14.
- Oliveira R, Vieira LHP, Martins A, Brito JP, Nalha M, Mendes B, et al. In-season internal and external workload variations between starters and non-starters—a case study of a top elite European soccer team. *Med*. 2021;57:1–15.
- Clemente FM, Mendes B, Nikolaidis PT, Calvete F, Carriço S, Owen AL. Internal training load and its longitudinal relationship with seasonal player wellness in elite professional soccer. *Physiol Behav*. 2017;179:262–7.
- Moalla W, Fessi MS, Farhat F, Nouira S, Wong DP, Dupont G. Relationship between daily training load and psychometric status of professional soccer players. *Res Sport Med*. 2016;24:387–94.
- Stevens T, Ruiter C, Twisk L, Savelsbergh G, Beek P. Quantification of in-season training load relative to match load in professional Dutch Eredivisie football players. *Sci Med Footb*. 2017;1:117–25.
- Clemente FM, Rabbani A, Conte D, Castillo D, Afonso J, Clark CCT, et al. Training/match external load ratios in professional soccer players: A full-season study. *Int J Environ Res Public Health*. 2019;16:1–11.
- Gabbett TJ. The training — injury prevention paradox : should athletes be training smarter and harder? 2016;50(5):273–80.
- Malone S, Owen A, Mendes B, Hughes B, Collins K, Gabbett TJ. High-speed running and sprinting as an injury risk factor in soccer: Can well-developed physical qualities reduce the risk? *J Sci Med Sport*. 2018;21:257–62.
- Casamichana D, Martín-García A, Díaz AG, Bradley PS, Castellano J. Accumulative weekly load in a professional football team: with special reference to match playing time and game position. *Biol Sport*. 2022;39:115–24.
- Malone J, Di Michele R, Morgans R, Burgess D, Morton J, Drust B. Seasonal Training-Load Quantification in Elite English Premier League Soccer Players. *Int J Sports Physiol Perform*. 2015;10:489–97.
- Gabbett TJ, Nielsen RO, Bertelsen ML, Bittencourt NFN, Fonseca ST, Malone S, et al. In pursuit of the “Unbreakable” Athlete: what is the role of moderating factors and circular causation? *Br J Sports Med*. 2019;53:394–5.
- Borg G. Perceived exertion and pain scales. Champaign IL, USA: Human Kinetics; 1998.
- Foster C, Florhaug JA, Franklin J, Gottschall L, Hrovatin LA, Parker S, et al. A New Approach to Monitoring Exercise Training. *J Strength Cond Res*. 2001;15:109–15.
- Haddad M, Stylianides G, Djaoui L, Dellal A, Chamari K. Session-RPE method for training load monitoring: Validity, ecological usefulness, and influencing factors. *Front Neurosci*. 2017;11:612.
- Gaudino P, Iaia FM, Strudwick AJ, Hawkins RD, Alberti G, Atkinson G, et al. Factors Influencing Perception of Effort (Session Rating of Perceived Exertion) During Elite Soccer Training. *Int J Sports Physiol Perform*. 2015;10:860–4.
- Foster C, Hector LL, Welsh R, Schrager M, Green MA, Snyder AC. Effects of specific versus cross-training on running performance. *Eur J Appl Physiol Occup Physiol*. 1995;70:367–72.
- Scott MTU, Scott TJ, Kelly VG. The Validity and Reliability of Global Positioning Systems in Team Sport: A Brief Review. *J strength Cond Res*. 2016;30:1470–90.
- Clavel P, Leduc C, Morin J-B, Owen C, Samozino P, Peeters A, et al. Concurrent Validity and Reliability of Sprinting Force-Velocity Profile Assessed With GPS Devices in Elite Athletes. *Int J Sports Physiol Perform*. 2022;17:1527–31.
- Crang ZL, Duthie G, Cole MH, Weakley J, Hewitt A, Johnston RD. The inter-device reliability of global navigation satellite systems during team sport movement across multiple days. *J Sci Med Sport*. 2021;25:340–4.
- Houtmeyers KC, Jaspers A, Brink MS, Vanrenterghem J, Varley MC, Helsen WF. External load differences between elite youth and professional football players: ready for take-off. *Sci Med Footb*. 2020;0:1–5.
- Gonçalves LGC, Kalva-Filho CA, Nakamura FY, Rago V, Afonso J, de Bedo BL S, et al. Effects of match-related contextual factors on weekly load responses in professional Brazilian soccer players. *Int J Environ Res Public Health*. 2020;17:1–13.

38. Hopkins WG, Marshall SW, Batterham AM, Hanin J. Progressive statistics for studies in sports medicine and exercise science. *Med Sci Sports Exerc.* 2009;41:3–13.
39. Batterham AM, Hopkins WG. Making Meaningful Inferences about Magnitudes. *Int J Sports Physiol Perform.* 2006;1:50–7.
40. Clemente FM, Lourenço Martins FM, Mendes RS. Developing aerobic and anaerobic fitness using small-sided soccer games: Methodological proposals. *Strength Cond J.* 2014;36:76–87.
41. Saw AE, Main LC, Gastin PB. Monitoring the athlete training response: subjective self-reported measures trump commonly used objective measures: a systematic review. *Br J Sports Med.* 2016;50:281–91.
42. Campbell PG, Stewart IB, Sirotic AC, Drovandi C, Foy BH, Minett GM. Analysing the predictive capacity and dose-response of wellness in load monitoring. *J Sports Sci.* 2021;39:1339–47.
43. Nobari H, Fani M, Pardos-mainer E, Pérez-gómez J. Fluctuations in well-being based on position in elite young soccer players during a full season. *Healthcare.* 2021;9(5):586.
44. Dalen T, Lorås H. Monitoring Training and Match Physical Load in Junior Soccer Players: Starters versus Substitutes. *Sports.* 2019;7:70.
45. Anderson L, Orme P, Di Michele R, Close GL, Milsom J, Morgans R, et al. Quantification of seasonal-long physical load in soccer players with different starting status from the English Premier League: Implications for maintaining squad physical fitness. *Int J Sports Physiol Perform.* 2016;11:1038–46.
46. Silva AF, Oliveira R, Cataldi S, Clemente FM, Latino F, Badicu G, et al. Weekly Variations of Well-Being and Interactions with Training and Match Intensities: A Descriptive Case Study in Youth Male Soccer Players. *Int J Environ Res Public Health.* 2022;19:1–12.
47. Robey E, Dawson B, Halson S, Gregson W, Goodman C, Eastwood P. Sleep quantity and quality in elite youth soccer players: A pilot study. *Eur J Sport Sci.* 2014;14:410–7.
48. Oliveira R, Brito JP, Martins A, Mendes B, Marinho DA, Ferraz R, et al. In-season internal and external training load quantification of an elite European soccer team. *PLoS ONE.* 2019;14:1–18.
49. Nobari H, Castillo D, Clemente FM, Carlos-Vivas J, Pérez-Gómez J. Acute, chronic and acute/chronic ratio between starters and non-starters professional soccer players across a competitive season. *Proc Inst Mech Eng Part P J Sport Eng Technol.* 2021;236(4):285–94.
50. Dalen T, Sandmæl S, Stevens TGA, Hjelde GH, Kjosnes TN, Wisløff U. Differences in Acceleration and High-Intensity Activities Between Small-Sided Games and Peak Periods of Official Matches in Elite Soccer Players. *J strength Cond Res.* 2021;35:2018–24.
51. Clemente FM, Sarmento H, Rabbani A, Van Der Linden CMIN, Kargarfard M, Costa IT. Variations of external load variables between medium- and large-sided soccer games in professional players. *Res Sports Med.* 2019;27:50–9.
52. Clemente FM, Silva R, Castillo D, Arcos AL, Mendes B, Afonso J. Weekly load variations of distance-based variables in professional soccer players: A full-season study. *Int J Environ Res Public Health.* 2020;17(9):3300.
53. Morgans R, Bezuglov E, Orme P, Burns K, Rhodes D, Babraj J, et al. The Physical Demands of Match-Play in Academy and Senior Soccer Players from the Scottish Premiership. *Sports.* 2022;10:1–11.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Ready to submit your research? Choose BMC and benefit from:

- fast, convenient online submission
- thorough peer review by experienced researchers in your field
- rapid publication on acceptance
- support for research data, including large and complex data types
- gold Open Access which fosters wider collaboration and increased citations
- maximum visibility for your research: over 100M website views per year

At BMC, research is always in progress.

Learn more biomedcentral.com/submissions

