

# Central Lancashire Online Knowledge (CLoK)

Title	Transcranial Temporal Interference Stimulation (tTIS) Influences Event-
	Related Alpha Activity During Mental Rotation
Туре	Article
URL	https://clok.uclan.ac.uk/52061/
DOI	https://doi.org/10.1111/psyp.14651
Date	2024
Citation	Thiele, Carsten, Rufener, Katherina S., Repplinger, Stefan, Zaehle, Tino and Ruhnau, Philipp (2024) Transcranial Temporal Interference Stimulation (tTIS) Influences Event-Related Alpha Activity During Mental Rotation. Psychophysiology, 61 (11). ISSN 0048-5772
Creators	Thiele, Carsten, Rufener, Katherina S., Repplinger, Stefan, Zaehle, Tino and Ruhnau, Philipp

It is advisable to refer to the publisher's version if you intend to cite from the work. https://doi.org/10.1111/psyp.14651

For information about Research at UCLan please go to <a href="http://www.uclan.ac.uk/research/">http://www.uclan.ac.uk/research/</a>

All outputs in CLoK are protected by Intellectual Property Rights law, including Copyright law. Copyright, IPR and Moral Rights for the works on this site are retained by the individual authors and/or other copyright owners. Terms and conditions for use of this material are defined in the <u>http://clok.uclan.ac.uk/policies/</u> Revised: 14 May 2024

WILEY

DOI: 10.1111/psyp.14651

### ORIGINAL ARTICLE

# Transcranial temporal interference stimulation (tTIS) influences event-related alpha activity during mental rotation

Carsten Thiele<sup>1,2</sup> | Katharina S. Rufener<sup>2,3</sup> | Stefan Repplinger<sup>1,2</sup> | Tino Zaehle<sup>1,2</sup> | Philipp Ruhnau<sup>2,4</sup>

<sup>1</sup>Department of Neurology, Otto-von-Guericke-University, University Clinic of Magdeburg, Magdeburg, Germany

<sup>2</sup>Center for Behavioral Brain Sciences (CBBS), Otto-von-Guericke-Universität Magdeburg, Magdeburg, Germany

<sup>3</sup>Department of Psychiatry, Psychotherapy and Psychosomatic Medicine of Childhood and Adolescents, Otto-von-Guericke-University, University Clinic of Magdeburg, Magdeburg, Germany

<sup>4</sup>School of Psychology and Humanities, University of Central Lancashire, Preston, UK

### Correspondence

Carsten Thiele, Medizinische Fakultät, Universitätsklinikum Magdeburg A.ö.R., Leipziger Str. 44, Magdeburg 39120, Germany. Email: carsten.thiele@med.ovgu.de

Philipp Ruhnau, School of Psychology and Humanities, University of Central Lancashire, Preston PR1 2HE, UK. Email: pruhnau@uclan.ac.uk

### Funding information

Center for Behavioral Brain Sciences (CBBS), Grant/Award Number: ZS/2016/04/78113

### Abstract

Non-invasive brain stimulation techniques offer therapeutic potential for neurological and psychiatric disorders. However, current methods are often limited in their stimulation depth. The novel transcranial temporal interference stimulation (tTIS) aims to overcome this limitation by non-invasively targeting deeper brain regions. In this study, we aimed to evaluate the efficacy of tTIS in modulating alpha activity during a mental rotation task. The effects of tTIS were compared with transcranial alternating current stimulation (tACS) and a sham control. Participants were randomly assigned to a tTIS, tACS, or sham group. They performed alternating blocks of resting and mental rotation tasks before, during, and after stimulation. During the stimulation blocks, participants received 20 min of stimulation adjusted to their individual alpha frequency (IAF). We assessed shifts in resting state alpha power, event-related desynchronization (ERD) of alpha activity during mental rotation, as well as resulting improvements in behavioral performance. Our results indicate tTIS and tACS to be effective in modulating cortical alpha activity during mental rotation, leading to an increase in ERD from pre- to poststimulation as well as compared to sham stimulation. However, this increase in ERD was not correlated with enhanced mental rotation performance, and resting state alpha power remained unchanged. Our findings underscore the complex nature of tTIS and tACS efficacy, indicating that stimulation effects are more observable during active cognitive tasks, while their impacts are less pronounced on resting neuronal systems.

PSYCHOPHYSIOLOGY

### K E Y W O R D S

event-related desynchronization, mental rotation, non-invasive brain stimulation, temporal interference stimulation, transcranial alternating current stimulation

This is an open access article under the terms of the Creative Commons Attribution License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

© 2024 The Author(s). Psychophysiology published by Wiley Periodicals LLC on behalf of Society for Psychophysiological Research.

## **1** | INTRODUCTION

Transcranial electrical stimulation (tES) is being researched as a promising therapeutic intervention for a diverse range of neurological and psychiatric disorders (Cho et al., 2022; Yang et al., 2021), including depression (Alexander et al., 2019; Brunoni et al., 2013; Wang et al., 2022), anxiety (Stein et al., 2020), dyslexia (Marchesotti et al., 2020; Rufener & Zaehle, 2021), schizophrenia (Brunelin et al., 2022; Pinault, 2017), attention deficit hyperactivity disorder (ADHD) (Salehinejad et al., 2019, 2020), Alzheimer's disease (Pini et al., 2022; Rajji, 2019), and stroke rehabilitation (Khan et al., 2022; Solomons & Shanmugasundaram, 2019). This noninvasive method involves the application of weak electric currents through electrodes attached to the scalp, modulating neuronal activity in underlying cortical regions, leading to both behavioral and electrophysiological changes (Antal et al., 2014; Guleyupoglu et al., 2013; Khadka & Bikson, 2023; Paulus et al., 2016; Vosskuhl et al., 2018; Zaghi et al., 2010).

One variant of tES, transcranial alternating current stimulation (tACS), applies weak sinusoidal currents to interact with endogenous brain oscillations via neural entrainment (Abd Hamid et al., 2015; Herrmann et al., 2016; Herrmann & Strüber, 2017). Entrainment refers to the synchronization of endogenous oscillations to the externally applied tACS signal, modulating ongoing rhythmic brain activity (Haegens et al., 2011; Reato et al., 2013; Thut et al., 2011). This interaction induces neuronal changes during stimulation (online effects) and persists poststimulation (offline effects) (Kasten et al., 2016; Veniero et al., 2015; Zaehle et al., 2010). Studies attribute these offline effects to changes in spike-timing-dependent plasticity (STDP) (Vossen et al., 2015, see Vogeti et al., 2022 for a discussion). This neuronal modulation enables tACS to affect a range of cognitive processes, such as working memory (Grover et al., 2022; Hoy et al., 2015; Pahor & Jaušovec, 2018; Reinhart & Nguyen, 2019), attention (Schuhmann et al., 2019), motor function (Wischnewski, Engelhardt, et al., 2019; Wischnewski, Schutter, & Nitsche, 2019), and speech perception (Rufener et al., 2016; Zoefel et al., 2018).

Despite the effectiveness of tACS at the cortical level, a significant limitation is its shallow stimulation depth. The electrical currents remain primarily superficial and quickly diminish in intensity when reaching deeper regions of the brain (Miranda et al., 2013; Opitz et al., 2016). Consequently, in the treatment of clinical conditions such as Parkinson's disease that arise from dysfunctions in deeper brain areas (Herrington et al., 2016; Limousin & Foltynie, 2019; Volkmann, 2004), invasively implanted electrodes remain the most viable approach for deep brain stimulation.

However, recent findings highlight the potential for a new method, which may be capable of non-invasive deep brain stimulation. Grossman et al. (2017) successfully demonstrated the feasibility of transcranial temporal interference stimulation (tTIS), showing that it could induce motor activity in rodents by stimulating cortical areas. This method simultaneously applies two high-frequency ( $\geq$ 1 kHz) sinusoidal electric fields  $\bar{E}_1$  and  $\bar{E}_2$  with a frequency offset  $\Delta f$  to the brain. The superposition of both fields creates an interference which results in a lowfrequency amplitude-modulated waveform at  $\Delta f$ , which is at the "envelope" or "beat" frequency (e.g.,  $\Delta f = 10$  Hz, generated via  $f_1 = 1000$  Hz,  $f_2 = f_1 + \Delta f = 1010$  Hz). This envelope frequency is within the receptive frequency range of neurons and leads to entrainment of endogenous oscillations, causing neuronal modulation (Cao et al., 2020; Conta et al., 2022; Esmaeilpour et al., 2021; Mirzakhalili et al., 2020; Rampersad et al., 2019; Wang et al., 2023). A key aspect of tTIS is that due to an inherent low-pass property, neurons are unable to entrain to oscillations ≥1 kHz (Hutcheon & Yarom, 2000). TTIS leverages this by intentionally setting its carrier frequency  $f_1$  above this threshold to limit its entrainment effect only to the low-frequency amplitude-modulated waveform. A further advantage of tTIS is its high focality (compared to tACS), as the locus of stimulation is limited to the area of maximum interference (i.e., where both electric fields have the same intensity). The intensity of the amplitude-modulated waveform is strongest only within this specific area, minimizing costimulation of adjacent areas and thereby enhancing the precision of the stimulation (Conta et al., 2021; Khatoun et al., 2021; Terasawa et al., 2022). Simulations and modeling approaches allow for the guidance of this interference point toward any region in the brain, including subcortical areas (Song 2019; Lee, 2021; Stoupis and Samaras 2022; Terasawa et al., 2022; Conta et al., 2021). The enhanced penetration depth is further facilitated by the higher conductance values of tissues for electric currents in the kHz range, enabling the currents to reach deeper regions (Gabriel et al., 1996). Therefore, tTIS could lay the foundation for non-invasive deep brain stimulation, potentially enabling the treatment of neurological deficits that currently require invasive procedures like deep brain stimulation (DBS) (Grossman et al., 2018).

A comparable method of electrostimulation was proposed half a century ago and tested for its capabilities in electro-anesthesia (Brown, 1975; Sachkov et al., 1967). While this method dubbed as "electrical interferential current therapy" was later used as a means to stimulate and treat muscle tissues (Goats 1990; Beatti et al. 2011), Grossman et al. (2017) were among the first to demonstrate its efficacy in cortical stimulation as well. Computational modeling studies deliver promising results, suggesting that tTIS can indeed induce neuronal entrainment (Karimi et al., 2019; Lee, 2021; Su et al., 2021; Zhu et al., 2022). But the body of in vivo research on tTIS in humans remains sparse and presents conflicting findings. Among the few studies, Ma et al. (2021) demonstrated improved motor functions after tTIS, while Zhu et al. (2022) observed an increased functional connectivity in the motor cortex. However, other studies using magnetoencephalography (MEG) (Conta et al., 2022) or electroencephalography (EEG) (Iszak et al., 2023) found no effect of tTIS on posterior  $\alpha$ -power, underscoring the need for further investigation. Consequently, our aim in this study was to gather empirical evidence on the stimulation effect of tTIS in an in vivo experiment in humans to provide a proof-of-concept.

In a recent study, we investigated the efficacy of amplitude-modulated waveforms, as used in tTIS, to cause neuronal activation. For this, we used induced retinal phosphenes as an indicator for stimulation efficacy (Thiele et al., 2021). Our results revealed that unlike tACS, amplitude-modulated waveforms failed to produce phosphenes even at intensities significantly higher than those required for tACS, which was also replicated in Iszak et al. (2023). These findings align with existing literature suggesting that amplitude-modulated waveforms necessitate greater stimulation intensities for effective neuronal activation compared to sinusoidal alternating currents, as used in tACS (Esmaeilpour et al., 2021; Rampersad et al., 2019). We concluded that although effects in the form of neuronal activation may not be achievable, the potential for neuronal modulation remains. Therefore, in the present study, we evaluated the efficacy of tTIS in humans by analyzing its capability for causing neuronal modulations.

For this, we induced and measured changes in cortical alpha activity after stimulation with tTIS. Studies on alpha oscillations generally refer many cognitive functions to alpha activity, including perception (Romei et al., 2012), intelligence (Doppelmayr et al., 2002), or top-down control (Sherman et al., 2016). The prevailing understanding of the role of alpha activity in cognition is that it directs the flow of information by gating task-irrelevant or distracting information, thereby allowing for more focused attention on task-relevant information (Foxe & Snyder, 2011; Klimesch et al., 2011). Consequently, an increase in alpha activity within a region is observed when there is active suppression of interfering information or processes. Conversely, a decrease in alpha activity is thought to facilitate information processing and cognitive performance (Foxe & Snyder, 2011; Haegens et al., 2011; Jensen

PSYCHOPHYSIOLOGY

& Mazaheri, 2010; Rihs et al., 2009; Sauseng et al., 2005; Zumer et al., 2014). TACS studies can use this link to their advantage by modulating alpha activity to enhance information processing (Schutter & Wischnewski, 2016). Studies have validated this concept using Shepard's mental rotation task (Shepard & Metzler, 1971), with tACS in the alpha range significantly improving task performance compared to sham stimulation (Kasten et al., 2018; Kasten & Herrmann, 2017). Building on these findings, the objective of this study was to explore whether tTIS could produce a similar effect. Specifically, we aimed to increase alpha power using tTIS, with the goal of enhancing performance in a mental rotation task. To differentiate between true stimulation effects and effects due to time-on-task, we included a sham stimulation group as a control. Additionally, a tACS group was included as a benchmark of stimulation effects using this paradigm (Kasten et al., 2018; Kasten & Herrmann, 2017) allowing for a direct comparison of the stimulation efficacy of tTIS and tACS.

We quantified the stimulation effect by calculating event-related spectral perturbation (ERSP) (Makeig et al., 2004; Pfurtscheller & Aranibar, 1977; Pfurtscheller & Da Silva, 1999) in the alpha band (8-13 Hz). ERSP measures the change in spectral power, or oscillatory activity, triggered by an event, like the presentation of a visual stimulus. This measurement distinguishes between two phenomena: an increase in spectral power, known as event-related synchronization (ERS); and a decrease in spectral power, referred to as event-related desynchronization (ERD). Klimesch et al. (2007) argue that an ERD in the alpha band is associated with a release of inhibitory processes, allowing for subsequent neuronal activation and facilitation of information processing (see also Pfurtscheller, 1997). This hypothesis has been supported by studies using transcranial magnetic stimulation (TMS) (Klimesch et al., 2003) and more recently, by studies using tACS to increase ERD and improve performance in a mental rotation task (Kasten et al., 2018; Kasten & Herrmann, 2017). However, conversely, some studies not using brain stimulation techniques have instead reported improved performance with reduced ERD, postulating a neural efficiency hypothesis, which suggests that skilled individuals use fewer brain resources during task performance (Chen et al., 2013; Riečanský & Katina, 2010). Given our use of brain stimulation techniques, we hypothesize our results to be in line with those found in the studies of Kasten et al. (2018; 2017), expecting an ERD increase in tTIS and tACS groups compared to the sham group, as well as a corresponding improvement in mental rotation performance.

Numerous studies have demonstrated the neuronal modulation effect of tACS and its associated increase in alpha power (Berger et al., 2018; Kasten et al., 2016; Veniero et al., 2015; Zaehle et al., 2010). Considering that

a mental rotation task naturally leads to significant alpha modulations by suppressing alpha activity during task execution, this could potentially disrupt the stimulation's entrainment effect. To measure the stimulation effect without the confounding influence of a complex cognitive task, we additionally included resting blocks utilizing a simple vigilance task, a setup also utilized by other studies (Conta et al., 2022; Kasten et al., 2018; Kasten & Herrmann, 2017; Zaehle et al., 2010). We hypothesized a significantly stronger increase in resting state alpha power from pre- to poststimulation in the tTIS and tACS groups compared to the sham group.

## 2 | METHODS

## 2.1 | Participants

Given that no studies on the effects of tTIS on mental rotation have been conducted, we could only approximate the required sample size. We based our estimate on a previous study by Kasten and Herrmann (2017), which reported an effect size of  $\eta^2 = 0.27$  for two groups (stimulation vs. sham). Using these parameters, we conducted a power analysis with G\*Power (Ver. 3.1.9.7, Faul et al., 2007), setting an alpha error probability of 0.05 and a power of 0.95 for a ttest between two independent means. This analysis suggested a required sample size of 16 participants per group. While we recognize that this is only an approximate solution, it provided a reasonable starting point for our research. In total, we recruited 67 participants who were randomly assigned to either the tTIS, tACS, or sham group. Of the 67 participants, 48 were eligible for data analysis. This was, on the one hand, due to technical issues (a coding error in the Matlab stimulation function) leading to the exclusion of nine individuals from the data analysis. Additionally, 10 participants were excluded due to the absence of a discernible peak in the alpha band (8-13 Hz) during the first block of the resting task, which was crucial to extract the individual alpha frequency (IAF) (see Electrical Stimulation) for subsequent stimulation. As a result, our analysis included data from 48 participants, divided into three stimulation conditions: tACS (n=18; 11 female, 7 male; mean age = 23.7, SD age = 4.52), tTIS (n = 16; 10 female, 6 male; mean age = 21.8, SD age = 2.46),and sham (n = 14; 11 female, 3 male; mean age = 22.9, SDage = 2.85).

Eligible participants for this study had to meet the following criteria: no history of epileptic seizures or psychiatric or neurological disorders, no metal or electric implants in their body, free of medication affecting the central nervous system, non-smoking, and normal or corrected-tonormal vision. Before the experiment, participants were informed about the experimental procedure and the potential adverse effects of electrostimulation and were required to give written informed consent. This study was approved by the local ethics committee of the University Clinic of Magdeburg and conducted in accordance with the guidelines of the Declaration of Helsinki.

## 2.2 | EEG

EEG data were acquired using passive Ag-AgCl electrodes (EasyCap, Brain Products, Gilching, Germany) positioned at Fz, Pz, P7, and P8 following the international 10–20 system. The reference electrode was placed on the tip of the participant's nose, while the ground electrode was positioned at AFz. In addition, electrodes were placed vertically (VEOG) and horizontally (HEOG) to the right eye and referenced to the nose electrode, to control for eye movements and eyeblinks. To increase electrode to skin conductivity, we applied a conductive paste (Abralyt 2000 abrasive electrolyte-gel, Brain Products, Gilching, Germany), ensuring impedances remained below  $5 \text{ k}\Omega$ . Data were sampled at a rate of 1000 Hz using a BrainAmp DC amplifier (Brain Products, Gilching, Germany).

## 2.3 | Electrical stimulation

We used a battery-operated stimulator system (DC-Stimulator Plus, NeuroConn GmbH, Ilmenau, Germany) connected to a PC via a digital-to-analog converter (NI USB-6212, National Instruments, Austin, TX, USA) to generate and send the stimulation signal through a custom Matlab (version 2020a, Mathworks, Natick, USA) script. The stimulation was administered using surface conductive rubber electrodes (NeuroConn GmbH, Ilmenau, Germany) attached to the subject's head using an adhesive, electrically conductive paste (Ten20, D.O. Weaver, Aurora, CO, USA). Electrode impedances were kept below 5 k $\Omega$  to ensure optimal conductivity.

As studies suggest that stimulation in the alpha band is much more effective if the stimulation frequency matches the participants' IAF (Huang et al., 2021; Kasten et al., 2019; Schutter & Wischnewski, 2016), we determined the IAF based on the EEG data recorded in the first resting block at EEG electrode Pz and used it as the stimulation frequency.

For the tACS and sham groups, we used a parietooccipital electrode montage consisting of a  $5 \times 7$  cm electrode placed over Cz and a  $4 \times 4$  cm electrode placed over Oz, with a stimulation intensity of 1 mA peak-topeak. This montage has shown to increase power in the alpha range in posterior areas (Kasten et al., 2019) and to improve performance in a mental rotation task (Kasten et al., 2018; Kasten & Herrmann, 2017). Using SimNIBS 3.0 (Thielscher et al., 2015), we simulated the electric field of this montage to confirm sufficient electric field strengths (>0.2 V/m) were being delivered to the targeted posterior brain areas (see Figure 1c Left). Participants in the tACS group received 20 min of stimulation in total, split between the Rest<sub>Stimulation</sub> and Rotation<sub>Stimulation</sub> blocks, whereas those in the sham group received stimulation for only 30 s at the beginning of a stimulation block. Stimulation intensities were ramped up and down for 10 s at the beginning and end, respectively.

For tTIS, we used a custom SimNIBS script based on the formula proposed by Grossman et al. (2017) to determine a tTIS montage targeting approximately the same posterior regions as our tACS montage. Our simulations indicated that two pairs of two round electrodes (34mm diameter) each allow for stimulation of posterior brain areas (see Figure 1c Right). The first pair of electrodes was positioned with one electrode on P4 and the other between I1 and O1. The second pair was arranged with one electrode on P3 and the other placed between I2 and O2. Since tTIS requires a higher stimulation intensity to achieve a comparable stimulation efficacy as tACS (Esmaeilpour et al., 2021), we used a stimulation intensity of 2mA peak-to-peak for stimulation with tTIS. This is still a safe (Cassarà et al., 2022; Piao et al., 2022) and tolerable stimulation intensity, as tTIS uses high-frequency stimulation, to which somatosensory perception is less sensitive (Fertonani et al., 2015; Hsu et al., 2021; Zeng et al., 2019) and thus can be tolerated by participants without adverse effects. Anecdotally, many participants in our study reported not feeling tTIS at all. We selected a carrier frequency of 1 kHz balancing the need to maximize stimulation efficacy and minimizing unintended stimulation effects related to the carrier frequency (Esmaeilpour et al., 2021; Grossman et al., 2017; Rampersad et al., 2019). For the envelope frequency, we used the IAF. Using tTIS, the envelope is determined by the difference in stimulation frequencies of  $E_1$  and  $E_2$ . This resulted in stimulation frequencies of  $f_1 = 1000 \text{ Hz}$  and  $f_2 = 1000 \text{ Hz} + \text{IAF}$ . As was the case for tACS, the stimulation duration for tTIS was 20 min in total, split between the Rest<sub>Stimulation</sub> and Rotation<sub>Stimulation</sub> blocks, with the stimulation being ramped up at the beginning and down at the end over a period of 10s.

## 2.4 Data analysis

The acquired electrophysiological data were processed and analyzed using Matlab 2020a (The MathWorks Inc., PSYCHOPHYSIOLOGY SPR

Natick, MA, USA) in conjunction with the Fieldtrip toolbox (Oostenveld et al., 2011). Statistical analysis was performed using JASP version 0.17.1 (JASP Team, 2023) and Jamovi version 2.3 (The Jamovi project 2022).

To assess behavioral performance in the mental rotation task, measures of task accuracy, that is, correct answers and reaction time (RT), were analyzed. Oneway analyses of variance (ANOVAs) were conducted on accuracy and RT of the Rotation<sub>Baseline</sub> Block, with the between-subject factor Stimulation (tTIS, tACS, sham) to identify potential baseline differences. Subsequently, group differences in task performance before, during, and after stimulation were examined using repeated measures ANOVAs (rmANOVAs). The rmANOVAs included the within-subject factor Block (Rotation<sub>Baseline</sub>, Rotation<sub>Stimulation</sub>, Rotation<sub>Poststim</sub>) and the betweensubject factor Stimulation (tTIS, tACS, sham). Further, to analyze the effects of stimulation on behavioral performance and adjust for individual baseline differences, we've calculated change measures  $\Delta$ Accuracy (Accuracy in Rotation<sub>Poststim</sub>-Accuracy in Rotation<sub>Baseline</sub>) and  $\Delta RT$  (RT in Rotation<sub>Poststim</sub> – RT in Rotation<sub>Baseline</sub>). This shows whether performance increased or decreased, comparing the pre- and poststimulation performance, to help us understand changes due to stimulation, regardless of initial performance levels. Change measures,  $\Delta$ Accuracy and  $\Delta$ RT, were then analyzed using separate ANOVAs using between-subject factor Stimulation (tTIS, tACS, sham).

In our ANOVAs, we report both partial eta squared  $(\eta_p^2)$ , which shows the variance explained in our model but may overestimate effects, and generalized eta squared  $(\eta_G^2)$ , which offers consistent effect sizes across studies but is less specific to our design. This approach balances detailed insights with broader comparability.

For EEG analysis of alpha activity, we focused on electrode Pz for the analysis similar to previous studies (Kasten et al., 2016; Kasten & Herrmann, 2017; Zaehle et al., 2010). To analyze changes in resting state alpha activity, data of all stimulation-free resting blocks (Rest<sub>Baseline</sub>, Rest<sub>Poststim 1</sub>, Rest<sub>Poststim 2</sub>, Rest<sub>Poststim 3</sub>) were analyzed. EEG data collected during the Rest<sub>Stimulation</sub> block could not be analyzed due to significant stimulation artifacts. Blocks were epoched into 1-second non-overlapping segments. Segments containing major artifacts (e.g., muscle artifacts) were removed after visual inspection via fieldtrips ft\_rejectvisual function. Afterward, power in the IAF band (IAF +/-1Hz) was calculated by performing fast Fourier transforms (FFTs) using a Hanning window and 2-second zero padding. In addition, using the fitting oscillations & oneover f (FOOOF) method (Donoghue et al., 2020), data were split into periodic and aperiodic components to



**FIGURE 1** Experimental design. (a) Experimental Timeline. Chronological sequence of events during the experiment. Participants began with a resting block, followed by alternating blocks of mental rotation and resting blocks. Stimulation was administered during the second resting block and the second mental rotation block (indicated with a lightning symbol). (b) Example Task Set. Illustration of example task sets used in the study. Geometric figures, based on Shepard's mental rotation task, were presented to participants. They were required to determine whether the figures were mirrored or not. (c) Electrode Montages. Depending on the (random) group assignment, participants received either tACS, sham, or tTIS. (d) Stimulation Signal Exemplified. The figure demonstrates the stimulation signals used in the study. On the left side, 10 Hz tACS sine waves are compared to a tTIS amplitude-modulated waveform with a 10 Hz envelope frequency. The right side provides a zoomed-in view of the waveforms, highlighting the high-frequency carrier component of tTIS.

distinguish rhythmic activity from concurrent powerspectral 1/f modulations. For data analysis, only periodic components were analyzed, as these reflect frequencyspecific rhythmic activity and are free of power changes affecting all frequency bands, which can be caused, for instance, by technical issues such as changes in electrode impedances. The periodic component in the IAF Band was then analyzed using an rmANOVA with the within-subject factor Block (Rest<sub>Baseline</sub>, Rest<sub>Poststim 1</sub>, Rest<sub>Poststim 2</sub>, Rest<sub>Poststim 3</sub>) and between-subject factor Stimulation (tTIS, tACS, sham) to detect changes in resting state alpha power. Additionally, we calculated a change measure,  $\Delta Alpha_{Rest}$  (=Alpha power in Rest<sub>Poststim3</sub> – Alpha power in Rest<sub>Baseline</sub>), and conducted an ANOVA with the between-subject factor Stimulation (tTIS, tACS, sham). This analysis focused specifically on increases in alpha activity, independent of individual alpha levels.

To capture changes in the IAF-band during the mental rotation task, ERSP was calculated for baseline and poststim mental rotation blocks. Again, data acquired during the stimulation block could not be analyzed due to the stimulation artifact. Data in the mental rotation blocks were segmented into 10-second epochs, spanning from 3 s before to 7 s after stimulus presentation. Epochs containing major artifacts were rejected after visual inspection using fieldtrips *ft\_rejectvisual* function. Remaining epochs were analyzed using FFTs with a Hanning-tapered sliding window with a fixed length of 1 s moving in steps of 50 ms along each segment and 2-second zero padding (as in Kasten & Herrmann, 2017). Afterward, ERSP values were calculated as:

$$\text{ERSP}_{\text{t}} = \frac{R_{\text{t}} - A_{\text{t}}}{R_{\text{t}}} \times 100$$

In this formula, R is defined as the power during a reference period, specifically before the stimulus is presented. In our study, R was the power in the IAF band (IAF +/-1 Hz) calculated from -2 to 0s relative to stimulus onset. On the other hand, A is defined as the power of the frequency of interest during a test period, which is after the stimulus presentation. For A, we defined it as the power in the IAF band from 0 to 3s following stimulus onset. Furthermore, t is defined as the trial number, as we calculated the ERSP on a trial-by-trial basis, based on prior studies (Kasten & Herrmann, 2017; Nakayashiki et al., 2014). To acquire an average ERSP for a block, ERSPs of all trials were averaged in each block for each subject. Resulting positive values indicate an ERD, reflecting a drop in power in the frequency band of interest at stimulus onset, whereas negative values indicate an ERS, reflecting an increase in power at stimulus onset. Based on previous PSYCHOPHYSIOLOGY SPR

work (Kasten et al., 2018; Kasten & Herrmann, 2017), we expected to observe an ERD. To explore potential differences in baseline ERD between stimulation groups, the ERD in the baseline block was analyzed using an ANOVA with between-subject factor *Stimulation* (tTIS, tACS, sham). ERD values were then fed into an rmANOVA with within-subject factor *Block* (Rotation<sub>Baseline</sub>, Rotation<sub>Poststim</sub>) and between-subject factor *Stimulation* (tTIS, tACS, sham). Again, a change measure  $\Delta$ ERD (ERD in Rotation<sub>Poststim</sub> – ERD in Rotation<sub>Baseline</sub>) was calculated and subjected to an ANOVA with between-subject factor *Stimulation* (tTIS, tACS, sham) to titrate changes in ERD from pre- to poststimulation.

Finally, we investigated if changes in task performance were correlated with changes in ERD values. To achieve this, we correlated  $\Delta$ ERD with  $\Delta$ Accuracy and  $\Delta$ RT. Prior to further analysis, we confirmed that all variables exhibited normal distribution based on the Shapiro–Wilk test ( $\Delta$ ERD: W=0.985, p=.796;  $\Delta$ Accuracy: W=0.972, p=.31;  $\Delta$ RT: W=0.973, p=.337). Subsequently, we performed Pearson's correlational analysis to examine the potential influence of changes in ERD on changes in task performance measures.

For rmANOVA results we report partial eta squared  $(\eta_p^2)$  to focus on effect size within our chosen design, as well as generalized eta squared  $(\eta_G^2)$  to facilitate comparing effect sizes across studies.

## 3 | RESULTS

## 3.1 | Experimental design

The experimental tasks were displayed on a Samsung SyncMaster SA450 placed at a distance of approximately 100 cm from the participants. For the experimental presentation, we utilized Psychtoolbox 3 (Kleiner et al., 2007) which was implemented using Matlab 2020a (The MathWorks Inc., Natick, MA, USA).

During the experiment, participants engaged in alternating resting blocks and mental rotation blocks (see Figure 1a). During the resting blocks (Rest<sub>Baseline</sub>, Rest<sub>Stimulation</sub>, Rest<sub>Poststim 1</sub>, Rest<sub>Poststim 2</sub>, Rest<sub>Poststim 3</sub>), participants engaged in a simple vigilance task as employed by previous studies (Kasten & Herrmann, 2017; Zaehle et al., 2010). In this task, participants were shown a fixation cross at the center of the screen, which could rotate by 45 degrees for a duration of 500 ms. The onset of the rotation varied, occurring randomly between 30 and 40 s. Upon noticing the rotation, participants were required to press a response button within a 2-second window following the stimulus onset. This task served a dual purpose: maintaining the subjects' attention and facilitating the

collection of clean EEG data. Resting blocks consisted of a 4-minute vigilance task followed by a 1-minute break. The exception was the Rest<sub>Stimulation</sub> block, whose duration was doubled to 10 min to accommodate the necessary 10-minute stimulation period.

(Rotation<sub>Baseline</sub>, In mental rotation blocks Rotation<sub>Stimulation</sub>, Rotation<sub>Poststim</sub>), participants performed a Shepard's mental rotation task (Shepard & Metzler, 1971) which involved presenting two geometrical figures (see Figure 1b) that could be (1) rotated and/ or (2) mirrored in relation to each other. Participants were tasked with mentally rotating the figures to matching angles, in order to determine whether the figures were mirrored or not. The figures could be rotated relative to each other by 0°, 50°, 100°, or 150°, with larger rotation angles increasing the task's difficulty. Participants indicated their responses by pressing the "J" key for mirrored trials and the "n" key for non-mirrored trials. Each rotation block comprised 48 trials, with 12 trials per rotation angle. Half of the trials contained mirrored figures. A trial began with the presentation of a fixation cross in the middle of the screen for 3s, followed by the presentation of the geometric figures which lasted for 7s. Participants were instructed to answer as fast and accurately as possible. A mental rotation block consisted of a 4-minute task, followed by a 1-minute break, then another 4-minute task, and a concluding 1-minute break. The stimuli for the task were selected from a published open-source stimulus set (Ganis & Kievit, 2015). During the Rotation<sub>Stimulation</sub> block, participants received 10 min of stimulation.

At the beginning of the experiment, participants completed a questionnaire to assess exclusion criteria and were informed about the planned experimental procedure and tasks. Following this, the montages for electrical stimulation (see Figure 1c) and EEG were set up. Participants then engaged in the alternating blocks of resting and mental rotation tasks. Depending on the group, electrical stimulation of either tTIS, tACS, or sham (see Figure 1d) was administered in the second resting block and in the second rotation task block (see Figure 1a). At the end of the experiment, participants filled out a questionnaire about possible adverse effects of the stimulation (nausea, headache, sensations of pain/burning/itching/reddening at the stimulated area) (Brunoni et al., 2011). Afterward, participants were debriefed about the aim of this study and reimbursed for their time, either with course credit or monetarily.

## 3.2 | Resting-state alpha power

Analysis of resting-state alpha power revealed a main effect *Block* [F(3,135) = 22.27, p < .001,  $\eta_p^2 = 0.331$ ,  $\eta_G^2 = 0.051$ ]. Subsequent post-hoc *t* tests indicated an increase in

alpha power throughout the experiment [Rest<sub>Baseline</sub> vs. Rest<sub>Poststim 1</sub>, Rest<sub>Baseline</sub> vs. Rest<sub>Poststim 2</sub>, Rest<sub>Baseline</sub> vs. Rest<sub>Poststim 3</sub>, Rest<sub>Poststim 1</sub> vs. Rest<sub>Poststim 2</sub>, Rest<sub>Poststim 1</sub> vs. Rest<sub>Poststim 3</sub>: all  $t(45) \ge 3.263$ ,  $p_{tukey} \ge 0.011$ ], with the exception of the comparison between blocks Rest<sub>Poststim 2</sub> and Rest<sub>Poststim 3</sub> [ $t(45) \ge 0.671$ ,  $p_{tukey} = 0.908$ ] (see Figure 2a). Neither a significant main effect of *Stimulation* on resting state alpha power [F(2,45)=1.19, p=.314,  $\eta_p^2=0.050$ ,  $\eta_G^2=0.045$ ] nor an interaction between *Stimulation* × *Block* [F(6,135)=0.931, p=.475,  $\eta_p^2=0.040$ ,  $\eta_G^2=0.004$ ] were observed. The ANOVA on  $\Delta$ Alpha<sub>Rest</sub> did not reveal a main effect *Stimulation* [F(2,45)=1.15, p=.327,  $\eta_p^2=0.048$ ]. In sum, these findings indicate that alpha power did increase over the course of the experiment, but this increase did not differ between stimulation groups.

### 3.3 Mental rotation accuracy

The analysis of baseline accuracy in the Rotation<sub>Baseline</sub> block did not reveal a statistically significant main effect for Stimulation [F(2,45)=1.38, p=.263,  $\eta_p^2=0.058$ ], indicating comparable baseline performance between groups. The rmANOVA across all mental rotation blocks (see Figure 2b) demonstrated a significant main effect Block  $[F(2,90)=7.14, p<.001, \eta_p^2=0.137, \eta_q^2=0.046]$ . Posthoc tests revealed this to be due to a statistically significant improvement in accuracy from Rotation<sub>Baseline</sub> to Rotation<sub>Poststim</sub> [ $t(45) = 3.77, p_{tukey} = .001$ ] but no statistically significant difference in accuracy between Rotation<sub>Baseline</sub> and Rotation<sub>Stimulation</sub> [t(45) = 1.96,  $p_{tukey} = .135$ ] or between Rotation<sub>Stimulation</sub> and Rotation<sub>Poststim</sub> blocks [t(45)=1.81,  $p_{\text{tukev}}$  = .176]. We found no main effect *Stimulation* on accuracy [F(2,45)=1.27, p=.291,  $\eta_p^2=0.053$ ,  $\eta_G^2=0.038$ ] nor a Block  $\times$  Stimulation interaction [F(4,90)=1.34, p=.260,  $\eta_{\rm p}^2 = 0.056, \eta_{\rm G}^2 = 0.018$ ]. For descriptive results of the mental rotation accuracy, see Table 1. The analysis of  $\Delta$ Accuracy did not reveal a main effect *Stimulation* [F(2,45)=0.653, $p = .525, \eta_p^2 = 0.028$ ], indicating that all groups experienced comparable increases in task accuracy.

Notably, accuracy levels in the Rotation<sub>Poststim</sub> block approached the ceiling across all stimulation groups, offering minimal room for observable improvements attributable to stimulation. This suggests that the lack of significant stimulation effects could be attributed to ceiling effects, limiting the ability to draw conclusions about the impact of stimulation on task accuracy. Collectively, the findings suggest that while accuracy on the mental rotation task significantly increased throughout the experiment, the rate of this improvement did not vary across the different stimulation groups. Therefore, it is most plausible that the observed improvement stems from a training effect rather than the stimulation itself.

-10

-25



**FIGURE 2** Overview of main results. (a) Mental rotation accuracy improves from the Baseline to the Poststim block, with no statistically significant differences observed between stimulation groups. (b) Similarly, reaction time in the mental rotation task shows improvement from the Baseline to the Poststim block. (c) In resting blocks, alpha power increases over the course of the experiment from Baseline to Poststim 2, where alpha activity plateaus and does not further increase in Poststim 3. No significant differences were observed between stimulation groups. (d) In the mental rotation task, ERD values increased from Baseline to Poststim blocks for tTIS and tACS groups, while remaining stable for the sham group. (e, f) Correlation analysis reveals no significant correlation between changes in ERD and changes in task accuracy (e) or RT (f).

50

25

Ó

ΔERD (%)

-25

Ó

ΔERD (%)

50

25

9 of 20

tTIS

tACS

Sham

Stimulation

## .....

TABLE 1 Desci

OPHYSIOLOG	SPR	ся.									
riptive statistics	of behavio	oral data du	aring mental rot	ation.							
	Accura	Accuracy (%)				RT (s)					
Block	Mean	SD	Minimum	Maxi	mum	Mean	SD	Minimu	ım N	laximum	
Baseline	91.1	7.70	70.8	97.9		2.64	0.59	1.75	4.	.06	
Stimulation	90.9	6.97	70.8	100.0		2.24	0.62	1.29	3.	.92	
Poststim	93.7	4.37	83.3	100.0		2.22	0.62	1.33	3.	3.71	
Baseline	87.2	7.15	75.0	95.8		3.18	0.68	1.78	4.	4.18	
Stimulation	89.8	7.39	68.8	100.0		2.91	0.76	1.58	4.	.21	
Poststim	91.9	4.89	81.3	97.9		2.60	0.55	1.67	3.	.76	
Baseline	90.3	7.43	75.0	100.0		2.82	0.60	1.79	3.	.89	
Stimulation	93.6	3.32	85.4	97.9		2.56	0.61	1.52	3.	.75	
Poststim	92.7	3.98	83.3	95.8		2.51	0.58	1.43	3.	.51	
m, poststimulatio	n, <i>SD</i> =stan the signific	dard deviati cant <i>Block</i> >	on. < <i>Stimulation</i> in	teractior	n effect or	n RT durin	g mental	rotation.			
imulation		Block	Stimulat	ion	Mean	diff.	SE	df	t	$p_{\mathrm{tukey}}$	
IS	-	Stim	tTIS		0.40		0.08	45	5.27	<.001	
IS	-	Poststim	tTIS		0.42		0.09	45	4.59	0.001	
IS	-	Poststim	tTIS		0.02		0.08	45	0.23	1.000	
CS		Stim	+1.05		0.27		0.07	45	2 0 2	0.011	

Abbreviations: Poststi

TABLE 2 Post-

Block	Stimulation		Block	Stimulation	Mean diff.	SE	df	t	<b>p</b> <sub>tukey</sub>
Baseline	tTIS	-	Stim	tTIS	0.40	0.08	45	5.27	<.001
Baseline	tTIS	-	Poststim	tTIS	0.42	0.09	45	4.59	0.001
Stim	tTIS	-	Poststim	tTIS	0.02	0.08	45	0.23	1.000
Baseline	tACS	-	Stim	tACS	0.27	0.07	45	3.82	0.011
Baseline	tACS	-	Poststim	tACS	0.58	0.09	45	6.80	<.001
Stim	tACS	-	Poststim	tACS	0.31	0.07	45	4.26	0.003
Baseline	Sham	-	Stim	Sham	0.27	0.08	45	3.31	0.044
Baseline	Sham	-	Poststim	Sham	0.32	0.10	45	3.27	0.048
Stim	Sham	-	Poststim	Sham	0.05	0.08	45	0.60	1.000

Abbreviations: df, degrees of freedom; mean diff., mean difference; Poststim, poststimulation; ptukey. p-value with Tukey correction; SE, standard error; t, t value.

#### 3.4 Mental rotation reaction time

Analysis of baseline differences of RT in the Rotation<sub>Baseline</sub> block revealed a statistically significant main effect Stimulation [F(2,45)=3.31, p=.046,  $\eta_p^2=0.128$ ]. Post-hoc ttests indicated that reaction times in the tACS group were significantly slower than in the tTIS group [t(45)=2.52, $p_{\text{tukey}} = .040$ ], but no differences were found between tTIS and sham  $[t(45)=0.81, p_{tukey}=.70]$  or tACS and sham groups  $[t(45)=1.60, p_{tukey}=.255]$ . Since this was RT in the Rotation<sub>Baseline</sub> block, where no stimulation had yet been administered, the observed differences are likely a random effect due to randomly assigning slower participants to the tACS group, faster subjects to the tTIS group, or a combination of both scenarios.

Comparing RT between all mental rotation blocks (see Figure 2c), we observed a main effect Block on RT  $[F(2,90)=45.67, p<.001, \eta_p^2=0.504, \eta_G^2=0.083]$ with post-hoc tests indicating an improvement in RT from Rotation<sub>Baseline</sub> to Rotation<sub>Stimulation</sub> [t(45) = 7.13, Rotation<sub>Baseline</sub> Rotation<sub>Poststim</sub>  $p_{\rm tukev} < .001],$ to

 $[t(45)=8.33, p_{tukey}<.001]$ , and Rotation<sub>Stimulation</sub> to Rotation<sub>Poststim</sub> [t(45)=2.81,  $p_{tukey}=.020$ ]. Further, a main effect Stimulation was found [F(2,45)=3.32, p=.045, $\eta_{\rm p}^2 = 0.129, \ \eta_{\rm C}^2 = 0.118$ ]. Post-hoc analysis revealed this to be due to the tACS group having significantly slower RTs compared to the tTIS group [t(45)=2.57,  $p_{tukey}=.035$ ], which is in line with the found baseline difference between these groups. No difference was found between tTIS and sham  $[t(45)=1.19, p_{tukey}=.464]$  or tACS and sham groups [t(45)=1.26,  $p_{tukev}=.426$ ]. Additionally, we observed a significant Block × Stimulation interaction on RT [F(4,90)=2.56, p=.044,  $\eta_p^2=0.102$ ,  $\eta_G^2=0.010$ ]. Post-hoc tests revealed that all stimulation groups exhibited improvements in RT from the Rotation<sub>Baseline</sub> to the Rotation<sub>Stimulation</sub> block, as well as from the Rotation<sub>Baseline</sub> to the Rotation<sub>Poststim</sub> block (see Table 2 for full post-hoc results). However, the tTIS and sham groups reached a plateau in RT improvement during the Rotation<sub>Stimulation</sub> block and did not show further improvement from Rotation<sub>Stimulation</sub> to Rotation<sub>Poststim</sub>. In contrast, the tACS group displayed continued RT improvement even after stimulation, showing further enhancements in the

Rotation<sub>Poststim</sub> block. For descriptive results of the RT, see Table 1. The analysis of  $\Delta$ RT did not reveal a main effect *Stimulation* [*F*(2,45)=2.20, p=.123,  $\eta_p^2$ =0.089], meaning no differential decrease in RT between stimulation groups could be observed.

Taken together, reaction times significantly improved over the course of the experiment, with RT in the tTIS and sham groups reaching a plateau during stimulation, while the tACS group continued to improve even poststimulation. It is important to note that the tACS group initially had the slowest RT, thereby having the greatest potential for improvement. This suggests that the difference observed between the tACS group and both the tTIS and sham groups could be attributed to the latter groups reaching an RT ceiling earlier in the Rotation<sub>Stimulation</sub> block, whereas the tACS group may have required the duration of the Rotation<sub>Poststim</sub> block to catch up.

## 3.5 | Mental rotation ERD

The analysis of baseline differences in ERD between stimulation groups did not reveal significant differences [F(2,45)=2.34, p=.108,  $\eta_p^2=0.094$ ]. The analysis of changes in ERD across mental rotation blocks revealed a main effect *Block* [F(1,45)=16.15, p<.001,  $\eta_p^2=0.264$ ,  $\eta_G^2=0.017$ ]. This was due to an increase in ERD from the Rotation<sub>Baseline</sub> to the Rotation<sub>Poststim</sub> block (see Figure 2d). This indicates a notable shift in task-relevant oscillatory activity. Though there was no statistically significant main effect *Stimulation* on ERD [F(2,45)=1.41, p=.254,  $\eta_p^2=0.059$ ,  $\eta_G^2=0.056$ ], we observed a significant *Stimulation*×*Block* interaction [F(2,45)=4.80, p=.013,  $\eta_p^2=0.176$ ,  $\eta_G^2=0.010$ ]. Post-hoc *t*tests revealed this interaction to be driven by an increase in ERD from the Rotation<sub>Baseline</sub> to the Rotation<sub>Poststim</sub> block for the

PSYCHOPHYSIOLOGY

tTIS  $[t(45)=3.99, p_{tukey}=.003]$  and tACS [t(45)=3.47, $p_{\text{tukev}}$  = .014] groups. In contrast, the sham group did not exhibit a change in ERD [t(45)=0.251,  $p_{tukey}=1.00$ ] (see Figure 3). To further titrate if the increases in ERD differed between stimulation groups, the change measure  $\Delta$ ERD was analyzed. This revealed a significant difference among the stimulation groups [F(2,45)=4.80, p=.013, $\eta_{\rm p}^2 = 0.176$ ], as was expected based on the significant interaction effect in the prior rmANOVA. Post-hoc analysis revealed that both the tTIS [t(45)=2.91,  $p_{tukev}=.015$ ] and tACS [t(45)=2.48,  $p_{tukey}=.043$ ] groups experienced significantly greater ERD increases compared to the sham group, with no notable difference between tTIS and tACS  $[t(45)=0.52, p_{tukey}=.861]$ . This suggests that both verum stimulations showed increased ERDs compared to sham, while no significant differences between verum stimulation conditions were evident.

Additionally, we performed a correlational analysis to determine if an increase in ERD values corresponded with an improvement in behavioral performance. This did not reveal a significant correlation between  $\Delta$ Accuracy and  $\Delta$ ERD [r(46)=.095, p=.259] (see Figure 2e), nor between  $\Delta$ RT and  $\Delta$ ERD [r(46)=.021, p=.556] (see Figure 2f). This leads to the conclusion that the change in ERD was not accompanied by a change in behavioral performance.

## 3.6 Exploratory analyses

Analyzing only ERDs has the disadvantage that one key information is missing: As the ERD is itself a ratio between the power in a reference period (i.e., the time period before stimulus presentation) and test period (i.e., the time period after stimulus presentation), it remains unclear if a rise in ERD is driven by a rise in power in the reference period, a loss of power in the test period or a combination



**FIGURE 3** Time-Frequency Representations (TFRs) depicting changes in ERD: Contrasted is power at the IAF between the Baseline and the Poststim block. TFRs were aligned at the IAF and averaged across participants within each stimulation group. The reference period for relative baseline correction was defined as the range from -2 to 0 before stimulus onset, as indicated by the white bar. The verum stimulation groups (tTIS, tACS), exhibit a distinct decrease in power around the IAF in the poststim block compared to the baseline block, resulting in increased ERD. The sham group does not exhibit this change in ERD.

of both. To test this, alpha power (IAF+-1Hz) was extracted for each trial for the reference period (2 to 0s before stimulus presentation) and test period (0 to 3s after stimulus presentation) and averaged for each block. Subsequently, we calculated changes in task alpha activity based on the Rotation<sub>Baseline</sub> and the Rotation<sub>Poststim</sub> blocks ( $\Delta$ Alpha=Alpha<sub>Poststim</sub> – Alpha<sub>Baseline</sub>) separately for both the reference period and the test period. Then, we performed correlational analyses between  $\Delta$ ERD and  $\Delta$ Alpha separately for the reference and the test period. This revealed a trend for a positive correlation between  $\Delta$ Alpha in the reference period with  $\Delta$ ERD [r(46)=0.279, p=.055] (see Figure 4b), but no significant correlation between

THIELE ET AL.

 $\Delta$ Alpha in the test period with  $\Delta$ ERD [r(46) = -0.181, p = .217] (see Figure 4a). This suggests that the changes in ERD are more likely due to power changes in the reference period than in the test period.

To further titrate stimulation effects on  $\Delta$ Alpha<sub>Rotation</sub>, we fed it into an rmANOVA with the factors *Period* (Reference, Test) and the between-subject factor *Stimulation* (tACS, tTIS, sham). This revealed a significant main effect *Period* [F(1,45)=6.10, p=.017,  $\eta_p^2=0.119$ ,  $\eta_G^2=0.052$ ] which is based on a significantly higher alpha increase in the reference period than in the test period (see Figure 4c). However, neither the main effect *Stimulation* [F(2,45)=1.36, p=.267,  $\eta_p^2=0.057$ ,  $\eta_G^2=0.035$ ] nor the



**FIGURE 4** Overview of Exploratory Results. (a, b) Correlational analysis of ERD and alpha activity changes. Analysis focused on the difference in ERD ( $\Delta$ ERD) and alpha activity changes ( $\Delta$ Alpha = Alpha<sub>Poststim</sub> – Alpha<sub>Baseline</sub>) during (a) the test period (0 to 3 s after stimulus presentation) and (b) the reference period (–2 to 0 s before stimulus presentation). Results indicated no significant correlation between  $\Delta$ Alpha in the test period and  $\Delta$ ERD, but a trending significance was observed between  $\Delta$ Alpha in the reference period and  $\Delta$ ERD. (c) Analysis of alpha changes during mental rotation split between reference and test periods. This indicates that alpha increases were significantly higher in the reference period compared to the test period. There was no difference between stimulation groups. (d) Analysis of IAF across tasks and blocks. The IAF remained stable in resting blocks, whereas in rotation tasks, it was significantly elevated compared to resting blocks and showed a significant slowing from baseline to the poststimulation block.

(a) - Test Period:  $\triangle$ Alpha<sub>Rotation</sub> vs  $\triangle$ ERD

# (b) - Ref. Period: $\Delta Alpha_{Rotation}$ vs $\Delta ERD$

interaction effect *Period*×*Stimulation* [F(1,45)=1.76, p=.183,  $\eta_p^2=0.073$ ,  $\eta_G^2=0.031$ ] reached significance. This suggests that although the reference period appears to be the primary factor influencing ERD changes, attributing the stimulation effect exclusively to either the reference or test period is overly simplistic. It likely indicates a complex interplay between stimulation effects and alpha activity alterations across both periods.

Further, we examined the changes in IAF over the course of the experiment aiming to assess the stability of IAF and to post-hoc verify the accuracy of the stimulation frequency. IAFs were extracted from the baseline blocks  $(\text{Rest}_{\text{Baseline}}, \text{ Rotation}_{\text{Baseline}})$  and the poststimulation blocks (Rest<sub>Poststim 3</sub>, Rotation<sub>Poststim</sub>). Descriptive analysis revealed varying IAF frequencies  $[M_{\text{RestBaseline}} = 10.2 \text{ Hz}]$ ,  $SD_{\text{RestBaseline}} = 0.91 \text{ Hz}, M_{\text{RestPoststim}} = 10.1 \text{ Hz}, SD_{\text{RestPoststim}} =$  $0.99 \,\mathrm{Hz}, M_{\mathrm{RotationBaseline}} = 10.9 \,\mathrm{Hz}, SD_{\mathrm{RotationBaseline}} = 1.38 \,\mathrm{Hz},$  $M_{\text{RotationPoststim}} = 10.4 \,\text{Hz}, \quad SD_{\text{RotationPoststim}} = 1.14 \,\text{Hz}].$  To analyze these differences, we employed an rmANOVA with within-subject factors of *Block* (Baseline, Poststim) and Task (Resting, Rotation). This revealed a significant main effect of *Block* on IAF [F(1,47)=17.71, $p < .001, \eta_p^2 = 0.274, \eta_G^2 = 0.019$ ], attributed to a slowing of IAF from baseline to poststim blocks (see Figure 4d). Additionally, a significant main effect of Task was observed [F(1,47) = 18.08, p < .001,  $\eta_p^2 = 0.278$ ,  $\eta_G^2 = 0.044$ ], indicating a faster IAF during the mental rotation task compared to the resting block. There was no significant interaction of  $Block \times Task$  [F(1,47)=1.92, p=.172,  $\eta_{\rm p}^2 = 0.039, \eta_{\rm G}^2 = 0.008$ ].

## 4 | DISCUSSION

To this day, studies researching in vivo effects of tTIS are still sparse, which is true for murine models but especially in humans. Our current study contributes a proofof-concept and new evidence for the neuromodulatory effect of tTIS in humans. We demonstrate an outlasting electrophysiological effect of tTIS in the form of an increase of ERD in the alpha range during a mental rotation task. Building on prior work that demonstrated the neuromodulatory potential of tACS on parieto-occipital alpha activity (Kasten et al., 2018; Kasten & Herrmann, 2017), we extend these findings to tTIS. Though contrary to our hypothesis, we observed no stimulation effect on mental rotation performance or resting alpha activity.

To further clarify the specific impact of the stimulation on ERD, we have conducted an additional analysis aimed at determining whether the stimulation effect could be attributed distinctly to alpha changes either in the reference or the test period or a combination of both (as discussed in Kasten & Herrmann, 2017). This analysis uncovered PSYCHOPHYSIOLOGY SPR

a significantly larger change in alpha power during the reference period than in the test period. Though importantly, we observed no interaction effect between the period-specific changes in alpha activity and the stimulation applied, which stands in contrast to the distinct interaction effect we identified between ERD changes and the stimulation. This suggests that while the alterations in ERD are primarily attributable to the reference period, these alterations alone cannot fully account for the effects of the stimulation. One possible explanation may be that the stimulation effect does not originate from changes in either period individually, but rather from an interaction between changes occurring in both periods.

Even though no specific stimulation effect was evident in the reference period, our correlational analysis suggests that the significant increase in alpha activity during this period might be the primary influence on the overall ERD changes observed. This elevated alpha activity sets the stage for a more marked drop in alpha activity when the stimulus is presented, suggesting that the dynamics within the reference period play a crucial role in shaping the brain's oscillatory response to stimuli. Generally speaking, the ERD is well known to be associated with task performance (Haegens et al., 2011; Kasten & Herrmann, 2017; Klimesch et al., 2003; Neubauer et al., 1995). Klimesch et al. (2003) used repetitive transcranial magnetic stimulation (rTMS) to apply stimulation time-locked to the reference period, in turn boosting alpha activity to result in an increase in ERD and task performance. Though they did not measure ERD, studies by Zoefel et al. (2011) and Hanslmayr et al. (2005) used neurofeedback training to demonstrate a performance advantage of increased parieto-occipital alpha activity on a mental rotation task. The rationale is based on the presumption that alpha range oscillations play a crucial role in regulating information flow. Specifically, high alpha oscillations represent the natural resting or idling state of certain brain areas. However, when a region becomes relevant to a task, alpha oscillations decrease, signaling that the area is transitioning to a state of active information processing, with the change in alpha activity quantified by ERD.

However, in our study, we did not observe an improvement in behavioral performance despite the increased ERD. One potential explanation for this discrepancy is the presence of a behavioral ceiling effect. This is particularly evident in task accuracy, where all groups achieved near-perfect hit rates in the poststim block. This indicates that participants, including those in the sham group, performed the task with ease, leaving little room for the potential enhancing effect of the stimulation to manifest in improved performance. Thus, the comparable improvement in task performance across all three group rather indicates a significant learning effect,

covering any specific effects of stimulation. To detect the specific benefits of stimulation on mental rotation, future studies should consider increasing the task difficulty. Interestingly, our findings, which indicate no significant effect of stimulation on reaction times, are consistent with existing literature that also did not observe a performance improvement following an increase in alpha activity (Klimesch et al., 2003).

A potential limitation in measuring stimulation effects during the mental rotation task is the experimental design, which may have led to an underestimation of the true stimulation effects. Specifically, our experimental design positioned the Rest<sub>Poststim 2</sub> Block in between the Rotation<sub>Stimulation</sub> and Rotation<sub>Poststimulation</sub> Blocks, resulting in a 5-minute delay between applied stimulation and measuring its effect on mental rotation. We cannot rule out that the stimulation effect is only strongest immediately after stimulation; thus, this delay could theoretically mean that the peak stimulation impact was not fully present during the Rotation<sub>Poststimulation</sub> block. Future studies should prioritize examining stimulation effects on mental rotation tasks by immediately following stimulation with a task to accurately assess the outcomes.

Contrary to our initial hypothesis, we did not observe any stimulation effect on alpha activity during the vigilance task. Though the alpha activity experienced an increase between resting blocks, the rate of this increase was consistent across all stimulation groups, indicating that the tTIS and tACS groups did not exhibit greater increases in alpha activity compared to the sham group. This suggests that the observed increase in alpha power over time might be attributed to an increase in mental fatigue throughout the experiment, which is known to increase alpha activity (Hsu & Wang, 2013; Käthner et al., 2014; Trejo et al., 2015), rather than a direct effect of the stimulation. In relation to tACS, our results did not align with the expected increase in alpha activity following parieto-occipital tACS, a well-documented effect in existing literature (Kasten et al., 2016; Kasten & Herrmann, 2017; Neuling et al., 2013; Vossen et al., 2015; Zaehle et al., 2010). Several reasons could account for the absence of observed stimulation effects. One consideration is the electrode montage's efficacy in targeting alpha oscillation generators. Despite using an established tACS montage and verifying field strengths at the region of interest (ROI) with SimNIBS simulations, the possibility of a non-optimal montage cannot be entirely dismissed. Furthermore, uncontrolled intra- and interindividual variations, including suboptimal brain states at the time of stimulation (Bergmann, 2018), could have influenced our results.

However, these potential explanations do not fully account for the discrepancy for a lack of stimulation effect on resting alpha activity and our observed stimulation effect on ERD or task-related alpha activity. It is possible that the alternation between resting and rotation blocks introduced significant interference, given the rotation task's requirement for constant adjustments in alpha activity. This hypothesis is supported by similar findings from Kasten and Herrmann (2017), who employed a similar design featuring alternating rest and task phases and were not able to demonstrate changes in resting alpha activity but were able to observe alterations in task-related alpha activity. Still, our study leaves open questions regarding the precise relationship between stimulation and alpha activity, underlining the necessity for further research to disentangle these complex dynamics.

While computational modeling studies have shown promising outcomes, indicating that tTIS can lead to neuronal entrainment (Karimi et al., 2019; Lee, 2021; Su et al., 2021; Zhu et al., 2022), some research highlights the potential for enhancing its effectiveness. These studies specifically propose enhancing tTIS by adopting an electrode configuration that utilizes multiple pairs of electrodes. This approach aims to increase the intensity at the target site and improve focality, potentially amplifying stimulation efficacy (Cao & Grover, 2019; Howell & McIntyre, 2021; Huang et al., 2020; Huang & Parra, 2019; Zhu et al., 2019). To validate these theoretical improvements, future research should investigate these strategies through in vivo studies.

A potential limitation of this study is the observed lack of robustness of the IAF. The aim of this study was to apply stimulation at each subject's IAF, as the Arnold tongue principle suggests that the efficacy of oscillatory stimulation can be enhanced when the target frequency and the applied frequency are as close as possible (Huang et al., 2021; Kasten et al., 2019; Schutter & Wischnewski, 2016). However, our exploratory analysis revealed that IAF fluctuates over the course of the experiment, a finding that aligns with other studies (Benwell et al., 2019; Stecher et al., 2017; Vossen et al., 2015). In this study, we estimated the IAF based on resting-state activity in the initial baseline resting block. Hence, the optimal stimulation frequency might have shifted by the time stimulation began. Furthermore, these studies suggest that IAF can vary depending on whether the neuronal system is at rest or under load. Accordingly, we might have misestimated the IAF for the mental rotation task, which led to the application of non-optimal stimulation frequencies. Future studies should consider (1) extracting the IAF immediately before the application of stimulation and (2) using a task that closely resembles the cognitive

demands of the target task, to reduce the mismatch between stimulation and target frequency and thus enhance stimulation efficacy. Additionally, adopting closed-loop methods (Karabanov et al., 2016) that adjust stimulation parameters in real-time based on the current brain state (Bergmann, 2018) could offer a more sophisticated approach to optimizing stimulation effectiveness.

In our study, we employed an inactive control stimulation (sham stimulation which was applied for only 30 seconds), which allowed us to differentiate between stimulation effects and effects not based on electrostimulation. Some other tTIS studies (Conta et al., 2022; Wessel et al., 2023) instead opt for an active control condition, where a high-frequency condition without a frequency offset  $(\Delta f = f_1 - f_2 = 2000 \text{ Hz} - 2000 \text{ Hz} = 0 \text{ Hz})$  is utilized, allowing to measure stimulation effects specific to the amplitude modulation and potential confounding effects due to the high-frequency signal of the stimulation. Indeed, recent studies are debating a potential effect of the high-frequency signal in the form of a conduction block in off-target areas, which should be considered (Mirzakhalili et al., 2020; Wang et al., 2023), though this is mainly when using highintensity, suprathreshold tTIS. However, an active control is not necessary to differentiate between entrainment effects and non-specific stimulation effects. We hypothesized that true entrainment would specifically boost power at the frequency of amplitude modulation (IAF), without affecting neighboring frequencies. In contrast, a non-specific stimulation effect would likely cause a broad increase across the frequency spectrum, raising overall power. Our results, as visualized in Figure 3, confirm that the power increase is indeed localized to the IAF and thus frequencyspecific, as per the entrainment principle. Nonetheless, future studies should ideally incorporate both an active as well as an inactive control condition to measure and control stimulation-unrelated effects (sham stimulation), non-specific stimulation effects due to the high-frequency signal (active control) as well as true entrainment effects due to the amplitude modulation (verum stimulation).

## 5 | CONCLUSION

This study offers early electrophysiological evidence for tTIS's in vivo effect on humans. We observed increased ERD during a mental rotation task after both tTIS and tACS stimulation, but no increase in alpha power at rest. This indicates that manifestation of tTIS's effects requires an active task-engaged neuronal network. Although higher ERD has been associated with enhanced information processing in prior studies, we did not see this translate to improved performance in the mental rotation task, likely due to a ceiling effect. Future research should focus

PSYCHOPHYSIOLOGY SPR

on identifying the best stimulation parameters and brain states for modulating behavior through tACS and tTIS.

## AUTHOR CONTRIBUTIONS

**Carsten Thiele:** Conceptualization; data curation; formal analysis; investigation; methodology; software; visualization; writing – original draft. **Katharina S. Rufener:** Conceptualization; writing – review and editing. **Stefan Repplinger:** Software; writing – review and editing. **Tino Zaehle:** Project administration; resources; writing – review and editing. **Philipp Ruhnau:** Conceptualization; funding acquisition; methodology; project administration; resources; supervision; writing – review and editing.

### ACKNOWLEDGMENTS

We thank all members of the Zaehle Lab for their valuable contributions to discussions. We would furthermore like to thank two anonymous reviewers for their helpful comments on an earlier version of this document.

## FUNDING INFORMATION

This study was supported by the federal state of Saxony-Anhalt and the European Regional Development Fund (ERDF) in the Center for Behavioral Brain Sciences (CBBS, ZS/2016/04/78113). The funding source had no direct involvement in this study.

## CONFLICT OF INTEREST STATEMENT

The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

## DATA AVAILABILITY STATEMENT

The data sets generated and analyzed during the current study are available from the corresponding author upon reasonable request.

## DECLARATION OF AI AND AI-ASSISTED TECHNOLOGIES IN THE WRITING PROCESS

During the preparation of this work, the authors used ChatGPT-4 in order to improve readability and language of the manuscript. After using this tool, the authors reviewed and edited the content as needed and take full responsibility for the content of the publication.

## ORCID

Philipp Ruhnau D https://orcid.org/0000-0001-6546-7312

## REFERENCES

Abd Hamid, A. I., Gall, C., Speck, O., Antal, A., & Sabel, B. A. (2015). Effects of alternating current stimulation on the healthy and diseased brain. *Frontiers in Neuroscience*, *9*, 391.

## PSYCHOPHYSIOLOGY SPR

- Alexander, M. L., Alagapan, S., Lugo, C. E., Mellin, J. M., Lustenberger, C., Rubinow, D. R., & Fröhlich, F. (2019). Double-blind, randomized pilot clinical trial targeting alpha oscillations with transcranial alternating current stimulation (tACS) for the treatment of major depressive disorder (MDD). *Translational Psychiatry*, 9(1), 1–12.
- Antal, A., Ambrus, G. G., & Chaieb, L. (2014). The impact of electrical stimulation techniques on behavior. Wiley Interdisciplinary Reviews: Cognitive Science, 5(6), 649–659.
- Beatti, A., Rayner, A., Chipchase, L., & Souvlis, T. (2011). Penetration and spread of interferential current in cutaneous, subcutaneous and muscle tissues. *Physiotherapy*, 97(4), 319–326. https:// doi.org/10.1016/j.physio.2011.01.008
- Benwell, C. S. Y., London, R. E., Tagliabue, C. F., Veniero, D., Gross, J., Keitel, C., & Thut, G. (2019). Frequency and power of human alpha oscillations drift systematically with time-on-task. *NeuroImage*, 192, 101–114.
- Berger, A., Pixa, N. H., Steinberg, F., & Doppelmayr, M. (2018). Brain oscillatory and hemodynamic activity in a bimanual coordination task following transcranial alternating current stimulation (tACS): A combined EEG-fNIRS study. *Frontiers in Behavioral Neuroscience*, 12, 67.
- Bergmann, T. O. (2018). Brain state-dependent brain stimulation. *Frontiers in Psychology*, *9*, 2108.
- Brown, C. C. (1975). Electroanesthesia and electrosleep. *The American Psychologist*, *30*(3), 402–410.
- Brunelin, J., Adam, O., & Mondino, M. (2022). Recent advances in noninvasive brain stimulation for schizophrenia. *Current Opinion in Psychiatry*, 35(5), 338–344.
- Brunoni, A. R., Amadera, J., Berbel, B., Volz, M. S., Rizzerio, B. G., & Fregni, F. (2011). A systematic review on reporting and assessment of adverse effects associated with transcranial direct current stimulation. *International Journal of Neuropsychopharmacology*, 14(8), 1133–1145.
- Brunoni, A. R., Valiengo, L., Baccaro, A., Zanão, T. A., de Oliveira, J. F., Goulart, A., Boggio, P. S., Lotufo, P. A., Benseñor, I. M., & Fregni, F. (2013). The sertraline vs. electrical current therapy for treating depression clinical study: Results from a factorial, randomized, controlled trial. *JAMA Psychiatry*, 70(4), 383–391.
- Cao, J., & Grover, P. (2019). STIMULUS: Noninvasive dynamic patterns of neurostimulation using Spatio-temporal interference. *IEEE Transactions on Biomedical Engineering*, 67(3), 726–737.
- Cao, J., Doiron, B., Goswami, C., & Grover, P. (2020). The mechanics of temporal interference stimulation. *bioRxiv*. https://doi.org/ 10.1101/2020.04.23.051870
- Cassarà, A. M., Newton, T. H., Zhuang, K., Regel, S. J., Achermann, P., Kuster, N., & Neufeld, E. (2022). Safety recommendations for temporal interference stimulation in the brain. *bioRxiv*. https:// doi.org/10.1101/2022.12.15.520077
- Chen, X., Bin, G., Daly, I., & Gao, X. (2013). Event-related desynchronization (ERD) in the alpha band during a hand mental rotation task. *Neuroscience Letters*, *541*, 238–242.
- Cho, H., Razza, L. B., Borrione, L., Bikson, M., Charvet, L., Dennis-Tiwary, T. A., Brunoni, A. R., & Sudbrack-Oliveira, P. (2022). Transcranial electrical stimulation for psychiatric disorders in adults: A primer. *Focus (American Psychiatric Publishing)*, 20(1), 19–31.
- Conta, J. v., Kasten, F. H., Schellhorn, K., Ćurčić-Blake, B., Aleman, A., & Herrmann, C. S. (2022). Benchmarking the effects of

transcranial temporal interference stimulation (tTIS) in humans. *Cortex*, *154*, 299–310.

- Conta, J. v., Kasten, F. H., Ćurčić-Blake, B., Aleman, A., Thielscher, A., & Herrmann, C. S. (2021). Interindividual variability of electric fields during transcranial temporal interference stimulation (tTIS). *Scientific Reports*, 11(1), 20357.
- Donoghue, T., Haller, M., Peterson, E. J., Varma, P., Sebastian, P., Gao, R., Noto, T., Lara, A. H., Wallis, J. D., Knight, R. T., Shestyuk, A., & Voytek, B. (2020). Parameterizing neural power spectra into periodic and aperiodic components. *Nature Neuroscience*, 23(12), 1655–1665.
- Doppelmayr, M., Klimesch, W., Stadler, W., Pöllhuber, D., & Heine, C. (2002). EEG alpha power and intelligence. *Intelligence*, *30*(3), 289–302.
- Esmaeilpour, Z., Kronberg, G., Reato, D., Parra, L. C., & Bikson, M. (2021). Temporal interference stimulation targets deep brain regions by modulating neural oscillations. *Brain Stimulation*, 14(1), 55–65.
- Faul, F., Erdfelder, E., Lang, A.-G., & Buchner, A. (2007). G\*power 3: A flexible statistical power analysis program for the social, behavioral, and biomedical sciences. *Behavior Research Methods*, 39(2), 175–191.
- Fertonani, A., Ferrari, C., & Miniussi, C. (2015). What do you feel if I apply transcranial electric stimulation? Safety, sensations and secondary induced effects. *Clinical Neurophysiology*, 126(11), 2181–2188.
- Foxe, J. J., & Snyder, A. C. (2011). The role of alpha-band brain oscillations as a sensory suppression mechanism during selective attention. *Frontiers in Psychology*, 2, 154.
- Gabriel, S., Lau, R. W., & Gabriel, C. (1996). The dielectric properties of biological tissues: II. Measurements in the frequency range 10 Hz to 20 GHz. *Physics in Medicine & Biology*, 41(11), 2251–2269.
- Ganis, G., & Kievit, R. (2015). A new set of three-dimensional shapes for investigating mental rotation processes: Validation data and stimulus set. *Journal of Open Psychology Data*, 3, 1–7. https:// doi.org/10.5334/jopd.ai
- Goats, G. C. (1990). Interferential current therapy. British Journal of Sports Medicine, 24(2), 87–92. https://doi.org/10.1136/bjsm. 24.2.87
- Grossman, N., Bono, D., Dedic, N., Kodandaramaiah, S. B., Rudenko, A., Suk, H.-J., Cassara, A. M., Neufeld, E., Kuster, N., & Tsai, L.-H. (2017). Noninvasive deep brain stimulation via temporally interfering electric fields. *Cell*, *169*(6), 1029–1041. e16.
- Grossman, N., Okun, M. S., & Boyden, E. S. (2018). Translating temporal interference brain stimulation to treat neurological and psychiatric conditions. *JAMA Neurology*, 75(11), 1307–1308.
- Grover, S., Wen, W., Viswanathan, V., Gill, C. T., & Reinhart, R. M. G. (2022). Long-lasting, dissociable improvements in working memory and long-term memory in older adults with repetitive neuromodulation. *Nature Neuroscience*, 25(9), 1237–1246.
- Guleyupoglu, B., Schestatsky, P., Edwards, D., Fregni, F., & Bikson, M. (2013). Classification of methods in transcranial electrical stimulation (tES) and evolving strategy from historical approaches to contemporary innovations. *Journal of Neuroscience Methods*, 219(2), 297–311.
- Haegens, S., Nácher, V., Luna, R., Romo, R., & Jensen, O. (2011). α-Oscillations in the monkey sensorimotor network influence discrimination performance by rhythmical inhibition of neuronal

spiking. Proceedings of the National Academy of Sciences of the United States of America, 108(48), 19377–19382.

- Hanslmayr, S., Sauseng, P., Doppelmayr, M., Schabus, M., & Klimesch, W. (2005). Increasing individual upper alpha power by neurofeedback improves cognitive performance in human subjects. *Applied Psychophysiology and Biofeedback*, 30(1), 1–10.
- Herrington, T. M., Cheng, J. J., & Eskandar, E. N. (2016). Mechanisms of deep brain stimulation. *Journal of Neurophysiology*, 115(1), 19–38.
- Herrmann, C. S., & Strüber, D. (2017). What can transcranial alternating current stimulation tell us about brain oscillations? *Current Behavioral Neuroscience Reports*, 4(2), 128–137.
- Herrmann, C. S., Strüber, D., Helfrich, R. F., & Engel, A. K. (2016). EEG oscillations: From correlation to causality. *International Journal of Psychophysiology*, 103, 12–21.
- Howell, B., & McIntyre, C. C. (2021). Feasibility of interferential and pulsed transcranial electrical stimulation for neuromodulation at the human scale. *Neuromodulation*, *24*(5), 843–853.
- Hoy, K. E., Bailey, N., Arnold, S., Windsor, K., John, J., Daskalakis, Z. J., & Fitzgerald, P. B. (2015). The effect of γ-tACS on working memory performance in healthy controls. *Brain and Cognition*, 101, 51–56.
- Hsu, B.-W., & Wang, M.-J. J. (2013). Evaluating the effectiveness of using electroencephalogram power indices to measure visual fatigue. *Perceptual and Motor Skills*, 116(1), 235–252.
- Hsu, G., Farahani, F., & Parra, L. C. (2021). Cutaneous sensation of electrical stimulation waveforms. *Brain Stimulation*, *14*(3), 693–702.
- Huang, W. A., Stitt, I. M., Negahbani, E., Passey, D. J., Ahn, S., Davey, M., Dannhauer, M., Doan, T. T., Hoover, A. C., Peterchev, A. V., Radtke-Schuller, S., & Fröhlich, F. (2021). Transcranial alternating current stimulation entrains alpha oscillations by preferential phase synchronization of fast-spiking cortical neurons to stimulation waveform. *Nature Communications*, *12*(1), 3151.
- Huang, Y., & Parra, L. C. (2019). Can transcranial electric stimulation with multiple electrodes reach deep targets? *Brain Stimulation*, *12*(1), 30–40.
- Huang, Y., Datta, A., & Parra, L. C. (2020). Optimization of interferential stimulation of the human brain with electrode arrays. *Journal of Neural Engineering*, 17(3), 36023.
- Hutcheon, B., & Yarom, Y. (2000). Resonance, oscillation and the intrinsic frequency preferences of neurons. *Trends in Neurosciences*, 23(5), 216–222.
- Iszak, K., Gronemann, S. M., Meyer, S., Hunold, A., Zschüntzsch, J., Bähr, M., Paulus, W., & Antal, A. (2023). Why temporal inference stimulation may fail in the human brain: A pilot research study. *Biomedicine*, 11(7), 1813.
- JASP Team. (2023). JASP (version 0.17) [Computer software].
- Jensen, O., & Mazaheri, A. (2010). Shaping functional architecture by oscillatory alpha activity: Gating by inhibition. *Frontiers in Human Neuroscience*, 4, 186.
- Karabanov, A., Thielscher, A., & Siebner, H. R. (2016). Transcranial brain stimulation: Closing the loop between brain and stimulation. *Current Opinion in Neurology*, 29(4), 397–404.
- Karimi, F., Attarpour, A., Amirfattahi, R., & Nezhad, A. Z. (2019). Computational analysis of non-invasive deep brain stimulation based on interfering electric fields. *Physics in Medicine and Biology*, 64(23), 235010.
- Kasten, F. H., & Herrmann, C. S. (2017). Transcranial alternating current stimulation (tACS) enhances mental rotation

performance during and after stimulation. *Frontiers in Human Neuroscience*, *11*, 2.

PSYCHOPHYSIOLOGY SPR

- Kasten, F. H., Dowsett, J., & Herrmann, C. S. (2016). Sustained aftereffect of α-tACS lasts up to 70 min after stimulation. *Frontiers in Human Neuroscience*, *10*, 245.
- Kasten, F. H., Duecker, K., Maack, M. C., Meiser, A., & Herrmann, C. S. (2019). Integrating electric field modeling and neuroimaging to explain inter-individual variability of tACS effects. *Nature Communications*, 10(1), 5427.
- Kasten, F. H., Maess, B., & Herrmann, C. S. (2018). Facilitated eventrelated power modulations during transcranial alternating current stimulation (tACS) revealed by concurrent tACS-MEG. *eNeuro*, 5(3), 1–15.
- Khadka, N., & Bikson, M. (2023). Noninvasive electrical brain stimulation of the central nervous system. In N. V. Thakor (Ed.), *Handbook of Neuroengineering* (pp. 2101–2133). Springer Nature. https://doi.org/10.1007/978-981-16-5540-1\_59
- Khan, A., Yuan, K., Bao, S.-C., Ti, C. H. E., Tariq, A., Anjum, N., & Tong, R. K.-Y. (2022). Can transcranial electrical stimulation facilitate post-stroke cognitive rehabilitation? A systematic review and meta-analysis. *Frontiers in Rehabilitation Sciences*, 3, 795737.
- Khatoun, A., Asamoah, B., & Mc, L. M. (2021). A computational modeling study to investigate the use of Epicranial electrodes to deliver interferential stimulation to subcortical regions. *Frontiers in Neuroscience*, *15*, 779271.
- Kleiner, M., Brainard, D., & Pelli, D. (2007). What's new in *Psychtoolbox-3?*.
- Klimesch, W., Fellinger, R., & Freunberger, R. (2011). Alpha oscillations and early stages of visual encoding. *Frontiers in Psychology*, *2*, 118.
- Klimesch, W., Sauseng, P., & Gerloff, C. (2003). Enhancing cognitive performance with repetitive transcranial magnetic stimulation at human individual alpha frequency. *The European Journal of Neuroscience*, 17(5), 1129–1133.
- Klimesch, W., Sauseng, P., & Hanslmayr, S. (2007). EEG alpha oscillations: The inhibition-timing hypothesis. *Brain Research Reviews*, 53(1), 63–88.
- Käthner, I., Wriessnegger, S. C., Müller-Putz, G. R., Kübler, A., & Halder, S. (2014). Effects of mental workload and fatigue on the P300, alpha and theta band power during operation of an ERP (P300) brain-computer interface. *Biological Psychology*, 102, 118–129.
- Lee, S. (2021). Temporal Interference Stimulation for Focused Stimulation of Deep Structures of Human Brain: A Computer-Based Feasibility Study: Hanyang University.
- Limousin, P., & Foltynie, T. (2019). Long-term outcomes of deep brain stimulation in Parkinson disease. *Nature Reviews Neurology*, 15(4), 234–242.
- Ma, R., Xia, X., Zhang, W., Lu, Z., Wu, Q., Cui, J., Song, H., Fan, C., Chen, X., & Zha, R. (2021). High gamma and Beta temporal interference stimulation in the human motor cortex improves motor functions. *Frontiers in Neuroscience*, 15, 800436.
- Makeig, S., Debener, S., Onton, J., & Delorme, A. (2004). Mining event-related brain dynamics. *Trends in Cognitive Sciences*, 8(5), 204–210.
- Marchesotti, S., Nicolle, J., Merlet, I., Arnal, L. H., Donoghue, J. P., & Giraud, A.-L. (2020). Selective enhancement of low-gamma activity by tACS improves phonemic processing and reading accuracy in dyslexia. *PLoS Biology*, 18(9), e3000833.

- Miranda, P. C., Mekonnen, A., Salvador, R., & Ruffini, G. (2013). The electric field in the cortex during transcranial current stimulation. *NeuroImage*, 70, 48–58.
- Mirzakhalili, E., Barra, B., Capogrosso, M., & Lempka, S. F. (2020). Biophysics of temporal interference stimulation. *Cell Systems*, 11(6), 557–572.e5.
- Nakayashiki, K., Saeki, M., Takata, Y., Hayashi, Y., & Kondo, T. (2014). Modulation of event-related desynchronization during kinematic and kinetic hand movements. *Journal of Neuroengineering and Rehabilitation*, 11(1), 90.
- Neubauer, A., Freudenthaler, H. H., & Pfurtscheller, G. (1995). Intelligence and spatiotemporal patterns of event-related desynchronization (ERD). *Intelligence*, 20(3), 249–266.
- Neuling, T., Rach, S., & Herrmann, C. S. (2013). Orchestrating neuronal networks: Sustained after-effects of transcranial alternating current stimulation depend upon brain states. *Frontiers in Human Neuroscience*, 7, 161.
- Oostenveld, R., Fries, P., Maris, E., & Schoffelen, J.-M. (2011). FieldTrip: Open source software for advanced analysis of MEG, EEG, and invasive electrophysiological data. *Computational Intelligence and Neuroscience*, 2011, 1–9.
- Opitz, A., Falchier, A., Yan, C.-G., Yeagle, E. M., Linn, G. S., Megevand, P., Thielscher, A., Deborah, A. R., Milham, M. P., Mehta, A. D., & Schroeder, C. E. (2016). Spatiotemporal structure of intracranial electric fields induced by transcranial electric stimulation in humans and nonhuman primates. *Scientific Reports*, 6(1), 31236.
- Pahor, A., & Jaušovec, N. (2018). The effects of theta and gamma tACS on working memory and electrophysiology. *Frontiers in Human Neuroscience*, 11, 651.
- Paulus, W., Nitsche, M. A., & Antal, A. (2016). Application of transcranial electric stimulation (tDCS, tACS, tRNS): From motorevoked potentials towards modulation of behaviour. *European Psychologist*, 21(1), 4–14.
- Pfurtscheller, G. (1997). EEG event-related desynchronization (ERD) and synchronization (ERS). *Electroencephalography and Clinical Neurophysiology*, 1(103), 26.
- Pfurtscheller, G., & Aranibar, A. (1977). Event-related cortical desynchronization detected by power measurements of scalp EEG. *Electroencephalography and Clinical Neurophysiology*, 42(6), 817–826.
- Pfurtscheller, G., & Da Silva, F. L. (1999). Event-related EEG/MEG synchronization and desynchronization: Basic principles. *Clinical Neurophysiology*, 110(11), 1842–1857.
- Piao, Y., Ma, R., Weng, Y., Fan, C., Xia, X., Zhang, W., Zeng, G. Q., Wang, Y., Lu, Z., Cui, J., Wang, X., Gao, L., Qiu, B., & Zhang, X. (2022). Safety evaluation of employing temporal interference transcranial alternating current stimulation in human studies. *Brain Sciences*, 12(9), 1–10.
- Pinault, D. (2017). A neurophysiological perspective on a preventive treatment against schizophrenia using transcranial electric stimulation of the corticothalamic pathway. *Brain Sciences*, 7(4), 34.
- Pini, L., Pizzini, F. B., Boscolo-Galazzo, I., Ferrari, C., Galluzzi, S., Cotelli, M., Gobbi, E., Cattaneo, A., Cotelli, M. S., Geroldi, C., Zanetti, O., Corbetta, M., van den Heuvel, M., Frisoni, G. B., Manenti, R., & Pievani, M. (2022). Brain network modulation in Alzheimer's and frontotemporal dementia with transcranial electrical stimulation. *Neurobiology of Aging*, 111, 24–34.

- Rajji, T. K. (2019). Transcranial magnetic and electrical stimulation in Alzheimer's disease and mild cognitive impairment: A review of randomized controlled trials. *Clinical Pharmacology & Therapeutics*, 106(4), 776–780.
- Rampersad, S., Roig-Solvas, B., Yarossi, M., Kulkarni, P. P., Santarnecchi, E., Dorval, A. D., & Brooks, D. H. (2019). Prospects for transcranial temporal interference stimulation in humans: A computational study. *NeuroImage*, 202, 116124.
- Reato, D., Rahman, A., Bikson, M., & Parra, L. C. (2013). Effects of weak transcranial alternating current stimulation on brain activity—A review of known mechanisms from animal studies. *Frontiers in Human Neuroscience*, 7, 687.
- Reinhart, R. M. G., & Nguyen, J. A. (2019). Working memory revived in older adults by synchronizing rhythmic brain circuits. *Nature Neuroscience*, 22(5), 820–827.
- Riečanský, I., & Katina, S. (2010). Induced EEG alpha oscillations are related to mental rotation ability: The evidence for neural efficiency and serial processing. *Neuroscience Letters*, 482(2), 133–136.
- Rihs, T. A., Michel, C. M., & Thut, G. (2009). A bias for posterior alpha-band power suppression versus enhancement during shifting versus maintenance of spatial attention. *NeuroImage*, 44(1), 190–199.
- Romei, V., Gross, J., & Thut, G. (2012). Sounds reset rhythms of visual cortex and corresponding human visual perception. *Current Biology*, 22(9), 807–813.
- Rufener, K. S., & Zaehle, T. (2021). Dysfunctional auditory gamma oscillations in developmental dyslexia: A potential target for a tACSbased intervention. *Progress in Brain Research, 264*, 211–232.
- Rufener, K. S., Zaehle, T., Oechslin, M. S., & Meyer, M. (2016). 40Hz-transcranial alternating current stimulation (tACS) selectively modulates speech perception. *International Journal of Psychophysiology*, 101, 18–24.
- Sachkov, V. I., Liventsev, N. M., Kuzin, M. I., & Zhukovskii, V. D. (1967). Experiences with combined electronarcosis with interference currents in clinical surgery. In *FM Wageneder, electrotherapeutic sleep and electroanaesthesia*. Excerpta Medica.
- Salehinejad, M. A., Nejati, V., Mosayebi-Samani, M., Mohammadi, A., Wischnewski, M., Kuo, M.-F., Avenanti, A., Vicario, C. M., & Nitsche, M. A. (2020). Transcranial direct current stimulation in ADHD: A systematic review of efficacy, safety, and protocolinduced electrical field modeling results. *Neuroscience Bulletin*, 36(10), 1191–1212.
- Salehinejad, M. A., Wischnewski, M., Nejati, V., Vicario, C. M., & Nitsche, M. A. (2019). Transcranial direct current stimulation in attention-deficit hyperactivity disorder: A meta-analysis of neuropsychological deficits. *PLoS One*, 14(4), e0215095.
- Sauseng, P., Klimesch, W., Stadler, W., Schabus, M., Doppelmayr, M., Hanslmayr, S., Gruber, W. R., & Birbaumer, N. (2005). A shift of visual spatial attention is selectively associated with human EEG alpha activity. *The European Journal of Neuroscience*, 22(11), 2917–2926.
- Schuhmann, T., Kemmerer, S. K., Duecker, F., de Graaf, T. A., ten Oever, S., de Weerd, P., & Sack, A. T. (2019). Left parietal tACS at alpha frequency induces a shift of visuospatial attention. *PLoS One*, 14(11), e0217729.
- Schutter, D. J., & Wischnewski, M. (2016). A meta-analytic study of exogenous oscillatory electric potentials in neuroenhancement. *Neuropsychologia*, 86, 110–118.

- Shepard, R. N., & Metzler, J. (1971). Mental rotation of threedimensional objects. *Science*, *171*(3972), 701–703.
- Sherman, M. T., Kanai, R., Seth, A. K., & VanRullen, R. (2016). Rhythmic influence of top-down perceptual priors in the phase of prestimulus occipital alpha oscillations. *Journal of Cognitive Neuroscience*, 28(9), 1318–1330.
- Solomons, C. D., & Shanmugasundaram, V. (2019). A review of transcranial electrical stimulation methods in stroke rehabilitation. *Neurology India*, 67(2), 417–423.
- Song, X., Zhao, X., Zhou, Y., Liu, S., He, F., Liu, Y., & Ming, D. (2019). Typical electrode configuration analysis for temporally interfering deep brain stimulation. 2019 9th International IEEE/EMBS Conference on Neural Engineering (NER). https://doi.org/10. 1109/NER.2019.8717013
- Stecher, H. I., Pollok, T. M., Strüber, D., Sobotka, F., & Herrmann, C. S. (2017). Ten minutes of  $\alpha$ -tACS and ambient illumination independently modulate EEG  $\alpha$ -power. *Frontiers in Human Neuroscience*, *11*, 257.
- Stein, D. J., Fernandes Medeiros, L., Caumo, W., & Torres, I. L. S. (2020). Transcranial direct current stimulation in patients with anxiety: Current perspectives. *Neuropsychiatric Disease and Treatment*, 16, 161–169.
- Stoupis, D., & Samaras, T. (2022). Non-invasive stimulation with temporal interference: Optimization of the electric field deep in the brain with the use of a genetic algorithm. *Journal of Neural Engineering*, 19(5), 056018. https://doi.org/10.1088/1741-2552/ac89b3
- Su, X., Guo, J., Zhou, M., Chen, J., Li, L., Chen, Y., Sui, X., Li, H., & Chai, X. (2021). Computational modeling of spatially selective retinal stimulation with temporally interfering electric fields. *IEEE Transactions on Neural Systems and Rehabilitation Engineering*, 29, 418–428.
- Terasawa, Y., Tashiro, H., Ueno, T., & Ohta, J. (2022). Precise temporal control of interferential neural stimulation via phase modulation. *IEEE Transactions on Bio-Medical Engineering*, 69(1), 220–228.
- The jamovi project. (2022). jamovi (Version 2.3) [Computer Software]. Retrieved from https://www.jamovi.org
- Thiele, C., Zaehle, T., Haghikia, A., & Ruhnau, P. (2021). Amplitude modulated transcranial alternating current stimulation (AM-TACS) efficacy evaluation via phosphene induction. *Scientific Reports*, 11(1), 1–10.
- Thielscher, A., Antunes, A., & Saturnino, G. B. (2015). Field modeling for transcranial magnetic stimulation: A useful tool to understand the physiological effects of TMS? 37th Annual International Conference of the IEEE Engineering in Medicine and Biology Society (EMBC).
- Thut, G., Schyns, P. G., & Gross, J. (2011). Entrainment of perceptually relevant brain oscillations by non-invasive rhythmic stimulation of the human brain. *Frontiers in Psychology*, *2*, 170.
- Trejo, L. J., Kubitz, K., Rosipal, R., Kochavi, R. L., & Montgomery, L. D. (2015). EEG-based estimation and classification of mental fatigue. *Psychology*, 6(5), 572–589.
- Veniero, D., Vossen, A., Gross, J., & Thut, G. (2015). Lasting EEG/ MEG aftereffects of rhythmic transcranial brain stimulation: Level of control over oscillatory network activity. *Frontiers in Cellular Neuroscience*, 9, 477.
- Vogeti, S., Boetzel, C., & Herrmann, C. S. (2022). Entrainment and spike-timing dependent plasticity-a review of proposed

mechanisms of transcranial alternating current stimulation. *Frontiers in Systems Neuroscience*, *16*, 827353.

PSYCHOPHYSIOLOGY SPR

- Volkmann, J. (2004). Deep brain stimulation for the treatment of Parkinson's disease. *Journal of Clinical Neurophysiology*, 21(1), 6–17.
- Vossen, A., Gross, J., & Thut, G. (2015). Alpha power increase after transcranial alternating current stimulation at alpha frequency (α-tACS) reflects plastic changes rather than entrainment. *Brain Stimulation*, 8(3), 499–508.
- Vosskuhl, J., Strüber, D., & Herrmann, C. S. (2018). Non-invasive brain stimulation: A paradigm shift in understanding brain oscillations. *Frontiers in Human Neuroscience*, 12, 211.
- Wang, B., Aberra, A. S., Grill, W. M., & Peterchev, A. V. (2023). Responses of model cortical neurons to temporal interference stimulation and related transcranial alternating current stimulation modalities. *Journal of Neural Engineering*, 19(6), 66047.
- Wang, H., Wang, K., Xue, Q., Peng, M., Yin, L., Gu, X., Leng, H., Lu, J., Liu, H., Wang, D., Xiao, J., Sun, Z., Li, N., Dong, K., Zhang, Q., Zhan, S., Fan, C., Min, B., Zhou, A., ... Wang, Y. (2022). Transcranial alternating current stimulation for treating depression: A randomized controlled trial. *Brain*, *145*(1), 83–91.
- Wessel, M. J., Beanato, E., Popa, T., Windel, F., Vassiliadis, P., Menoud, P., Beliaeva, V., Violante, I. R., Abderrahmane, H., Dzialecka, P., Park, C.-H., Maceira-Elvira, P., Morishita, T., Cassara, A. M., Steiner, M., Grossman, N., Neufeld, E., & Hummel, F. C. (2023). Noninvasive theta-burst stimulation of the human striatum enhances striatal activity and motor skill learning. *Nature Neuroscience*, 26, 1–12.
- Wischnewski, M., Engelhardt, M., Salehinejad, M. A., Schutter, D. J. L. G., Kuo, M.-F., & Nitsche, M. A. (2019). NMDA receptormediated motor cortex plasticity after 20 Hz transcranial alternating current stimulation. *Cerebral Cortex*, 29(7), 2924–2931.
- Wischnewski, M., Schutter, D. J. L. G., & Nitsche, M. A. (2019). Effects of beta-tACS on corticospinal excitability: A metaanalysis. *Brain Stimulation*, 12(6), 1381–1389.
- Yang, D., Shin, Y.-I., & Hong, K.-S. (2021). Systemic review on transcranial electrical stimulation parameters and EEG/fNIRS features for brain diseases. *Frontiers in Neuroscience*, 15, 629323.
- Zaehle, T., Rach, S., & Herrmann, C. S. (2010). Transcranial alternating current stimulation enhances individual alpha activity in human EEG. *PLoS One*, *5*(11), e13766.
- Zaghi, S., Acar, M., Hultgren, B., Boggio, P. S., & Fregni, F. (2010). Noninvasive brain stimulation with low-intensity electrical currents: Putative mechanisms of action for direct and alternating current stimulation. *The Neuroscientist*, 16(3), 285–307.
- Zeng, F.-G., Tran, P., Richardson, M., Sun, S., & Xu, Y. (2019). Human sensation of transcranial electric stimulation. *Scientific Reports*, 9(1), 1–12.
- Zhu, X., Li, Y., Zheng, L., Shao, B., Liu, X., Li, C., Huang, Z.-G., Liu, T., & Wang, J. (2019). Multi-point temporal interference stimulation by using each electrode to carry different frequency currents. *IEEE Access*, 7, 168839–168848.
- Zhu, Z., Xiong, Y., Chen, Y., Jiang, Y., Qian, Z., Lu, J., Liu, Y., & Zhuang, J. (2022). Temporal interference (TI) stimulation boosts functional connectivity in human motor cortex: A comparison study with transcranial direct current stimulation (tDCS). *Neural Plasticity*, 2022, 7605046.

- Zoefel, B., Archer-Boyd, A., & Davis, M. H. (2018). Phase entrainment of brain oscillations causally modulates neural responses to intelligible speech. *Current Biology*, *28*(3), 401–408.e5.
- Zoefel, B., Huster, R. J., & Herrmann, C. S. (2011). Neurofeedback training of the upper alpha frequency band in EEG improves cognitive performance. *NeuroImage*, *54*(2), 1427–1431.
- Zumer, J. M., Scheeringa, R., Schoffelen, J.-M., Norris, D. G., & Jensen, O. (2014). Occipital alpha activity during stimulus processing gates the information flow to object-selective cortex. *PLoS Biology*, *12*(10), e1001965.

**How to cite this article:** Thiele, C., Rufener, K. S., Repplinger, S., Zaehle, T., & Ruhnau, P. (2024). Transcranial temporal interference stimulation (tTIS) influences event-related alpha activity during mental rotation. *Psychophysiology*, *00*, e14651. <u>https://doi.org/10.1111/psyp.14651</u>